


UNIVERSITY OF TORONTO



3 1761 01350188 7

Digitized by the Internet Archive  
in 2007 with funding from  
Microsoft Corporation











21

370 cr

370 Train

11



AN INTRODUCTION  
TO THE  
ALGEBRA OF QUANTICS

*EDWIN BAILEY ELLIOTT*

London

HENRY FROWDE

OXFORD UNIVERSITY PRESS WAREHOUSE  
AMEN CORNER, E.C.



New York

MACMILLAN & CO., 66 FIFTH AVENUE

465i

# AN INTRODUCTION

TO THE

# ALGEBRA OF QUANTICS

BY

EDWIN BAILEY ELLIOTT, M.A., F.R.S.

WAYNFLETE PROFESSOR OF PURE MATHEMATICS, AND  
FELLOW OF MAGDALEN COLLEGE, OXFORD

39381  
21/6/97

Oxford

AT THE CLARENDON PRESS

1895

QA  
201  
E44

Oxford

PRINTED AT THE CLARENDON PRESS

BY HORACE HART, PRINTER TO THE UNIVERSITY



## PREFACE

---

THE present work is an expansion of a course of lectures which I have annually delivered for some years past at Queen's College, Oxford.

Its primary object is, as was the case in the lecture room, that of explaining with all the clearness at my command the leading principles of invariant algebra, in the hope of making it evident to the junior student that the subject is attractive as well as important, and that its early difficulties are only such as he can readily surmount. Lucidity in mathematical works has often suffered from undue compression. My constant aim has been to guard against such a possibility here. In a book of moderate size dealing with a great subject much must remain unsaid, if the fundamental considerations are to be presented with the thoroughness and the perspicuity necessary to enable the student adequately to realize them, and give him the interest in them which will prepare him to pursue for himself the study to which they introduce him.

But, while the interests of the beginner have thus been given precedence, I am not without hope that the mathematician who is not new to the higher algebra will, especially in the chapters near the middle of the

book, find in its pages matter of value to him as an aid to his researches. In some branches of the theory, which though of really elementary character are of comparatively recent investigation, as for instance in much of the algebra of differential operators, it is believed that a welcome supplement to previous treatises is offered.

The title 'Algebra of Quantics' is perhaps one of my own introduction. It probably needs no defence, and can hardly fail to convey the right meaning. The mathematical world has now for half a century associated the algebra of invariants and covariants with the name of Cayley, and with his 'Memoirs on Quantics,' so that it may perhaps be regarded as appropriate that a new work, appearing in the year which has seen the close of the labours of the renowned author of those memoirs, and dealing with their subject, should bear a name which recalls his memory.

To Salmon's *Higher Algebra* and his other works it is impossible to say how much I am indebted, both for direct reference and for guidance to the use of other authorities. Faà de Bruno's *Formes Binaires* has also been constantly before me. Of Clebsch's *Binäre Formen* and Gordan's *Invariantentheorie* less use has been made, as their symbolical method, and their successful application of it to the great problem of the investigation of complete irreducible systems, have been reluctantly passed over with little more than an allusion in the following pages. A scanty chapter or two on this subject would have been utterly inadequate, and inconsistent with the general plan, as stated above, of an introductory treatise which prefers to omit rather

than to obscure by condensation. A whole work which shall present to the English reader in his own language a worthy exposition of the method of the great German masters remains a desideratum.

The reader will not, however, find that the present work is a compilation from others which have preceded it, great as has been the help which those others have afforded. Constant recourse has been had to the original authorities, particularly of course to Cayley's series of memoirs, and to Sylvester's writings in the Cambridge and Dublin Mathematical Journal, the American Journal of Mathematics, and elsewhere.

No bibliography of works and memoirs on the subject has been introduced. All mathematicians who wish to go deeply into the study of original authorities will have in their hands Dr. F. Meyer's 'Bericht über den gegenwärtigen Stand der Invariantentheorie' in the 'Jahresbericht der Deutschen Mathematiker-Vereinigung' for 1890-91, which is so full and thorough a bibliography and analysis of what has been done, especially in the later period of the history of the invariant theory, that it is hard to see how more can be desired. With regard to the originators of particular results, the difficulty continues, and has grown with the multitude of investigators, which was felt by Dr. Salmon when he wrote, 'I can scarcely pretend to assign to their proper authors the merits of the several steps; and, as between Messrs. Cayley and Sylvester, perhaps these gentlemen themselves, who were in constant communication with each other at the time, would now find it hard to say how much properly belongs to each.' To the difficulty with

regard to Cayley and Sylvester may in particular be added that of discriminating between what in Salmon's work should be ascribed to them or others at all and what to Salmon himself. Throughout the following pages discoverers' names are very frequently attached to results; but it is too much to hope, though all care has been taken, that there are not cases in which the names given are those of authors in whose writings the results in question have certainly occurred, rather than those of the authors who first gave them.

I am indebted to several friends for suggestions and other help. Among them there is one, Mr. J. Hammond, M.A., one of the most distinguished of living researchers in the higher Algebra, to whom my especial thanks are due for a manuscript on the binary quintic which has been exceedingly helpful.

Some students, approaching the subject for the first time, will be advised to omit chapters vii to xi till part of what follows them has been read.

E. B. ELLIOTT.

OXFORD,

*September, 1895.*

# TABLE OF CONTENTS

---

## CHAPTER I.

### PRINCIPLES AND DIRECT METHODS.

	PAGE
Definitions (Arts. 1-4) . . . . .	I
Invariants and covariants to be expected (5-7) . . . . .	6
Jacobians, Hessians, eliminants, and discriminants (8-13, 15). . . . .	10
The modulus irresoluble (14) . . . . .	19
A theorem on powers of the modulus proved and applied (16, 17) . . . . .	21
Intermediate invariants and covariants (18-20) . . . . .	27

## CHAPTER II.

### ESSENTIAL QUALITIES OF INVARIANTS.

Homogeneity (Arts. 21, 22) . . . . .	31
The factor a power of the modulus (23, 24) . . . . .	33
Isobarism (25-28) . . . . .	36
Absolute invariants (29) . . . . .	40
Number of independent invariants limited (30) . . . . .	41
Invariants of more quantics than one (31-33) . . . . .	43

## CHAPTER III.

### ESSENTIAL QUALITIES OF COVARIANTS.

Homogeneity (Arts. 34-36) . . . . .	46
The factor a power of the modulus (37) . . . . .	47
Isobarism (38-40) . . . . .	48
Absolute covariants (41) . . . . .	52
Number of independent covariants limited (42, 43) . . . . .	53
Covariants of more quantics than one (44) . . . . .	56
Invariants and covariants of covariants (45) . . . . .	57

## CHAPTER IV.

## COGREDIENT AND CONTRAGREDIENT QUANTITIES.

	PAGE
Cogredency of $\frac{d}{dy}$ and $-\frac{d}{dx}$ with $x$ and $y$ (Arts. 46-50) . . . . .	59
Cogredency defined (51) . . . . .	64
Emanants (52, 53) . . . . .	65
Geometry of binary systems (54, 55). . . . .	68
Covariants derived from emanants (56-58) . . . . .	71
Symbolical representation. Hyperdeterminants. Transvectants (59-61) . . . . .	75
Contragredency (62-65) . . . . .	82
Contravariants and mixed concomitants. Evectants (66, 67) . . . . .	85
Covariants of a binary quantic as contravariants and as invariants (68, 69) . . . . .	87

## CHAPTER V.

## BINARY QUANTICS. INVARIANTS, ETC. AS FUNCTIONS OF DIFFERENCES.

Convention as to numerical multiples (Art. 71) . . . . .	90
Covariancy of the factors of a binary quantic (72, 73) . . . . .	91
Invariants as functions of differences (74-76) . . . . .	93
Discriminants (77, 79) . . . . .	97
Invariants of a quadratic, of a cubic, and of a quartic (78, 80, 81) . . . . .	98
Covariants as functions of differences (82-84) . . . . .	104
Covariants of a quadratic, and of a cubic (85-88) . . . . .	108
Several binary quantics (89) . . . . .	111

## CHAPTER VI.

## BINARY QUANTICS CONTINUED. ANNIHILATORS. SEMINVARIANTS.

The annihilators $\Omega$ and $O$ of invariants (Arts. 90-94) . . . . .	112
Symmetry of invariants. Skew invariants (95, 96) . . . . .	117
Sufficiency of conditions of annihilation (97) . . . . .	118
Differential operations which effect a linear transformation (98-100) . . . . .	119
Formation of invariants by aid of $\Omega$ (101, 102) . . . . .	124
Invariants of several binary quantics (103) . . . . .	126
Annihilators of covariants (104-108) . . . . .	128
A covariant is given by one of its coefficients (109, 110) . . . . .	131
Seminvariants, and the covariants which they lead (111, 112, 115) . . . . .	134
Determination of seminvariants (113, 114) . . . . .	136
—	
Additional examples . . . . .	142

CHAPTER VII.

FURTHER THEORY OF THE OPERATORS  $\Omega$  AND  $O$ . RECIPROCITY.

	PAGE
Gradients, and the differential equations they satisfy (Arts. 116, 117) . . . . .	145
Seminvariants and invariants as classes of gradients (118-120) . . . . .	146
Repeated operations with $\Omega$ and $O$ . Alternants (121-125) . . . . .	149
There are no seminvariants of negative excess (126) . . . . .	154
The exact number of given positive excess (128, 129) . . . . .	156
Reciprocity. Hermite's law (130, 131) . . . . .	160
Gradients in more sets than one (132) . . . . .	163

CHAPTER VIII.

GENERATING FUNCTIONS.

Generating functions for numbers of partitions and of seminvariants (Arts. 133-137) . . . . .	165
Irreducible concomitants of linear form, of quadratic, and of cubic (138-141) . . . . .	171
Irreducible invariants of quartic, of quintic and of sextic (142, 143) . . . . .	176
Generating functions for concomitants of given degree and order (144) . . . . .	178
Reduced and representative generating functions (145-147) . . . . .	179
Real generating functions (148) . . . . .	186

CHAPTER IX.

HILBERT'S PROOF OF GORDAN'S THEOREM.

Lemmas on solutions of Diophantine equations (Art. 150) . . . . .	188
Proof of Gordan's theorem for invariants of one binary quantic (151-154) . . . . .	193
Proof for covariants of one binary quantic (155, 156) . . . . .	200
Proof for several binary quantics (157) . . . . .	201

CHAPTER X.

PROTOMORPHS, ETC.

Recapitulation (Art. 158) . . . . .	204
Substitution of derived quantics for arguments in seminvariants, &c. (159-161) . . . . .	205
Systems of protomorphs (162-168) . . . . .	209

Irreducible systems of concomitants. The cubic and quartic (169-171) . . . . .	218
Seminvariants are given by their non-unitary terms (172, 173) . . . . .	221
Non-unitary terms of invariants are determinate (174, 175) . . . . .	224
Seminvariants as quantics in $a_p : 1$ . (176-179) . . . . .	229
Operators which generate seminvariants (180-182) . . . . .	232

## CHAPTER XI.

FURTHER THEORY OF SEMINVARIANTS. THE BINARY  
QUANTIC OF INFINITE ORDER.

Expressions for $\Omega$ , &c. Roots and sums of their powers (Arts. 183-187) . . . . .	237
Non-unitary symmetric functions (188) . . . . .	245
Generating functions. Perpetuants (189, 190) . . . . .	247
Power ending products (191) . . . . .	250
Annihilators of all gradients (192, 193) . . . . .	251
Generators of all seminvariants (194) . . . . .	253
<hr/>	
Additional Examples (Miscellaneous) . . . . .	255

## CHAPTER XII.

## CANONICAL FORMS, ETC.

Definitions (Arts. 195-197) . . . . .	260
Canonical form of binary cubic (199-202) . . . . .	263
Canonical forms of binary quantics of odd order (203-205) . . . . .	268
Canonizants (206, 207) . . . . .	272
Quantics of even order. Catalecticants (208-210) . . . . .	275
Canonical form of binary quartic (211-220) . . . . .	277
Solution of a quartic equation (221, 222) . . . . .	289
Geometry of concomitants of quartic (223) . . . . .	291
Binary sextics, octavics, and $2n$ -ics (224-228) . . . . .	294
Canonical form of ternary cubic (229) . . . . .	300
Catalecticants of ternary, quaternary, &c. quartics (230, 231) . . . . .	301

## CHAPTER XIII.

## INVARIANTS AND COVARIANTS OF THE BINARY QUINTIC AND SEXTIC.

The twenty-three concomitants of a quintic (Arts. 232, 234, 235) . . . . .	304
Hammond's canonical and semi-canonical forms (233, 235) . . . . .	305
Discriminant of Quintic (236) . . . . .	311
Syzygy among invariants of Quintic (237) . . . . .	311
Hermite's Formes-types (238, 240) . . . . .	313



	PAGE
Quintics which satisfy invariant conditions (239-243) . . . . .	314
The twenty-six concomitants of a sextic (244, 245) . . . . .	322
<hr/>	
Additional examples . . . . .	325

## CHAPTER XIV.

## SEVERAL BINARY QUANTICS.

Concomitants of several binary quantics (Art. 247) . . . . .	328
Linear form and $p$ -ic (248, 249, 252) . . . . .	329
Cases of $p = 1$ and $p = 2$ (250, 251) . . . . .	331
Irreducible system for linear form and cubic (253, 254) . . . . .	333
Linear form and quartic (255) . . . . .	338
Several linear forms (256) . . . . .	339
System of two quadratics (257, 258) . . . . .	340
Linear form and two quadratics (259) . . . . .	345
Quadratic and cubic (260-262) . . . . .	346
System of quantics with orders in A. P. (263) . . . . .	349
Quantics of one order. Combinants (264, 265) . . . . .	351

## CHAPTER XV.

## RESTRICTED SUBSTITUTIONS. METRICAL GEOMETRY OF PENCILS.

The substitutions of Cartesian geometry (Art. 266) . . . . .	354
Boolean and orthogonal invariants, &c. (267-269) . . . . .	355
Orthogonal cogrediency and contragrediency identical (270) . . . . .	358
Annihilators of orthogonal and Boolean concomitants (271, 272) . . . . .	361
Boolean systems for quantics of low orders (273-278) . . . . .	364

## CHAPTER XVI.

## TERNARY QUANTICS. THE QUADRATIC AND CUBIC.

Triple arrangement of a ternary quantic (Art. 280) . . . . .	369
The annihilators of invariants and covariants (281-283, 288, 289) . . . . .	370
The alternants of these annihilators (284-287) . . . . .	374
The invariant of a ternary quadratic (290) . . . . .	380
The invariants of a ternary cubic (291-295) . . . . .	381
Covariants of ternary quantics (296-299) . . . . .	389
Has a ternary quadratic covariants? (300) . . . . .	398
Covariants of a ternary cubic (301-304) . . . . .	399
Contravariants (305) . . . . .	405
Contravariants of ternary quadratic and of ternary cubic (306-308) . . . . .	407
Mixed concomitants (309, 310) . . . . .	410
Quantics in more than three variables (311) . . . . .	412

#### ERRATA.

- P. 11, line 6, for covariant read invariant.
- P. 53, line 9, for invariant read covariant.
- P. 56, line 20, for  $p$ -ics read quantics.
- P. 58, line 2, for  $i' i$  read  $i' ip$ .
- P. 95, line 6, for  $x_r y_s - x y_r$  read  $x_r y_s - x_s y_r$ .
- P. 246, line 4 from bottom, for  $(a_0, a_1, a_2, \dots a_p)^p$  read  $(a_0, a_1, a_2, \dots a_p)$ .
- P. 281, line 8, for  $J27^2$  read  $27J^2$ .

AN INTRODUCTION  
TO  
THE ALGEBRA OF QUANTICS.

CHAPTER I.

PRINCIPLES AND DIRECT METHODS.

1.] **Quantics** or **Forms**. A function of any number of variables  $x, y, z, \dots$ , which is rational, integral, and homogeneous in those variables, is called a *quantic* in  $x, y, z, \dots$ . The coefficients in a quantic are constants as far as  $x, y, z, \dots$  are concerned. The idea of the variability of  $x, y, z, \dots$  is rarely introduced. We call them variables only to have a distinctive name for them.

If there be only two variables  $x, y$ , the quantic is spoken of as a *binary* quantic; if three  $x, y, z$ , as a *ternary* quantic; if four, as a *quaternary* quantic; and so on. If there are  $q$  variables, where  $q$  is any number, we may call it a *q-ary* quantic.

The degree of a quantic in the variables  $x, y, z, \dots$  is generally spoken of as its *order*. Quantics of the first, second, third, fourth,  $\dots$ ,  $p$ th orders are called briefly *linear, quadratics, cubics, quartics, \dots p-ics*.

Thus for instance  $ax^3 + 3bx^2y + 3cxy^2 + dy^3$  is a binary cubic, and  $ax^2 + by^2 + cz^2 + 2fyz + 2gzx + 2hxy$  is a ternary quadratic.

By some English and most foreign writers the word *Form* is used as synonymous with and instead of the word *Quantic*. Both words being well established, they will be used almost indiscriminately in this work.

Attention will often be concentrated on *binary* quantics alone. The binary  $p$ -ic will almost invariably be considered in the form

$$a_0 x^p + p a_1 x^{p-1} y + \frac{p(p-1)}{1 \cdot 2} a_2 x^{p-2} y^2 + \frac{p(p-1)(p-2)}{1 \cdot 2 \cdot 3} a_3 x^{p-3} y^3 + \dots + p a_{p-1} x y^{p-1} + a_p y^p,$$

which it is usual shortly to symbolize by

$$(a_0, a_1, a_2, a_3, \dots a_p) (x, y)^p.$$

It is of course clear that, if  $a_0, a_1, a_2, a_3, \dots a_p$  be capable of receiving any values whatever, this is neither more nor less general than the form

$$b_0 x^p + b_1 x^{p-1} y + b_2 x^{p-2} y^2 + b_3 x^{p-3} y^3 + \dots + b_{p-1} x y^{p-1} + b_p y^p,$$

which it is the custom to denote by

$$(b_0, b_1, b_2, b_3, \dots b_p) \uparrow (x, y)^p.$$

The advantages gained by use of the first form, in which the numerical coefficients  $1, p, \frac{p(p-1)}{1 \cdot 2}, \dots$  in the  $p$ th power of a binomial are explicitly introduced as factors of the coefficients in order in the binary  $p$ -ic, will become apparent in the sequel.

Analogous advantages are gained in general, when quantics in higher numbers of variables are being dealt with, by the explicit introduction of multinomial coefficients. Thus in the general  $q$ -ary  $p$ -ic in the variables  $x_1, x_2, \dots x_q$  it is convenient to consider each coefficient to be the product of a factor denoted by a letter, to which any value whatever may be assigned, and the coefficient of the corresponding term in the expansion of  $(x_1 + x_2 + \dots + x_q)^p$ .

When speaking of the *coefficients* in a binary quantic  $(a_0, a_1, a_2, \dots a_p) (x, y)^p$ , we as a rule mean  $a_0, a_1, a_2, \dots a_p$ , and not  $a_0, p a_1, \frac{p(p-1)}{1 \cdot 2} a_2, \dots$ ; and analogously for quantics in higher numbers of variables.

2.] **Linear transformation.** If in a quantic we replace each of the variables by a sum of multiples of first powers of an equally numerous set of new variables, if for instance,

the variables originally involved being  $x, y, z, \dots$ , we substitute for them according to the scheme

$$\begin{aligned} x &= l X + m Y + n Z + \dots, \\ y &= l' X + m' Y + n' Z + \dots, \\ z &= l'' X + m'' Y + n'' Z + \dots, \\ &\dots \end{aligned}$$

where there are just as many of  $X, Y, Z, \dots$  as of  $x, y, z, \dots$ , we are said to make a *linear substitution* in the quantic, or to *linearly transform* the quantic; and the new quantic in  $X, Y, Z, \dots$  which we obtain is spoken of as a *linear transformation* of the original quantic.

The determinant

$$\begin{vmatrix} l, & m, & n, & \dots \\ l', & m', & n', & \dots \\ l'', & m'', & n'', & \dots \\ \dots & \dots & \dots & \dots \end{vmatrix},$$

whose constituents are the coefficients, in their natural order, of the new variables  $X, Y, Z, \dots$  in the expression for the old ones  $x, y, z, \dots$ , and which accordingly consists of as many rows and columns as there are variables in either set, is called the *modulus* of the substitution or of the transformation. It will often be convenient to denote it by a single letter. The letter which will be as a rule chosen is  $M$ .

The original variables  $x, y, z, \dots$  are as a rule taken to be all independent. It is unlawful then to substitute for them any expressions in terms of new variables which are not all independent. Now if  $X, Y, Z, \dots$  are all independent the linear expressions

$$\begin{aligned} l X + m Y + n Z + \dots, \\ l' X + m' Y + n' Z + \dots, \\ l'' X + m'' Y + n'' Z + \dots, \\ \dots \end{aligned}$$

are or are not all independent according as the modulus  $M$  does not or does vanish. We must impose then on the generality of the coefficients in a lawful scheme of linear substitution the one limitation that the modulus  $M$  do not vanish.

We are now in a position to define invariants and covariants.

### 3.] Invariants and Covariants.

An *invariant of a single quantic* is a function of the coefficients in that quantic which needs at most to be multiplied by a factor which is a function only of the coefficients in a scheme of linear substitution to be made equal to the same function of the corresponding coefficients in the quantic into which the given quantic is transformed by that scheme.

An *invariant of two or more quantics* in the same variables is a function of the two or more sets of coefficients in those quantics which needs at most to be multiplied by a factor which is a function only of the coefficients in a scheme of linear substitution to be made equal to the same function of the corresponding coefficients in the quantics into which the given quantics are transformed by that scheme.

A *covariant* of a single quantic, or of two or more quantics in the same variables, is a function of the variables and of the coefficients in that quantic or those quantics which has the like property; namely that of needing at most to be multiplied by a factor which is a function only of the coefficients in a scheme of linear substitution to be made equal to the same function of the new variables and of the corresponding coefficients in the quantic or quantics into which the given quantic or quantics are transformed by that scheme.

For instance, let the binary  $p$ -ic

$$(a_0, a_1, a_2, \dots a_p)(x, y)^p$$

be transformed by the linear substitution

$$x = lX + mY,$$

$$y = l'X + m'Y,$$

and become

$$(A_0, A_1, A_2, \dots A_p)(X, Y)^p,$$

where  $A_0, A_1, A_2, \dots A_p$  are functions of  $a_0, a_1, a_2, \dots a_p$ , and  $l, m, l', m'$ : then  $f(a_0, a_1, a_2, \dots a_p)$  will be an invariant if an identity hold of the form

$$f(A_0, A_1, A_2, \dots A_p) = \phi(l, m, l', m')f(a_0, a_1, a_2, \dots a_p),$$

and  $F(a_0, a_1, a_2, \dots, a_p, x, y)$  will be a covariant if an identity hold of the form

$$F(A_0, A_1, A_2, \dots, A_p, X, Y) \\ = \phi(l, m, l', m') F(a_0, a_1, a_2, \dots, a_p, x, y).$$

Again, if the same substitution transforms another binary quantic in  $x, y$

$$(a'_0, a'_1, a'_2, \dots, a'_{p'}) (x, y)^{p'},$$

into

$$(A'_0, A'_1, A'_2, \dots, A'_{p'}) (X, Y)^{p'},$$

$f(a_0, a_1, \dots, a_p, a'_0, a'_1, \dots, a'_{p'})$  will be an invariant of the  $p$ -ic and the  $p'$ -ic jointly if an identity hold of the form

$$f(A_0, A_1, \dots, A_p, A'_0, A'_1, \dots, A'_{p'}) \\ = \phi(l, m, l', m') f(a_0, a_1, \dots, a_p, a'_0, a'_1, \dots, a'_{p'}),$$

and  $F(a_0, a_1, \dots, a_p, a'_0, a'_1, \dots, a'_{p'}, x, y)$  will be a covariant if an identity hold of the form

$$F(A_0, A_1, \dots, A_p, A'_0, A'_1, \dots, A'_{p'}, X, Y) \\ = \phi(l, m, l', m') F(a_0, a_1, \dots, a_p, a'_0, a'_1, \dots, a'_{p'}, x, y).$$

It will be noticed that covariants include invariants as a particular case.

4.] In every case the factor depending only on the coefficients in the scheme of substitution in the identity which expresses the fact of invariancy or covariancy is as a matter of fact a power of the modulus  $M$ . In particular for any invariant or covariant of a *binary* quantic or *binary* quantics the  $\phi(l, m, l', m')$  above is a power of  $lm' - l'm$ . It is a departure from usual practice not to apparently narrow the definition of invariants and covariants by stating this as a requirement. It will probably be granted that the departure is a proper one, for the necessity is a proposition which can and will be proved hereafter, and, were there any functions such as contemplated in the definitions for which the factor was other than a power of the modulus, their property would be none the less appropriately described as invariantic. The fact that there are not really such functions is one of sufficient interest in itself and of sufficient importance in its applications to deserve proof and prominence.

No limitation requiring functions defined as invariants and

covariants to be rational and integral has been imposed in the definitions. There are in fact irrational and fractional functions which have the property of invariancy and covariancy, as well as others which are rational and integral. The main quest in this work will be however for invariants which are rational and integral, and for covariants which are rational and integral both in the coefficients and in the variables, and the words *invariant* and *covariant* will as a rule be used as meaning *rational integral invariant* and *rational integral covariant*.

There is a greater completeness about a system of rational integral invariants and covariants than is at first sight apparent, in that all invariants and covariants can be expressed in terms of such as are rational and integral. The present is not the stage at which to attempt to prove this fact, but the case of covariants of a single binary  $p$ -ic may be mentioned as an instance. It will be seen in Chapter III that there cannot be more than  $p$  absolutely independent covariants, including the  $p$ -ic itself and invariants. If then we have  $p-1$  absolutely independent ones, any other can be expressed as a function of them. Also in Chapter X a system of  $p$  absolutely independent covariants will be found which are all rational and integral. It will follow that all other covariants, including such as are irrational or fractional, can be expressed in terms of them.

5.] A little careful consideration will show that we ought not to be surprised at the existence of invariants and covariants. Consider for instance a binary quantic. It is equivalent to a product of linear factors, to grant which is only to grant the fundamental theorem of algebra that every rational integral equation has a root, and therefore  $p$  roots if its order be  $p$ . A relation in the coefficients of the quantic will be equivalent to the expression of some special fact with regard to those linear factors. In particular there will be some relations which express kinds of interdependence among two or more factors which are not altered by the application of a linear transformation of the variables. Such a relation will necessitate the corresponding relation among the coefficients in the transformed quantic. In other words the



function of the coefficients of the given quantic whose vanishing gives the relation in question must be a factor of the same function of the corresponding coefficients of the transformed quantic.

Thus in particular the vanishing of the discriminant of a binary quantic is the condition, sufficient and necessary, that the quantic have two identical linear factors. Now, if it have, so clearly must the result of replacing in it  $x$  and  $y$  by  $lX + mY$  and  $l'X + m'Y$ . Consequently, if the discriminant of the given quantic vanish, so too must that of the linearly transformed quantic. In other words the first discriminant must be a factor of the second. That the remaining factor must be a function of  $l, m, l', m'$  only is to be expected because, the discriminant being homogeneous, and each coefficient in the transformed quantic being linear in the coefficients of the untransformed, the degrees of the two discriminants in the coefficients of the untransformed quantic are the same. It will presently be proved with complete rigour that the discriminants of all quantics, and not of binary quantics only, are invariants.

Again, by thinking of the eliminant or resultant of two binary quantics we can realize that invariants of two or more quantics jointly are with equal reason to be expected. The vanishing of the eliminant of two binary quantics is the necessary and sufficient condition for those quantics to have a common factor. If they have, so equally must their linear transformations. In other words, if the eliminant of two binary quantics vanishes, so must that of the two transformed quantics. The former eliminant is then a factor of the latter.

6.] To convince ourselves of the a priori reasonableness of expecting *covariants* to exist, we shall do well to avail ourselves of geometrical representation.

Let us take axes of Cartesian coordinates inclined at any angle, which it is best to regard as unknown, since otherwise we may be in danger of introducing or implying its value in functions with which we deal, and so bringing in ideas not afforded by the quantics and transformation that are before us. The factors of a binary quantic or quantics correspond each to a straight line through the origin, the straight line

in each case whose equation is obtained by equating to zero the factor under consideration.

Let us now consider what is effected by the linear substitution  $x = lX + mY$ ,  $y = l'X + m'Y$ , regarding the substitution not as implying a change of axes but as expressing the coordinates of one point  $(x, y)$  in terms of those of another  $(X, Y)$  with regard to the same axes. The first point  $(x, y)$  being definite, so is the second  $(X, Y)$ . Moreover to different points  $(x, y)$  on a straight line through the origin correspond different points  $(X, Y)$  on another line through the origin, for  $\frac{Y}{X}$  is uniquely determined in terms of  $\frac{y}{x}$ . In fact we have

$$\frac{y}{x} = \frac{l' + m' \frac{Y}{X}}{l + m \frac{Y}{X}}, \quad \text{so that} \quad \frac{Y}{X} = \frac{l \frac{y}{x} - l'}{m' - m \frac{y}{x}}.$$

Now the student of geometry will recognize from this that the two lines on which  $(x, y)$  and  $(X, Y)$  lie have a definite homographic or projective correspondence for given values of  $l, m, l', m'$ . The effect then of the linear transformation is to replace points on lines through the origin by corresponding points on projectively corresponding lines through the origin.

The pencil of lines representative of any given binary quantic or quantics is accordingly replaced by any linear transformation by a projectively corresponding pencil of lines. Is there any other pencil of lines associated with the first pencil, whose projective correspondents are associated with the second pencil exactly as they themselves are with the first? If so, then the equation of their correspondents may be formed either by applying the linear transformation to their equation or by forming an equation from the transformed quantic or quantics in precisely the same way as their equation was formed from the given quantic or quantics. In other words, the derived quantic which equated to zero gives their equation, and that derived in like manner from the transformed quantic or quantics, are identical, but for a possible factor independent of the coordinates. Such a derived quantic will be a covariant if only the factor involve merely the constants  $l, m, l', m'$  of the transformation and not also the coefficients in the quantic or quantics. But

the factor must be of no dimensions in the coefficients of the quantic or quantics, for each coefficient in the transformed quantic or quantics is homogeneous and of one dimension in them. Thus it is to be expected that what is required will be the case.

Now as a rule there are of course lines associated with a given pencil in such a way that if they and the pencil are replaced by others by projective transformation the character of the association is preserved. In particular harmonic properties are unaltered by projective transformation.

Thus, for instance, it suggests itself that a binary quadratic and a linear form have jointly a linear covariant, namely the harmonic conjugate of the linear with regard to the quadratic—see Ex. 6 below: or again, that two binary quadratics have jointly a quadratic covariant, their common pair of harmonic conjugates—see Ex. 7 below: or once more, that a binary cubic has a cubic covariant, composed of the three harmonic conjugates of the three factors singly each with regard to the other two factors—a fact which will be established later.

7.] We have now suggested to us a number of classes of functions which are likely to be invariants and covariants, and which may be examined by the direct method of substitution.

Ex. 1. To verify that  $ac - b^2$  is an invariant of the binary quadratic  $ax^2 + 2bxy + cy^2$ , of which it is the discriminant.

If by the substitution  $x = lX + mY$ ,  $y = l'X + m'Y$ ,

$$ax^2 + 2bxy + cy^2 \text{ become } AX^2 + 2BXY + CY^2,$$

we have

$$A = al^2 + 2bl'l' + cl'^2,$$

$$B = alm + b(lm' + l'm) + cl'm',$$

$$C = am^2 + 2bmm' + cm'^2.$$

Hence at once

$$\begin{aligned} AC - B^2 &= (l^2m'^2 - 2l'mm' + m^2l'^2)(ac - b^2) \\ &= M^2(ac - b^2). \end{aligned}$$

Ex. 2. Verify that the eliminant  $ab' - a'b$  is an invariant of the two binary linear forms  $ax + by$ ,  $a'x + b'y$ .

$$\text{Ans. } AB' - A'B = M(ab' - a'b).$$

Ex. 3. Verify that the eliminant  $ab'^2 - 2ba'b' + ca'^2$  is an invariant of the quadratic and linear forms  $ax^2 + 2bxy + cy^2$ ,  $a'x + b'y$ .

$$\text{Ans. } AB'^2 - 2BA'B' + CA'^2 = M^2(ab'^2 - 2ba'b' + ca'^2).$$

Ex. 4. Verify that  $ac' + a'c - 2bb'$  is an invariant of the two binary quadratics  $ax^2 + 2bxy + cy^2$ ,  $a'x^2 + 2b'xy + c'y^2$ .

*Ans.*  $AC' + A'C - 2BB' = M^2(ac' + a'c - 2bb')$ . The vanishing of this invariant is the condition that the two quadratics denote pairs of harmonic conjugates.

Ex. 5. Verify that  $ae - 4bd + 3c^2$  is an invariant of the binary quartic  $ax^4 + 4bx^3y + 6cx^2y^2 + 4dxy^3 + ey^4$ .

$$\text{Ans. } AE - 4BD + 3C^2 = M^4(ae - 4bd + 3c^2).$$

\* Ex. 6. Verify that  $b'(ax + by) - a'(bx + cy)$  is a covariant of the binary quadratic and linear forms  $ax^2 + 2bxy + cy^2$ ,  $a'x + b'y$ .

$$\begin{aligned} \text{Ans. } B'(AX + BY) - A'(BX + CY) \\ = M\{b'(ax + by) - a'(bx + cy)\}. \end{aligned}$$

This covariant is the harmonic conjugate of the linear with regard to the quadratic form.

Ex. 7. Verify that  $(ab' - a'b)x^2 + (ac' - a'c)xy + (bc' - b'c)y^2$  is a covariant of the two quadratics  $ax^2 + 2bxy + cy^2$ ,  $a'x^2 + 2b'xy + c'y^2$ .

$$\begin{aligned} \text{Ans. } (AB' - A'B)X^2 + (AC' - A'C)XY + (BC' - B'C)Y^2 \\ = M\{(ab' - a'b)x^2 + (ac' - a'c)xy + (bc' - b'c)y^2\}. \end{aligned}$$

This covariant is the common pair of harmonic conjugates with regard to the two quadratics.

Ex. 8. Verify that  $(ac - b^2)x^2 + (ad - bc)xy + (bd - c^2)y^2$  is a covariant of the binary cubic  $ax^3 + 3bx^2y + 3cxy^2 + dy^3$ .

$$\begin{aligned} \text{Ans. } (AC - B^2)X^2 + (AD - BC)XY + (BD - C^2)Y^2 \\ = M^2\{(ac - b^2)x^2 + (ad - bc)xy + (bd - c^2)y^2\}. \end{aligned}$$

8.] Several of the above examples are particular cases of general facts, the proof of which will next occupy us.

Thus Example 1 is a particular case of the general theorem that the discriminant, or eliminant of the various first partial differential coefficients, of any quantic whatever, is an invariant of that quantic.

Again, Examples 2 and 3 are cases of the general fact that the eliminant or resultant of any number of quantics in as many variables is an invariant of those quantics jointly.

Examples 6 and 7 are cases of the theorem that the Jacobian or Functional Determinant (§ 10) of any number of quantics in as many variables is a covariant.

Once more, Example 8 is a case of the fact that the Hessian

of a quantic, i.e. the Jacobian of its first partial differential coefficients, is a covariant of the quantic.

9.] **Eliminant of linear forms.** We may at once prove a first extension of Example 2, that the eliminant of any number of *linear* forms in that same number of variables is ~~the~~ <sup>an</sup> *covariant* of those linear forms.

Let there be  $n$  variables  $x_1, x_2, x_3, \dots, x_n$ , and let the  $n$  linear forms be

$$\begin{aligned} & a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n, \\ & a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n, \\ & \dots \dots \dots \dots \dots \dots \dots \dots \\ & a_{n1}x_1 + a_{n2}x_2 + \dots + a_{nn}x_n; \end{aligned}$$

and let the scheme of linear substitution

$$\begin{aligned} x_1 &= l_{11}X_1 + l_{12}X_2 + \dots + l_{1n}X_n, \\ x_2 &= l_{21}X_1 + l_{22}X_2 + \dots + l_{2n}X_n, \\ &\dots \dots \dots \dots \dots \dots \dots \dots \\ x_n &= l_{n1}X_1 + l_{n2}X_2 + \dots + l_{nn}X_n, \end{aligned}$$

transform them into

$$\begin{aligned} & A_{11}X_1 + A_{12}X_2 + \dots + A_{1n}X_n, \\ & A_{21}X_1 + A_{22}X_2 + \dots + A_{2n}X_n, \\ & \dots \dots \dots \dots \dots \dots \dots \dots \\ & A_{n1}X_1 + A_{n2}X_2 + \dots + A_{nn}X_n. \end{aligned}$$

Then we see at once that, each of  $r$  and  $s$  being any number between 1 and  $n$  inclusive,

$$A_{rs} = l_{1s}a_{r1} + l_{2s}a_{r2} + l_{3s}a_{r3} + \dots + l_{ns}a_{rn};$$

so that, by the ordinary theorem for the multiplication of determinants,

$$\begin{aligned} \begin{vmatrix} A_{11}, A_{12}, \dots, A_{1n} \\ A_{21}, A_{22}, \dots, A_{2n} \\ \dots \dots \dots \dots \dots \\ A_{n1}, A_{n2}, \dots, A_{nn} \end{vmatrix} &= \begin{vmatrix} l_{11}, l_{12}, \dots, l_{1n} \\ l_{21}, l_{22}, \dots, l_{2n} \\ \dots \dots \dots \dots \dots \\ l_{n1}, l_{n2}, \dots, l_{nn} \end{vmatrix} \times \begin{vmatrix} a_{11}, a_{12}, \dots, a_{1n} \\ a_{21}, a_{22}, \dots, a_{2n} \\ \dots \dots \dots \dots \dots \\ a_{n1}, a_{n2}, \dots, a_{nn} \end{vmatrix} \\ &= M \begin{vmatrix} a_{11}, a_{12}, \dots, a_{1n} \\ a_{21}, a_{22}, \dots, a_{2n} \\ \dots \dots \dots \dots \dots \\ a_{n1}, a_{n2}, \dots, a_{nn} \end{vmatrix}. \end{aligned}$$

10.] **Jacobians are covariants.** If  $u, v, w, \dots$  are any number of quantities in that same number of variables  $x, y, z, \dots$ , the Jacobian or Functional Determinant of  $u, v, w, \dots$  is the determinant

$$\begin{vmatrix} \frac{du}{dx} & \frac{du}{dy} & \frac{du}{dz} & \dots \\ \frac{dv}{dx} & \frac{dv}{dy} & \frac{dv}{dz} & \dots \\ \frac{dw}{dx} & \frac{dw}{dy} & \frac{dw}{dz} & \dots \\ \dots & \dots & \dots & \dots \end{vmatrix},$$

which it is usual more shortly to write

$$\frac{d(u, v, w, \dots)}{d(x, y, z, \dots)}.$$

That it is a covariant of  $u, v, w, \dots$  may be seen as follows. If in any function  $u$  of  $x, y, z, \dots$  we substitute for these variables according to the linear scheme

$$\begin{aligned} x &= lX + mY + nZ + \dots, \\ y &= l'X + m'Y + n'Z + \dots, \\ z &= l''X + m''Y + n''Z + \dots, \\ &\dots \end{aligned}$$

in this way expressing it as a function of  $X, Y, Z, \dots$ , we have at once

$$\begin{aligned} \frac{du}{dX} &= \frac{du}{dx} \cdot \frac{dx}{dX} + \frac{du}{dy} \cdot \frac{dy}{dX} + \frac{du}{dz} \cdot \frac{dz}{dX} + \dots \\ &= l \frac{du}{dx} + l' \frac{du}{dy} + l'' \frac{du}{dz} + \dots, \end{aligned}$$

all the differential coefficients being partial.

Similarly

$$\begin{aligned} \frac{du}{dY} &= m \frac{du}{dx} + m' \frac{du}{dy} + m'' \frac{du}{dz} + \dots, \\ \frac{du}{dZ} &= n \frac{du}{dx} + n' \frac{du}{dy} + n'' \frac{du}{dz} + \dots, \\ &\text{\&c., \&c.} \end{aligned}$$

Thus, by the rule for multiplication of determinants of the same order,

$$\begin{vmatrix} l, & m, & n, & \dots \\ l', & m', & n', & \dots \\ l'', & m'', & n'', & \dots \\ \dots & \dots & \dots & \dots \end{vmatrix} \times \begin{vmatrix} \frac{du}{dx}, & \frac{du}{dy}, & \frac{du}{dz}, & \dots \\ \frac{dv}{dx}, & \frac{dv}{dy}, & \frac{dv}{dz}, & \dots \\ \frac{dw}{dx}, & \frac{dw}{dy}, & \frac{dw}{dz}, & \dots \\ \dots & \dots & \dots & \dots \end{vmatrix} \\
 = \begin{vmatrix} l \frac{du}{dx} + l' \frac{du}{dy} + l'' \frac{du}{dz} + \dots, & m \frac{du}{dx} + m' \frac{du}{dy} + m'' \frac{du}{dz} + \dots, & \dots \\ l \frac{dv}{dx} + l' \frac{dv}{dy} + l'' \frac{dv}{dz} + \dots, & m \frac{dv}{dx} + m' \frac{dv}{dy} + m'' \frac{dv}{dz} + \dots, & \dots \\ \dots & \dots & \dots \\ \dots & \dots & \dots \end{vmatrix} \\
 = \begin{vmatrix} \frac{du}{dX}, & \frac{du}{dY}, & \frac{du}{dZ}, & \dots \\ \frac{dv}{dX}, & \frac{dv}{dY}, & \frac{dv}{dZ}, & \dots \\ \frac{dw}{dX}, & \frac{dw}{dY}, & \frac{dw}{dZ}, & \dots \\ \dots & \dots & \dots & \dots \end{vmatrix} .$$

And this, transposing the right and left-hand sides, is

$$\frac{d(u, v, w, \dots)}{d(X, Y, Z, \dots)} = M \frac{d(u, v, w, \dots)}{d(x, y, z, \dots)},$$

where on the right  $u, v, w, \dots$  are expressed in terms of  $x, y, z, \dots$ , i.e. in their original forms, and on the left in terms of  $X, Y, Z, \dots$ , i.e. in their transformed forms. Thus the Jacobian of  $u, v, w, \dots$  is a covariant.

Covariants, as stated already, include invariants as a particular case. Should the Jacobian not involve the variables it is an invariant. This is the case when  $u, v, w, \dots$  are all of the first order in  $x, y, z, \dots$ . Thus the present result includes that of the preceding article.

Ex. 9. Obtain from this result examples 2, 6 and 7 of § 7.

Ex. 10. Obtain a linear covariant of the ternary quadratic and two linear forms

$$\begin{aligned} ax^2 + by^2 + cz^2 + 2fyz + 2gzx + 2hxy, \\ a'x + b'y + c'z, \\ a''x + b''y + c''z; \end{aligned}$$

and interpret it geometrically, by taking for  $X$  and  $Y$  the two linear forms, or otherwise.

$$\begin{aligned} \text{Ans. } (b'c'' - b''c') (ax + hy + gz) + (c'a'' - c''a') (hx + by + fz) \\ + (a'b'' - a''b') (gx + fy + cz). \end{aligned}$$

The polar of the intersection of two straight lines with regard to a conic.

Ex. 11. Two ternary quadratics and a linear form have a quadratic covariant.

Ex. 12. Obtain and interpret geometrically a linear covariant of the quaternary quadratic and three linear forms

$$\begin{aligned} ax^2 + by^2 + cz^2 + d^2w^2 + 2fyz + 2gzx + 2hxy + 2pxw + 2qyw + 2rzw, \\ a'x + b'y + c'z + d'w, \\ a''x + b''y + c''z + d''w, \\ a'''x + b'''y + c'''z + d'''w. \end{aligned}$$

11.] **Hessians are covariants.** To prove that the *Hessian*

$$\begin{vmatrix} \frac{d^2u}{dx^2}, & \frac{d^2u}{dxdy}, & \frac{d^2u}{dxdz}, & \dots \\ \frac{d^2u}{dxdy}, & \frac{d^2u}{dy^2}, & \frac{d^2u}{dydz}, & \dots \\ \frac{d^2u}{dxdz}, & \frac{d^2u}{dydz}, & \frac{d^2u}{dz^2}, & \dots \\ \dots & \dots & \dots & \dots \end{vmatrix}$$

of a quantic  $u$  in the variables  $x, y, z, \dots$  is a covariant of  $u$ .

A natural but erroneous form of argument must first be guarded against. The Hessian of  $u$  is the Jacobian of  $\frac{du}{dx}, \frac{du}{dy}, \frac{du}{dz}, \dots$ . Hence by the last article it is a covariant of the system of quantics  $\frac{du}{dx}, \frac{du}{dy}, \frac{du}{dz}, \dots$ . It would be unjustifiable hence to conclude that it is a covariant of  $u$ , for when  $u$  is transformed by a linear substitution  $\frac{du}{dx}, \frac{du}{dy}, \frac{du}{dz}, \dots$  are not transformed into  $\frac{du}{dX}, \frac{du}{dY}, \frac{du}{dZ}, \dots$ .



A correct method of proving the theorem is the following. Multiply the Hessian written above by the modulus

$$\begin{vmatrix} l, & m, & n, & \dots \\ l', & m', & n', & \dots \\ l'', & m'', & n'', & \dots \\ \dots & \dots & \dots & \dots \end{vmatrix}, \text{ or } M,$$

of the transforming linear substitution. Using the facts, employed in § 10, that, when the operations are on any function of  $x, y, z, \dots$  upon the right, and on its equivalent in terms of  $X, Y, Z, \dots$  upon the left,

$$\frac{d}{dX} = l \frac{d}{dx} + l' \frac{d}{dy} + l'' \frac{d}{dz} + \dots,$$

$$\frac{d}{dY} = m \frac{d}{dx} + m' \frac{d}{dy} + m'' \frac{d}{dz} + \dots$$

&c., &c.

we see at once that the product may be written

$$\begin{vmatrix} \frac{d}{dX} \cdot \frac{du}{dx}, & \frac{d}{dY} \cdot \frac{du}{dx}, & \frac{d}{dZ} \cdot \frac{du}{dx}, & \dots \\ \frac{d}{dX} \cdot \frac{du}{dy}, & \frac{d}{dY} \cdot \frac{du}{dy}, & \frac{d}{dZ} \cdot \frac{du}{dy}, & \dots \\ \frac{d}{dX} \cdot \frac{du}{dz}, & \frac{d}{dY} \cdot \frac{du}{dz}, & \frac{d}{dZ} \cdot \frac{du}{dz}, & \dots \\ \dots & \dots & \dots & \dots \end{vmatrix},$$

or, since the order of differentiations in such an operator as

$$\frac{d}{dX} \cdot \frac{d}{dx} \equiv \left( l \frac{d}{dx} + l' \frac{d}{dy} + l'' \frac{d}{dz} + \dots \right) \frac{d}{dx}$$

may be interchanged, that it may be written

$$\begin{vmatrix} \frac{d}{dx} \cdot \frac{du}{dX}, & \frac{d}{dx} \cdot \frac{du}{dY}, & \frac{d}{dx} \cdot \frac{du}{dZ}, & \dots \\ \frac{d}{dy} \cdot \frac{du}{dX}, & \frac{d}{dy} \cdot \frac{du}{dY}, & \frac{d}{dy} \cdot \frac{du}{dZ}, & \dots \\ \frac{d}{dz} \cdot \frac{du}{dX}, & \frac{d}{dz} \cdot \frac{du}{dY}, & \frac{d}{dz} \cdot \frac{du}{dZ}, & \dots \\ \dots & \dots & \dots & \dots \end{vmatrix}.$$

Now multiply again by  $M$ , taking this time columns with columns in forming the product. The same equivalences of operators as before tell us that the result is

$$\begin{vmatrix} \frac{d^2u}{dX^2} & \frac{d^2u}{dXdY} & \frac{d^2u}{dXdZ} & \cdots \\ \frac{d^2u}{dXdY} & \frac{d^2u}{dY^2} & \frac{d^2u}{dYdZ} & \cdots \\ \frac{d^2u}{dXdZ} & \frac{d^2u}{dYdZ} & \frac{d^2u}{dZ^2} & \cdots \\ \dots & \dots & \dots & \dots \end{vmatrix}$$

Thus upon multiplying the Hessian of the untransformed quantic by  $M^2$ , the square of the modulus, we have obtained the Hessian of the transformed. The Hessian is then a covariant.

When the quantic  $u$  is binary only the Hessian is

$$\frac{d^2u}{dx^2} \frac{d^2u}{dy^2} - \left( \frac{d^2u}{dxdy} \right)^2.$$

Ex. 13. Apply this result to prove Ex. 8 of § 7.

Ex. 14. If the covariant  $(ac-b^2)x^2 + (ad-bc)xy + (bd-c^2)y^2$  of the binary cubic  $ax^3 + 3bx^2y + 3cxy^2 + dy^3$  be broken up into factors  $(px+qy)(p'x+q'y)$ , and if these factors be taken for  $X$  and  $Y$ , so that the formulae of linear transformation are

$$x = \frac{q'X - qY}{pq' - p'q}, \quad y = \frac{-p'X + pY}{pq' - p'q},$$

show that the cubic takes the form  $AX^3 + DY^3$ .

Ex. 15. Hence solve the cubic  $ax^3 + 3bx^2 + 3cx + d = 0$ . (Cf. § 200.)

Ex. 16. Find a covariant of degree 2 in the coefficients and order 4 in the variables of the binary quartic  $(a, b, c, d, e)(x, y)^4$ .

$$\text{Ans. } (ax^2 + 2bxy + cy^2)(cx^2 + 2dxy + ey^2) - (bx^2 + 2cxy + dy^2)^2.$$

Ex. 17. Find a covariant of degree 2 and order 6 of the binary quintic  $(a, b, c, d, e, f)(x, y)^5$ .

Ex. 18. The Hessian of a quartic, in any number of variables, which has a cubed factor is the fourth power of that factor, multiplied by a function of the coefficients. (*Cayley*.)

*Ans.* Take the cubed factor for  $X^3$ .

12.] **Discriminants of Quadratics.** The Hessian of a quantic, proved above to be in general a covariant, is in particular an invariant if it be free from the variables. This is the case if the quantic be a quadratic in any number of variables.

We have accordingly the proof of a first generalization of § 7, Ex. 1, namely that the discriminant of any quadratic is an invariant of that quadratic. For the Hessians of the binary, ternary, and quaternary quadratics

$$\begin{aligned} & ax^2 + 2bxy + cy^2, \\ & ax^2 + by^2 + cz^2 + 2fyz + 2gzx + 2hxy, \\ & ax^2 + by^2 + cz^2 + dw^2 + 2fyz + 2gzx + 2hxy \\ & \qquad \qquad \qquad + 2pxw + 2qyw + 2rzw, \end{aligned}$$

are, after rejection of the numerical factors  $2^2$ ,  $2^3$ ,  $2^4$ ,

$$\begin{vmatrix} a, & b \\ b, & c \end{vmatrix}, \quad \begin{vmatrix} a, & h, & g \\ h, & b, & f \\ g, & f, & c \end{vmatrix}, \quad \begin{vmatrix} a, & h, & g, & p \\ h, & b, & f, & q \\ g, & f, & c, & r \\ p, & q, & r, & d \end{vmatrix},$$

and, quite generally, that of the  $q$ -ary quadratic

$$\sum_{n=1}^{n=q} a_{nn} x_n^2 + 2 \sum_{m=1}^{m=q-1} \sum_{n=m+1}^{n=q} a_{mn} x_m x_n,$$

is, after rejection of the numerical factor  $2^q$ ,

$$\begin{vmatrix} a_{11}, & a_{12}, & a_{13}, & \dots & a_{1q} \\ a_{12}, & a_{22}, & a_{23}, & \dots & a_{2q} \\ a_{13}, & a_{23}, & a_{33}, & \dots & a_{3q} \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ a_{1q}, & a_{2q}, & a_{3q}, & \dots & a_{qq} \end{vmatrix}.$$

Now these are the eliminants of the first partial differential coefficients, each divided by 2, of the various quadratics; i. e. they are the discriminants of the quadratics.

13.] **Eliminants are invariants.** Let the  $q$  quantics  $u, v, w, \dots$  in as many variables  $x, y, z, \dots$  become  $U, V, W, \dots$

when  $x, y, z, \dots$  are replaced according to the linear scheme

$$x = l X + m Y + n Z + \dots,$$

$$y = l' X + m' Y + n' Z + \dots,$$

$$z = l'' X + m'' Y + n'' Z + \dots,$$

$$\dots \dots \dots$$

Let  $R(a, b, a', \dots)$  denote the eliminant or resultant of  $u, v, w, \dots$ , and  $R(A, B, A', \dots)$  that of  $U, V, W, \dots$ ,  $a$  and  $A, b$  and  $B, a'$  and  $A', \dots$  being corresponding coefficients in untransformed and transformed quantities. It is to be proved that  $R(a, b, a', \dots)$  is an invariant of  $u, v, w, \dots$

If  $R(A, B, A', \dots)$  vanish,  $U, V, W, \dots$  are made simultaneously to vanish by some set of values of  $X, Y, Z, \dots$  which are not all zero. The above expressions for  $x, y, z, \dots$  in terms of  $X, Y, Z, \dots$  then determine a corresponding set of values of  $x, y, z, \dots$  which make  $u, v, w, \dots$  vanish simultaneously. If these are not all zero it must follow that  $R(a, b, a', \dots)$  vanishes. On the other hand, if they be all zero it is necessitated that

$$\begin{vmatrix} l, m, n, \dots \\ l', m', n', \dots \\ l'', m'', n'', \dots \\ \dots \end{vmatrix},$$

i. e.  $M$ , vanishes.

Thus if  $R(A, B, A', \dots) = 0$  it follows that either

$$R(a, b, a', \dots) = 0 \text{ or } M = 0.$$

Again, if  $M = 0$ , whether  $R(a, b, a', \dots) = 0$  or not, the  $q$  linear functions

$$lX + mY + nZ + \dots, l'X + m'Y + n'Z + \dots, l''X + m''Y + n''Z + \dots$$

are not linearly independent, but all vanish when for  $X, Y, Z, \dots$  are taken the solutions of any  $q-1$  of them. Consequently, if this be so,  $U, V, W, \dots$  can be made simultaneously to vanish by values not all zero of  $X, Y, Z, \dots$ , and therefore

$$R(A, B, A', \dots) = 0.$$

Also if  $R(a, b, a', \dots) = 0$ , even when  $M$  does not vanish, there are values of  $x, y, z, \dots$ , not all zero, which satisfy  $u = 0, v = 0, w = 0, \dots$  simultaneously; and these determine by the above equations corresponding values of  $X, Y, Z, \dots$

which satisfy  $U = 0, V = 0, W = 0, \dots$  simultaneously, so that  $R(A, B, A', \dots) = 0$ .

Thus the condition  $R(A, B, A', \dots) = 0$  expresses exactly the same special state of things as do the alternative conditions  $M = 0, R(a, b, a', \dots) = 0$ .

Hence, assuming, as we shall prove in the next article, that the algebraic function  $M$  is not resolvable into simpler algebraic factors, but not assuming the unproved fact that  $R(a, b, a', \dots)$  is not so resolvable,

$$R(A, B, A', \dots) = \text{power of } M \times F(a, b, a', \dots),$$

where  $F(a, b, a', \dots)$ , if not  $R(a, b, a', \dots)$  or a power of it, is at any rate a product of powers of all the factors of  $R(a, b, a', \dots)$ , supposing for safety that it may have simpler factors.

This result is proved for all linear substitutions. It holds then for every particular linear substitution. Now take  $l, m', n'', \dots$  all units and the other coefficients in the scheme all zeros, so that the scheme becomes simply  $x = X, y = Y, z = Z, \dots$ , and  $M = 1$ , while  $A, B, A', \dots$  are merely  $a, b, a', \dots$ : then our general result gives

$$R(a, b, a', \dots) = F(a, b, a', \dots),$$

so that  $F(a, b, a', \dots)$  is really the eliminant of  $u, v, w, \dots$  itself.

Consequently the general result is

$$R(A, B, A', \dots) = \text{power of } M \times R(a, b, a', \dots),$$

which proves that the eliminant  $R(a, b, a', \dots)$  is an invariant.

We now give the proof that  $M$  is irresolvable.

14.] **The modulus irresolvable into factors.** Let us use a double suffix notation, and suppose, if possible, that

$$M \equiv \begin{vmatrix} l_{11}, l_{12}, l_{13}, \dots \\ l_{21}, l_{22}, l_{23}, \dots \\ l_{31}, l_{32}, l_{33}, \dots \\ \dots \end{vmatrix}$$

can be written as a product of two rational factors  $\theta\phi$ .

The determinant is of the first degree in every constituent. Thus  $l_{11}$  cannot occur in both factors  $\theta, \phi$ . Suppose that it occurs in  $\theta$ .

In the expansion of the determinant no term occurs in which  $l_{11}$  is multiplied by any constituent belonging to its row or its column. Thus  $\phi$  can involve no constituent belonging to the first row or the first column. Let  $l_{r,s}$  be a constituent which does occur in  $\phi$ . By similar reasoning no constituent belonging to the  $r$ th row or  $s$ th column can occur in  $\theta$ .

Thus two constituents,  $l_{r1}$  and  $l_{1s}$ , cannot occur either in  $\theta$  or in  $\phi$ . But the expansion of the determinant involves every constituent. Our supposition that  $M$  can be written as a product of factors  $\theta\phi$  is therefore untenable.

15.] **All discriminants are invariants.** Of this proposition, already proved for quadratics, a general demonstration will now be given.

If  $u$  be a quantic in  $q$  variables  $x, y, z, \dots$  we have to prove that its discriminant, i.e. the eliminant of its  $q$  first differential coefficients  $\frac{du}{dx}, \frac{du}{dy}, \frac{du}{dz}, \dots$ , is an invariant of  $u$ .

The scheme of linear substitution being the usual one, we have, as in § 10,

$$\frac{du}{dX} = l \frac{du}{dx} + l' \frac{du}{dy} + l'' \frac{du}{dz} + \dots,$$

$$\frac{du}{dY} = m \frac{du}{dx} + m' \frac{du}{dy} + m'' \frac{du}{dz} + \dots,$$

$$\frac{du}{dZ} = n \frac{du}{dx} + n' \frac{du}{dy} + n'' \frac{du}{dz} + \dots,$$

$$\dots \dots \dots$$

Now, in accordance with the definition, the discriminant  $\Delta(A, B, \dots)$  of the transformed form of  $u$  will vanish if and only if  $\frac{du}{dX}, \frac{du}{dY}, \frac{du}{dZ}, \dots$  are made simultaneously to vanish by some set of values, not all zero, of  $X, Y, Z, \dots$ . But the above equivalences tell us that this will be the case if and only if either (1)  $\frac{du}{dx}, \frac{du}{dy}, \frac{du}{dz}, \dots$  can be made simultaneously

to vanish by values of  $x, y, z, \dots$  not all zero, i.e. if the discriminant  $\Delta(a, b, \dots)$  of the untransformed  $u$  vanishes, or (2) if

$$\begin{vmatrix} l, & l', & l'', & \dots \\ m, & m', & m'', & \dots \\ n, & n', & n'', & \dots \\ \cdot & \cdot & \cdot & \cdot \end{vmatrix},$$

i. e.  $M$  the modulus of the substitution, vanishes.

It follows therefore, since  $M$  is irresoluble, that

$$\Delta(A, B, \dots) = \text{power of } M \times \Delta'(a, b, \dots),$$

where  $\Delta'(a, b, \dots)$ , if not  $\Delta(a, b, \dots)$  itself or a power of it, is at any rate the product of powers of the factors into which we might allow the possibility of  $\Delta(a, b, \dots)$  breaking up.

Apply however the general result to the case of the particular substitution  $x = X, y = Y, z = Z, \dots$ , for which  $M = 1$  and  $A = a, B = b, \&c.$  It becomes

$$\Delta(a, b, \dots) = \Delta'(a, b, \dots).$$

Thus our general conclusion is that

$$\Delta(A, B, \dots) = \text{power of } M \times \Delta(a, b, \dots).$$

Consequently the discriminant  $\Delta(a, b, \dots)$  is an invariant.

16.] **Determinant expressions for powers of  $lm' - l'm$ .** For purposes of direct proofs that large classes of functions in determinant form are invariants and covariants of *binary* quantics, a simple theorem, due to Faa de Bruno, as to a certain class of determinants, is of great utility. The first three cases of the theorem are

$$\begin{vmatrix} l, & m \\ l', & m' \end{vmatrix} = lm' - l'm,$$

$$\begin{vmatrix} l^2, & lm, & m^2 \\ 2ll', & lm' + l'm, & 2mm' \\ l'^2, & l'm', & m'^2 \end{vmatrix} = (lm' - l'm)^3,$$

$$\begin{vmatrix} l^3, & l^2m, & lm^2, & m^3 \\ 3l^2l', & 2ll'm + l^2m', & 2lmm' + l'm^2, & 3m^2m' \\ 3ll'^2, & l'^2m + 2ll'm', & lm'^2 + 2l'mm', & 3mm'^2 \\ l'^3, & l'^2m', & l'm'^2, & m'^3 \end{vmatrix} = (lm' - l'm)^6;$$

and the general theorem is that the determinant whose first row consists of the constituents

$$l^r, l^{r-1}m, l^{r-2}m^2, \dots, lm^{r-1}, m^r,$$

and whose other rows are obtained in succession by operating on the constituents of this first row with

$$l' \frac{d}{dl} + m' \frac{d}{dm}, \quad \frac{1}{1 \cdot 2} \left( l' \frac{d}{dl} + m' \frac{d}{dm} \right)^2, \quad \dots \quad \frac{1}{r!} \left( l' \frac{d}{dl} + m' \frac{d}{dm} \right)^r,$$

is a power, namely the  $\frac{1}{2}r(r+1)$ th power, of  $lm' - l'm$ .

It will be readily seen that we might equally write down first the last row

$$l'^r, l'^{r-1}m', l'^{r-2}m'^2, \dots, l'm'^{r-1}, m'^r,$$

and obtain the other rows in succession upwards by operations on it with

$$l \frac{d}{dl'} + m \frac{d}{dm'}, \quad \frac{1}{1 \cdot 2} \left( l \frac{d}{dl'} + m \frac{d}{dm'} \right)^2, \quad \dots \quad \frac{1}{r!} \left( l \frac{d}{dl'} + m \frac{d}{dm'} \right)^r.$$

For the constituents in the  $(s+1)$ th column, read downwards, are the coefficients of the various powers of  $t$  in the expansion of

$$(l + tl')^{r-s} (m + tm')^s$$

by Taylor's theorem; and the same, read upwards, are the coefficients of powers of  $\tau$  in the expansion of

$$(\tau l + l')^{r-s} (\tau m + m')^s.$$

We speak below of the two modes of forming the determinant as the first and second ways of writing it down.

The first case of the theorem is immediate. The second is at once proved by adding to the first row  $-\frac{m}{m'}$  times the second and  $\frac{m^2}{m'^2}$  times the third: and the third case is easily proved in a similar manner. The general theorem is an easy exercise on the theory of Lagrange's solution of linear partial differential equations, as we proceed to show.

By the ordinary rule for differentiation of products we know that the result of differentiating a determinant of the



$r$ th order can be written as a sum of  $r$  determinants, each obtained by differentiating the constituents of one row, leaving the constituents of all the other rows unaltered. Now operate on the given determinant, thinking of it as written down in its first way, with  $l' \frac{d}{dl} + m' \frac{d}{dm}$ . The result is a sum of  $r$  determinants all of which vanish. For the result of operating on any row except the last is to produce a numerical multiple of the following row, and the result of operating on the last row is to produce a row of zeros. If then  $D$  denote the determinant, we have

$$l' \frac{dD}{dl} + m' \frac{dD}{dm} = 0.$$

Hence by Lagrange's theory  $D$  involves  $l$  and  $m$  only in the connexion  $lm' - l'm$ .

Again, think of  $D$  as written down in its second way, and operate on it with  $l \frac{d}{dl'} + m \frac{d}{dm'}$ . We obtain in like manner

$$l \frac{dD}{dl'} + m \frac{dD}{dm'} = 0,$$

so that  $D$  involves  $l'$  and  $m'$  only in the connexion  $lm' - l'm$ .

Thus  $D$  is a function of  $lm' - l'm$  only; and, being homogeneous, must consist of a single power of  $lm' - l'm$ , with a possible numerical factor. But this numerical factor is unity, as we see for instance by taking  $l = m' = 1$ ,  $l' = m = 0$ , for which  $lm' - l'm$  is unity and  $D$  consists of a principal diagonal of units with all other constituents zero.

That the power of  $lm' - l'm$  is the  $\frac{1}{2}r(r+1)$ th follows from the fact that  $D$  is of dimensions  $r(r+1)$  in  $l, m, l', m'$ .

17.] As a typical application of this theorem let us prove that

$$\begin{vmatrix} \frac{d^4 u}{dx^4} & \frac{d^4 u}{dx^3 dy} & \frac{d^4 u}{dx^2 dy^2} \\ \frac{d^4 u}{dx^3 dy} & \frac{d^4 u}{dx^2 dy^2} & \frac{d^4 u}{dx dy^3} \\ \frac{d^4 u}{dx^2 dy^2} & \frac{d^4 u}{dx dy^3} & \frac{d^4 u}{dy^4} \end{vmatrix}$$

is a covariant of a binary quantic  $u$ , or in particular an invariant if  $u$  is a quartic.

We will multiply twice, taking columns with columns, by the determinant expression above for  $(lm' - l'm)^3$ , i. e.  $M^3$ .

The first multiplication produces, since

$$\left(l \frac{d}{dx} + l' \frac{d}{dy}\right)^2 = \frac{d^2}{dX^2}, \quad \left(l \frac{d}{dx} + l' \frac{d}{dy}\right) \left(m \frac{d}{dx} + m' \frac{d}{dy}\right) = \frac{d^2}{dXdY},$$

$$\left(m \frac{d}{dx} + m' \frac{d}{dy}\right)^2 = \frac{d^2}{dY^2},$$

$$\begin{vmatrix} \frac{d^2}{dX^2} & \frac{d^2u}{dx^2} & \frac{d^2}{dXdY} & \frac{d^2u}{dx^2} & \frac{d^2}{dY^2} & \frac{d^2u}{dx^2} \\ \frac{d^2}{dX^2} & \frac{d^2u}{dx dy} & \frac{d^2}{dXdY} & \frac{d^2u}{dx dy} & \frac{d^2}{dY^2} & \frac{d^2u}{dx dy} \\ \frac{d^2}{dX^2} & \frac{d^2u}{dy^2} & \frac{d^2}{dXdY} & \frac{d^2u}{dy^2} & \frac{d^2}{dY^2} & \frac{d^2u}{dy^2} \end{vmatrix},$$

and the second multiplication of this, with the order of differentiation in each constituent changed, produces

$$\begin{vmatrix} \frac{d^4u}{dX^4} & \frac{d^4u}{dX^3dY} & \frac{d^4u}{dX^2dY^2} \\ \frac{d^4u}{dX^3dY} & \frac{d^4u}{dX^2dY^2} & \frac{d^4u}{dXdY^3} \\ \frac{d^4u}{dX^2dY^2} & \frac{d^4u}{dXdY^3} & \frac{d^4u}{dY^4} \end{vmatrix}.$$

Thus the fact stated is proved.

Ex. 19. Prove that

$$\begin{vmatrix} a, & b, & c \\ b, & c, & d \\ c, & d, & e \end{vmatrix},$$

$$\text{i. e. } ace + 2bcd - ad^2 - b^2e - c^3,$$

is an invariant of the binary quartic  $(a, b, c, d, e)(x, y)^4$ .

*Ans.* Factor  $M^6$ . This important invariant, usually denoted by  $J$ , is called the *Catalecticant* of the quartic.

Ex. 20. Obtain a covariant of the third order and degree of the binary quintic  $(a, b, c, d, e, f)(x, y)^5$ .

Ans. Its so called *canonizant*

$$\begin{vmatrix} ax+by, & bx+cy, & cx+dy \\ bx+cy, & cx+dy, & dx+ey \\ cx+dy, & dx+ey, & ex+fy \end{vmatrix}.$$

Ex. 21. If  $u_{r,s}$  denote  $\frac{d^{r+s}u}{dx^r dy^s}$ , prove that

$$\begin{vmatrix} u_{60}, & u_{51}, & u_{42}, & u_{33} \\ u_{51}, & u_{42}, & u_{33}, & u_{24} \\ u_{42}, & u_{33}, & u_{24}, & u_{15} \\ u_{33}, & u_{24}, & u_{15}, & u_{06} \end{vmatrix}$$

is a covariant of a binary quantic  $u$  of order greater than 6.

Ans. Factor  $M^{12}$ .

Ex. 22. Prove that the *catalecticant*

$$\begin{vmatrix} a, & b, & c, & d \\ b, & c, & d, & e \\ c, & d, & e, & f \\ d, & e, & f, & g \end{vmatrix}$$

is an invariant of the binary sextic  $(a, b, c, d, e, f, g)(x, y)^6$ .

Ex. 23. Every binary quantic of even order  $2n$  has an invariant, its *catalecticant*, of degree  $n+1$ .

Ans. Factor  $M^{n(n+1)}$ .

Ex. 24. Prove that

$$\begin{vmatrix} \frac{d^2u}{dx^2}, & \frac{d^2u}{dxdy}, & \frac{d^2u}{dy^2} \\ \frac{d^2v}{dx^2}, & \frac{d^2v}{dxdy}, & \frac{d^2v}{dy^2} \\ \frac{d^2w}{dx^2}, & \frac{d^2w}{dxdy}, & \frac{d^2w}{dy^2} \end{vmatrix}$$

is a covariant of three binary quantics  $u, v, w$ .

Ans. Factor  $M^3$ .

Ex. 25. Obtain and geometrically interpret the invariant

$$\begin{vmatrix} a, & b, & c \\ a', & b', & c' \\ a'', & b'', & c'' \end{vmatrix}$$

of three binary quadratics

$$(a, b, c)(x, y)^2, (a', b', c')(x, y)^2, (a'', b'', c'')(x, y)^2.$$

Ans. Criterion of an involution.

Ex. 26. Prove that

$$\begin{vmatrix} \frac{d^2u}{dx^2}, & \frac{d^2u}{dxdy}, & \frac{d^2u}{dy^2} \\ \frac{d^2v}{dx^2}, & \frac{d^2v}{dxdy}, & \frac{d^2v}{dy^2} \\ y^2, & -xy, & x^2 \end{vmatrix}$$

is a covariant of two binary quantics  $u, v$ .

*Ans.* Factor  $\frac{M^3}{M^2} = M$ .

Ex. 27. Deduce Ex. 7 of § 7.

Ex. 28. Prove that

$$\begin{vmatrix} \frac{d^3u}{dx^3}, & \frac{d^3u}{dx^2dy}, & \frac{d^3u}{dxdy^2} \\ \frac{d^3u}{dx^2dy}, & \frac{d^3u}{dxdy^2}, & \frac{d^3u}{dy^3} \\ y^2, & -xy, & x^2 \end{vmatrix}$$

is a covariant of a binary quantic  $u$ .

*Ans.* Factor  $M^2$ . Multiply first by the determinant expression for  $M^3$ , and then by  $M$  in the form

$$\begin{vmatrix} l, m, 0 \\ l', m', 0 \\ 0, 0, 1 \end{vmatrix}.$$

Ex. 29. Prove that

$$\begin{vmatrix} u_{50}, & u_{41}, & u_{32}, & u_{23} \\ u_{41}, & u_{32}, & u_{23}, & u_{14} \\ u_{32}, & u_{23}, & u_{14}, & u_{05} \\ y^3, & -xy^2, & x^2y, & -x^3 \end{vmatrix}$$

is a covariant of a binary quantic  $u$ .

*Ans.* Factor  $M^6$ . Multiply first by the determinant expression for  $M^6$ , and then by that for  $M^3$ .

Ex. 30. Prove that

$$\begin{vmatrix} a, & b, & c, & d \\ b, & c, & d, & e \\ c, & d, & e, & f \\ y^3, & -xy^2, & x^2y, & -x^3 \end{vmatrix}$$

is a covariant of the binary quintic  $(a, b, c, d, e, f)(x, y)^5$ .

Ex. 31. Prove that this covariant of the quintic is, but for sign, the same as the canonizant (Ex. 20).

Ans. Show that the form of Ex. 30 multiplied by

$$\begin{vmatrix} 1, & 0, & 0, & 0 \\ x, & y, & 0, & 0 \\ 0, & x, & y, & 0 \\ 0, & 0, & x, & y \end{vmatrix}$$

is the form of Ex. 20 multiplied by  $-y^3$ .

18.] **Intermediate invariants and covariants.** From a given invariant or covariant of a quantic can always be derived a series of invariants or covariants, as the case may be, of two or more quantics of the same order in the same variables. The method may be illustrated by the deduction of the result of Ex. 4 (§ 7) from that of Ex. 1.

By the substitution  $x = lX + mY$ ,  $y = l'X + m'Y$  let  $ax^2 + 2bxy + cy^2$  and  $a'x^2 + 2b'xy + c'y^2$  be transformed into  $AX^2 + 2BXY + CY^2$  and  $A'X^2 + 2B'XY + C'Y^2$  respectively. Then, whatever constant  $k$  be,

$$(a + ka')x^2 + 2(b + kb')xy + (c + kc')y^2$$

is transformed into  $(A + kA')X^2 + 2(B + kB')XY + (C + kC')Y^2$ .

Consequently, by Ex. 1,

$$\begin{aligned} (A + kA')(C + kC') - (B + kB')^2 \\ = M^2\{(a + ka')(c + kc') - (b + kb')^2\}; \end{aligned}$$

i. e.

$$\begin{aligned} AC - B^2 + k(AC' + A'C - 2BB') + k^2(A'C' - B'^2) \\ = M^2\{ac - b^2 + k(ac' + a'c - 2bb') + k^2(a'c' - b'^2)\}. \end{aligned}$$

This is true for all values of  $k$ . The multipliers of different powers of  $k$  on the two sides must then be separately equal each to each. Accordingly

$$\begin{aligned} AC - B^2 &= M^2(ac - b^2), \\ AC' + A'C - 2BB' &= M^2(ac' + a'c - 2bb'), \\ A'C' - B'^2 &= M^2(a'c' - b'^2). \end{aligned}$$

Of these three equalities the first and third are merely expressive of the fact of invariancy from which we started. The second however gives us the additional fact that

$$ac' + a'c - 2bb'$$

is an invariant of the quadratics

$$ax^2 + 2bxy + cy^2, a'x^2 + 2b'xy + c'y^2$$

jointly. It is said to be the invariant *intermediate* between  $ac - b^2$  and  $a'c' - b'^2$ .

This result is one of great historic interest. With Boole's discovery of it in 1841 the era of systematic investigation in the algebra of invariants began. In his original memoir (*Cambridge Math. Journal*, Vol. III) he showed how to find from any discriminant the intermediate invariants between the discriminants of two quantics of the same kind and order.

For another well-known example of the method reference may be made to the investigation (Salmon's *Conic Sections*, § 370) of the intermediate invariants  $\Theta$ ,  $\Theta'$  between the discriminants  $\Delta$ ,  $\Delta'$  of two conics (ternary quadratics).

19.] The method is clearly one of perfectly general application when we are given any invariant or covariant whatever of any quantic whatever. Let  $P$  be any invariant or any covariant of a  $q$ -ary  $p$ -ic in which the coefficients are  $a, b, c, \dots$  and the variables  $x, y, z, \dots$ . Consider also another  $q$ -ary  $p$ -ic, in the same variables, whose coefficients in the same order are  $a', b', c', \dots$ . Put for  $a, b, c, \dots$ , in  $P$ ,  $a + ka'$ ,  $b + kb'$ ,  $c + kc'$ ,  $\dots$ , and expand in powers of  $k$ . The multiplier of every power of  $k$  in the result is an invariant or covariant, as the case may be, of the two  $q$ -ary  $p$ -ics, the factor, which is a function of the constants in the scheme of linear substitution, in the relation expressive of the fact of invariancy or covariancy being the same as that in the relation which expresses the fact of invariancy or covariancy of  $P$ . The multiplier of the highest power of  $k$  which occurs is  $P'$ , the result of replacing  $a, b, c, \dots$  by  $a', b', c', \dots$  in  $P$ , and the multipliers of other powers of  $k$  are invariants, or covariants, *intermediate* between  $P$  and  $P'$ .

The general form of the invariants or covariants thus derived from  $P$  is

$$\frac{1}{r!} \left( a' \frac{d}{da} + b' \frac{d}{db} + c' \frac{d}{dc} + \dots \right)^r P,$$

for this is, by Taylor's theorem, the coefficient of  $k^r$ . Or, again, it may be written

$$\frac{1}{(i-r)!} \left( a \frac{d}{da'} + b \frac{d}{db'} + c \frac{d}{dc'} + \dots \right)^{i-r} P,$$

where  $i$  is the degree in the coefficients  $a, b, c, \dots$  of  $P$ . The values  $1, 2, 3, \dots, i-1$  of  $r$  give the intermediates between  $P$  and  $P'$ . The values  $0$  and  $i$  give  $P$  and  $P'$  respectively. Greater values of  $r$  than  $i$  are unproductive, for the differential operation  $a' \frac{d}{da} + b' \frac{d}{db} + c' \frac{d}{dc} + \dots$  repeated more than  $i$  times annihilates  $P$ .

In like manner invariants and covariants of systems of more than two quantics of the same order in the same variables are derived from invariants and covariants  $P$  of a single quantic of that type. We have only to put in  $P$ , for  $a, a+k_1a_1+k_2a_2+\dots$ , for  $b, b+k_1b_1+k_2b_2+\dots$ , and similarly for  $c, d, \dots$ , to expand according to powers and products of powers of  $k_1, k_2, \dots$ , and to take the multipliers of these powers and products separately. We thus obtain that, for any positive integral or zero values of  $r_1, r_2, r_3, \dots$  whose sum lies between  $0$  and  $i$ ,

$$\left(a_1 \frac{d}{da} + b_1 \frac{d}{db} + \dots\right)^{r_1} \left(a_2 \frac{d}{da} + b_2 \frac{d}{db} + \dots\right)^{r_2} \left(a_3 \frac{d}{da} + b_3 \frac{d}{db} + \dots\right)^{r_3} \dots P$$

is an invariant or covariant of the system of  $q$ -ary  $p$ -ics whose coefficients in the same order are  $a, b, c, \dots; a_1, b_1, c_1, \dots; a_2, b_2, c_2, \dots; a_3, b_3, c_3, \dots; \dots$ , according as  $P$  is an invariant or covariant of the first  $q$ -ary  $p$ -ic. The corresponding invariants or covariants  $P_1, P_2, \dots$  of the second, third, &c.  $q$ -ary  $p$ -ics, as well as their intermediates, and the corresponding invariants or covariants of triads, &c. of  $q$ -ary  $p$ -ics chosen from among the entire system, are all included.

20.] The method admits of a limited application to quantics of different orders in the same variables; namely to the case when the order of one quantic is a multiple of the order of every other quantic of the system. For instance, if two quantics  $u, v$  in the same variables be of orders  $p', p$  respectively, and if  $a, b, c, \dots$  are the coefficients in  $u$  and  $\alpha, \beta, \gamma, \dots$  the corresponding coefficients in  $v^{p'}$ , then the functions

$$\left(\alpha \frac{d}{da} + \beta \frac{d}{db} + \gamma \frac{d}{dc} + \dots\right)^{r} P,$$

where  $P$  is any invariant or covariant of  $u$ , are invariants or covariants of  $u$  and  $v^{p'}$ , and therefore of  $u$  and  $v$ .

Ex. 32. From the invariant  $ae - 4bd + 3c^2$  of the quartic

$$(a, b, c, d, e) (x, y)^4$$

obtain an invariant of that quartic and the quadratic  $(a', b', c') (x, y)^2$  of the first degree in the coefficients of the quartic and of the second in those of the quadratic.

Ans.  $a'^2e - 4a'b'd + 2(a'c' + 2b'^2)c - 4b'c'b + c'^2a$ . Factor  $M^4$ .

Ex. 33. If  $P$  be an invariant or covariant of  $(a_0, a_1, a_2, \dots, a_p) (x, y)^p$ , prove that the functions

$$\left( \xi^p \frac{d}{da_0} + \xi^{p-1} \eta \frac{d}{da_1} + \xi^{p-2} \eta^2 \frac{d}{da_2} + \dots + \eta^p \frac{d}{da_p} \right)^r P,$$

for values of  $r$  between 1 and  $i - 1$  inclusive, where  $i$  is the degree of  $P$  in  $a_0, a_1, a_2, \dots, a_p$ , are invariants of the  $p$ -ic and the linear form  $\xi x + \eta y$  jointly.

The importance of *evectants*, as the functions obtained in this manner from invariants are called, will be seen hereafter.

Ex. 34. From any invariant or covariant of several quantities of the same order in the same variables the operation

$$a' \frac{d}{da} + b' \frac{d}{db} + c' \frac{d}{dc} + \dots,$$

repeated till a vanishing result is obtained, produces a series of invariants or covariants, as the case may be. Here  $a, b, c, \dots$  and  $a', b', c', \dots$  are corresponding coefficients in any two of the quantities.

Ex. 35. The effect of replacing  $a', b', c', \dots$  by  $a, b, c, \dots$  in an invariant or covariant of two quantities  $u, v$  of the same order in the same variables, where  $a, b, c, \dots$  and  $a', b', c', \dots$  are corresponding coefficients in  $u$  and  $v$ , is to give an invariant or covariant of  $u$  alone, or else a vanishing result.



## CHAPTER II.

### ESSENTIAL QUALITIES OF INVARIANTS.

21.] In the present chapter we shall, at the expense of some repetition hereafter, confine our attention to invariants, reserving till the next the analogous consideration of covariants.

Except where otherwise stated, rational integral invariants are alone dealt with, the words 'rational integral' being as a rule omitted.

And first we consider invariants of a single quantic only.

Let us denote constantly by  $u$  the quantic under consideration, by  $p$  its order in the variables, by  $q$  the number of those variables, by small letters  $a, b, c, \dots, x, y, \dots$  the coefficients and variables in its original form, and by capitals  $A, B, C, \dots, X, Y, \dots$  the corresponding coefficients and variables in the transformed form to which it is reduced by a linear substitution. Also let us, except where otherwise stated, consider the scheme of linear substitution perfectly general as in § 2, and denote by  $l, m, \dots, l', m', \dots$  the assemblage of the coefficients of  $X, Y, \dots$  in the expressions for  $x, y, \dots$ . These coefficients we will speak of as the constants of the substitution, or of the transformation.

Taking the identical equality

$$F(A, B, \dots) = \phi(l, m, \dots, l', m', \dots) F(a, b, \dots),$$

which expresses that  $F(a, b, \dots)$  is an invariant of  $u$ , our immediate aim will be to prove

- (1) that  $F(a, b, \dots)$  is necessarily homogeneous, and
- (2) that  $\phi(l, m, \dots, l', m', \dots)$  is necessarily a power of the modulus  $M$  of the transformation, defined in § 2.

A knowledge of the first fact must precede a proof of the second.

22.] An invariant necessarily homogeneous in the coefficients. We shall speak of the dimensions of a homogeneous function of the coefficients in those coefficients as its *degree*<sup>1</sup>.

If possible let the invariant  $F(a, b, \dots)$  consist of a sum of parts

$$H_1(a, b, \dots) + H_2(a, b, \dots) + H_3(a, b, \dots) + \dots$$

of different degrees  $i_1, i_2, i_3, \dots$

Since  $F(a, b, \dots)$  is an invariant for all possible schemes of linear substitution, it is so of course for a particular scheme. Let us express the fact of invariancy for the scheme of substitution

$$x = \lambda X, y = \lambda Y, z = \lambda Z, \dots,$$

which, it is to be observed, has only the effect of multiplying the  $p$ -ic  $u$  by  $\lambda^p$  and replacing  $x, y, z, \dots$  by  $X, Y, Z, \dots$ . The coefficients  $A, B, C, \dots$  in the transformed  $p$ -ic have then in this case the values  $\lambda^p a, \lambda^p b, \lambda^p c, \dots$ . Any homogeneous function of degree  $i$  in them is accordingly  $\lambda^{ip}$  times the same function of  $a, b, c, \dots$ .

Thus, if  $\psi(\lambda)$  be the form taken by  $\phi(l, m, \dots l', m', \dots)$  for the particular substitution we are using, the identical equality expressive of the invariancy gives us

$$\begin{aligned} \lambda^{i_1 p} H_1(a, b, \dots) + \lambda^{i_2 p} H_2(a, b, \dots) + \lambda^{i_3 p} H_3(a, b, \dots) + \dots \\ = \psi(\lambda) \{ H_1(a, b, \dots) + H_2(a, b, \dots) + H_3(a, b, \dots) + \dots \}. \end{aligned}$$

This is an identity, true for all values of  $a, b, \dots$ . Consequently the terms of each degree in  $a, b, \dots$  on the left are the same as the corresponding terms in each case on the right. Hence we must have simultaneously

$$\lambda^{i_1 p} = \psi(\lambda),$$

$$\lambda^{i_2 p} = \psi(\lambda),$$

$$\&c., \&c.,$$

<sup>1</sup> I should have preferred to use the older term *order* for this characteristic. But the practice of speaking of a function (in particular of a covariant), whose dimensions are  $i$  in the coefficients and  $\varpi$  in the variables, as of *degree*  $i$  and *order*  $\varpi$  has of late become almost universal. While regretting this I feel bound to adopt it consistently throughout.

which are inconsistent if  $i_1, i_2, i_3, \dots$  are different. The supposition was therefore unsound, and the invariant  $F(a, b, \dots)$  is of the same degree  $i$  throughout.

The proof holds for irrational invariants.

23.] The factor a power of the modulus. The formulae of the general linear substitution

$$\left. \begin{aligned} x &= lX + mY + nZ + \dots, \\ y &= l'X + m'Y + n'Z + \dots, \\ z &= l''X + m''Y + n''Z + \dots, \\ &\dots \dots \dots \end{aligned} \right\} \dots (1)$$

may we know, by solution for  $X, Y, Z, \dots$ , be reversed and written

$$\left. \begin{aligned} X &= M^{-1} \left\{ \frac{dM}{dl} x + \frac{dM}{dl'} y + \frac{dM}{dl''} z + \dots \right\}, \\ Y &= M^{-1} \left\{ \frac{dM}{dm} x + \frac{dM}{dm'} y + \frac{dM}{dm''} z + \dots \right\}, \\ Z &= M^{-1} \left\{ \frac{dM}{dn} x + \frac{dM}{dn'} y + \frac{dM}{dn''} z + \dots \right\}, \\ &\dots \dots \dots \end{aligned} \right\} \dots (2)$$

$\frac{dM}{dl} = \text{minor det.}$

where  $M$  denotes the modulus

$$\begin{vmatrix} l, & m, & n, & \dots \\ l', & m', & n', & \dots \\ l'', & m'', & n'', & \dots \\ \dots & \dots & \dots & \dots \end{vmatrix},$$

and (cf. § 2) must not vanish.

Looking upon the formulae of substitution for  $x, y, z, \dots$  in terms of  $X, Y, Z, \dots$  as those of the standard substitution, we may speak of the formulae for  $X, Y, Z, \dots$  in terms of  $x, y, z, \dots$  as those of the reversed substitution. The reversal of the reversed substitution reproduces the standard substitution. The modulus of the reversed substitution is  $M^{-1}$ , the reciprocal of the modulus of the standard substitution, as immediately follows from the known fact (cf. Burnside and Panton's *Theory of Equations*, § 124) that the determinant reciprocal to a given determinant of  $q$  rows and  $q$  columns is its  $(q-1)$ th power.

D  $\frac{dM}{dl} = \dots = M^{-1} \dots$

Our present object is to prove that the factor  $\phi(l, m, \dots l', m', \dots)$  in the equality (§ 21) expressive of the fact of invariancy of  $F(a, b, \dots)$  is a power of  $M$ . We have seen in the last article that  $F(a, b, \dots)$  is homogeneous in  $a, b, \dots$ , and therefore  $F(A, B, \dots)$  homogeneous in  $A, B, \dots$ . Now  $A, B, \dots$  are homogeneous and of degree  $p$  in  $l, m, \dots l', m', \dots$ . For our quantic  $u$  is transformed from the form

$$ax^p + pbx^{p-1}y + \dots$$

to the form  $AX^p + pBX^{p-1}Y + \dots$

by the scheme (1) in which  $x, y, \dots$  are homogeneous and linear in  $l, m, \dots l', m', \dots$ , so that  $x^p, x^{p-1}y, \dots$  are homogeneous and of degree  $p$  in  $l, m, \dots l', m', \dots$ . Thus  $F(A, B, \dots)$ , being homogeneous, and of degree  $i$  say, in its arguments  $A, B, \dots$ , which are themselves all homogeneous and of degree  $p$  in  $l, m, \dots l', m', \dots$ , is itself homogeneous, and of degree  $ip$ , in  $l, m, \dots l', m', \dots$ . Seeing then that it is equal to

$$\phi(l, m, \dots l', m', \dots) F(a, b, \dots),$$

where the second factor  $F(a, b, \dots)$  is free from  $l, m, \dots l', m', \dots$ , we conclude that  $\phi(l, m, \dots l', m', \dots)$  is homogeneous and of degree  $ip$  in its arguments.

We now use the fact that the effect of the reversed substitution (2) is to bring the  $q$ -ary  $p$ -ic  $u$  back from its second form  $AX^p + \dots$  to its first  $ax^p + \dots$ . The invariant equality

$$F(A, B, \dots) = \phi(l, m, \dots l', m', \dots) F(a, b, \dots), \quad \dots (3)$$

applying as it does to all linear transformations of all  $q$ -ary  $p$ -ics, must hold when we interchange  $a, b, \dots$  and  $A, B, \dots$ , and replace  $l, m, \dots l', m', \dots$  by the corresponding coefficients in the scheme (2). Thus

$$\begin{aligned} & F(a, b, \dots) \\ &= \phi\left(M^{-1} \frac{dM}{dl}, M^{-1} \frac{dM}{dl'}, \dots M^{-1} \frac{dM}{dm}, M^{-1} \frac{dM}{dm'}, \dots\right) F(A, B, \dots) \\ &\hat{=} M^{-ip} \phi\left(\frac{dM}{dl}, \frac{dM}{dl'}, \dots \frac{dM}{dm}, \frac{dM}{dm'}, \dots\right) F(A, B, \dots), \quad \dots (4) \end{aligned}$$

in virtue of the homogeneity of degree  $ip$  possessed by the function  $\phi$ . Accordingly, by combination of (3) and (4), we arrive at the identity

$$\phi(l, m, \dots l', m', \dots) \phi\left(\frac{dM}{dl}, \frac{dM}{dl'}, \dots \frac{dM}{dm}, \frac{dM}{dm'}, \dots\right) = M^{ip} \dots (5)$$

Thus  $M^{ip}$  breaks up into two rational integral factors, of which  $\phi(l, m, \dots l', m', \dots)$  is one. But (§ 14)  $M$  has no factors but unity and itself. Consequently  $\phi(l, m, \dots l', m', \dots)$  is a power of  $M$ , or a numerical multiple of such a power.

Suppose then that

$$\phi(l, m, \dots l', m', \dots) = kM^r.$$

By (5) it follows that

$$\phi\left(\frac{dM}{dl}, \frac{dM}{dl'}, \dots, \frac{dM}{dm}, \frac{dM}{dm'}, \dots\right) = \frac{1}{k} M^{ip-r}.$$

But  $\frac{dM}{dl}, \frac{dM}{dl'}, \dots, \frac{dM}{dm}, \frac{dM}{dm'}, \dots$  are all of  $q-1$  dimensions in  $l, m, \dots l', m', \dots$ , so that the dimensions in  $l, m, \dots l', m', \dots$  of the second  $\phi$  are  $q-1$  times those of the first. Hence

$$ip - r = (q-1)r,$$

i. e. 
$$r = \frac{ip}{q}.$$

Accordingly the equality expressive of the fact that  $F(a, b, \dots)$  is an invariant is of the form

$$F(A, B, \dots) = kM^{\frac{ip}{q}} F(a, b, \dots),$$

where  $k$  is a numerical constant. That this constant is necessarily unity we see at once by application to the case of the substitution

$$x = X, y = Y, z = Z, \dots,$$

for which  $A, B, \dots$  are the same as  $a, b, \dots$ , and  $M = 1$ .

We have proved, then, completely that if  $F(a, b, \dots)$  is an invariant of a  $q$ -ary  $p$ -ic it is necessarily homogeneous, and that, if its degree is  $i$ , the identity expressive of the fact of its invariancy is

$$F(A, B, \dots) = M^{\frac{ip}{q}} F(a, b, \dots).$$

The proof holds for irrational invariants, if we raise the two sides of (5), before reasoning from that equivalence, to such a power  $\mu$  as to make  $\mu ip$ , the index of the power of  $M$ , an integer.

24.] A consequence of this last conclusion is that,  $i$  being

the *degree* in the coefficients of any rational integral invariant,  $\frac{ip}{q}$  must necessarily be integral. For the left-hand member  $F(A, B, \dots)$ , when expressed in terms of  $a, b, \dots$  and  $l, m, \dots l', m', \dots$ , is rational and integral in  $l, m, \dots l', m', \dots$  as well as in  $a, b, \dots$ . So too must the right-hand member be. Thus  $M^{\frac{ip}{q}}$  is rational in  $l, m, \dots l', m', \dots$ . But  $M$  is not a power of any rational function, seeing that it has no factors but unity and itself. Hence  $\frac{ip}{q}$  is an integer.

The particular form which this conclusion takes when  $q = 2$ , i.e. for the case of binary quantics, should be at once noticed. It is that  $i$  and  $p$  cannot both be odd. Hence the theorem :

*No binary quantic of odd order can have any invariant of odd degree.*

In the next few articles an interpretation will be given to the integer  $\frac{ip}{q}$ , first in the case  $q = 2$  of binary quantics, and afterwards generally.

It will be seen, in fact, that there is another characteristic which is constant throughout an invariant, and equal to this integer; namely, its *weight*.

25.] **Weight.** In the binary  $p$ -ic

$$(a_0, a_1, a_2, \dots a_p) (x, y)^p$$

we have, as is usually done, given to every coefficient a suffix equal to the defect below the order  $p$  of the index of the power of  $x$  which it multiplies.

This suffix is, it will be remembered, in each case equal to the dimensions, in the roots of the equation in  $x : y$  obtained by equating the  $p$ -ic to zero, of the symmetric function of the roots which is equal to the ratio of the coefficient in question to the first coefficient  $a_0$ . Or, if we choose, as we may, to regard  $a_0$  as merely denoting a number  $a_0$  of abstract units, and so as being of no dimensions in the roots, we may say that the suffix attached to every coefficient exactly measures the dimensions in the roots of that coefficient. The suffix or degree in the roots of a coefficient is designated its *weight*.

The *weight* of any product of coefficients is the sum of the weights of its various factors, i. e. the sum of their suffixes, and measures the dimensions in the roots of the product in question. A repeated factor in a product must be reckoned as many times as it is repeated in estimating the product's weight. Thus, for instance, the product  $a_r^\rho a_s^\sigma a_t^\tau \dots$  is of weight  $\rho r + \sigma s + \tau t + \dots$ .

An invariant of degree  $i$  of a binary  $p$ -ic has been proved to be homogeneous, i. e. to consist of a sum of positive and negative numerical multiples of products of  $i$  factors chosen from among  $a_0, a_1, a_2, \dots, a_p$ , repeated factors being allowed. The theorem now to be established is that all these products have the same weight  $\frac{1}{2} ip$ .

A function which is thus of one weight throughout is said to be *isobaric*.

26.] **An invariant of a binary quantic is isobaric.** Apply to

$$(a_0, a_1, a_2, \dots, a_p)(x, y)^p$$

the particular linear substitution  $x = X, y = \lambda Y$ , of which the modulus is

$$\begin{vmatrix} 1, & 0 \\ 0, & \lambda \end{vmatrix} = \lambda.$$

This transforms the quantic into

$$(a_0, a_1\lambda, a_2\lambda^2, \dots, a_p\lambda^p)(X, Y)^p.$$

Consequently, if  $F(a_0, a_1, a_2, \dots, a_p)$  be an invariant of degree  $i$ , the identity expressive of the fact, viz.

$$F(A_0, A_1, A_2, \dots, A_p) = M^{\frac{1}{2}ip} F(a_0, a_1, a_2, \dots, a_p),$$

tells us that

$$F(a_0, a_1\lambda, a_2\lambda^2, \dots, a_p\lambda^p) = \lambda^{\frac{1}{2}ip} F(a_0, a_1, a_2, \dots, a_p).$$

The right-hand member here is entirely of degree  $\frac{1}{2} ip$  in  $\lambda$ . So therefore must the left be. Now the term on the left corresponding to a term

$$a_r^\rho a_s^\sigma a_t^\tau \dots \text{ in } F(a_0, a_1, a_2, \dots, a_p)$$

is  $(a_r\lambda^r)^\rho (a_s\lambda^s)^\sigma (a_t\lambda^t)^\tau \dots,$

i. e.  $a_r^\rho a_s^\sigma a_t^\tau \dots \lambda^{\rho r + \sigma s + \tau t + \dots}$ .

Consequently for every such term

$$\rho r + \sigma s + \tau t + \dots = \frac{1}{2}ip.$$

Thus  $F(a_0, a_1, a_2, \dots, a_p)$  is isobaric throughout, the constant weight of its terms such as  $a_r^\rho a_s^\sigma a_t^\tau \dots$  being  $\frac{1}{2}ip$ .

This applies even when the invariant is irrational, for an irrational invariant may be expressed as a sum, not necessarily finite, of terms to which the reasoning may be applied.

Ex. 1. If  $p = 2n$  or  $2n + 1$  there is no term in any invariant of the binary  $p$ -ic which has not at least one of  $a_0, a_1, a_2, \dots, a_n$  for a factor.

Ex. 2. Every invariant vanishes for a binary  $p$ -ic which has a linear factor raised to the  $r$ th power if  $2r > p$ . (*Cayley.*)

*Ans.* Take the linear factor for  $Y$ .

27.] **Weight generalized.** A like method and the analogous conclusion apply in general to a quantic in  $q$  variables. Of these variables call one, singled out as the last,  $\omega$ , and the others  $x, y, z, \dots$ .

In our  $q$ -ary  $p$ -ic let the suffix given to each coefficient be the index of the power of  $\omega$  which it multiplies. Thus, for instance,

the coefficients of $x^p, y^p, z^p, x^{p-1}y, y^{p-1}z, \dots$	have the suffix 0,
" $x^{p-1}\omega, y^{p-1}\omega, z^{p-1}\omega, x^{p-2}z\omega, \dots$	" 1,
" $x^{p-2}\omega^2, y^{p-2}\omega^2, x^{p-3}y\omega^2, y^2z^{p-4}\omega^2, \dots$	" 2,
" $\dots \dots \dots$	"
" $x\omega^{p-1}, y\omega^{p-1}, z\omega^{p-1}, \dots$	" $p-1,$

and the coefficient of  $\omega^p$  has the suffix  $p$ .

Our definition of *weight* is that every coefficient is of weight measured by its suffix, and that every product of coefficients is of weight measured by the sum of the suffixes of its various factors.

Our ideas of the import of weight according to this definition are made more definite by supposing that the result of equating our  $q$ -ary  $p$ -ic to zero is a relation in  $q-1$  quantities of the same kind,  $x : \omega, y : \omega, z : \omega, \dots$ . To be intelligible, and not imply more relations than one, it must be of



the same dimensions throughout in that kind of quantity. For this to be the case the coefficients which multiply products of  $p$  factors  $x, y, z, \dots$  without  $\omega$ , those which multiply products of  $\omega$  and  $p-1$  factors  $x, y, z, \dots$ , those which multiply  $\omega^2$  and  $p-2$  factors  $x, y, z, \dots$ , and so on, must be of dimensions in that kind of quantity which form an ascending arithmetic progression of common difference unity. If then, as implies no real loss of generality, we choose to regard the first class of coefficients as of no dimensions in the kind of quantity, the dimensions of the other classes will be  $1, 2, 3, \dots, p$  respectively. In other words, the dimensions of the various coefficients are measured by the suffixes assigned according to the convention from which we started. The idea of such dimensions is then identical with that of weight.

28.] **All invariants isobaric.** We can now prove the constancy and equality to  $\frac{ip}{q}$  of the weight, defined as above, for all terms of an invariant of a  $q$ -ary  $p$ -ic.

Transform the quantic by the substitution

$$x = X, y = Y, z = Z, \dots, \omega = \lambda \Omega,$$

which leaves every variable unaltered except  $\omega$ . Its modulus is  $\lambda$ .

The coefficients in the transformed quantic are at once seen to be the same as those in the untransformed, except that those with suffixes  $0, 1, 2, 3, \dots, p$  are multiplied by  $1, \lambda, \lambda^2, \lambda^3, \dots, \lambda^p$  respectively. Thus, if  $F(a_0, b_0, \dots, a_1, b_1, \dots, a_2, b_2, \dots, a_p)$  be an invariant of degree  $i$ ,

$$\begin{aligned} F(a_0, b_0, \dots, a_1\lambda, b_1\lambda, \dots, a_2\lambda^2, b_2\lambda^2, \dots, a_p\lambda^p) \\ = \lambda^{\frac{ip}{q}} F(a_0, b_0, \dots, a_1, b_1, \dots, a_2, b_2, \dots, a_p). \end{aligned}$$

Here the left-hand member must be, like the right, a multiple of a single power, the  $\frac{ip}{q}$ th, of  $\lambda$ . The index of every power of  $\lambda$  which occurs as multiplying a product in the expanded left, and consequently the weight of every product of coefficients in  $F$ , must therefore be constant and equal to  $\frac{ip}{q}$ .

This applies even when the invariant is irrational.

Ex. 3. Every term in any invariant of a  $q$ -ary  $p$ -ic must contain at least one factor with a suffix less than  $r$  if  $qr > p$ .

Ex. 4. No quadratic in more than two variables can have any invariant which does not vanish when the quadratic breaks up into two linear factors.

Ex. 5. Every term in any invariant of a  $q$ -ary  $p$ -ic must contain at least one factor with a suffix greater than  $r$  if  $qr < p$ .

29.] **Absolute invariants.** For integral invariants the degree  $i$ , and consequently the weight  $\frac{ip}{q}$ , are essentially positive and different from zero. Thus the power of  $M$  in the equality expressive of invariancy

$$F(A, B, \dots) = M^{\frac{ip}{q}} F(a, b, \dots)$$

is essentially a positive power. We cannot then discover any integral function of the coefficients of a quantic which is what is called an *absolute invariant*, that is to say a function of the coefficients which is absolutely equal to the same function of the coefficients in the transformed quantic. For an absolute invariant the power of  $M$  above would have to be  $M^0$ , or the degree  $i$ , and consequently the weight  $\frac{ip}{q}$ , would have to be zero.

If, however, a quantic have two or more distinct integral invariants, i.e. two invariants which are not powers of the same invariant, it will have one or more absolute fractional invariants. For, if  $F_1(a, b, \dots)$  and  $F_2(a, b, \dots)$  are two invariants of the same degree  $i$  of a  $q$ -ary  $p$ -ic, we have

$$F_1(A, B, \dots) = M^{\frac{ip}{q}} F_1(a, b, \dots),$$

and 
$$F_2(A, B, \dots) = M^{\frac{ip}{q}} F_2(a, b, \dots);$$

so that 
$$\frac{F_1(A, B, \dots)}{F_2(A, B, \dots)} = \frac{F_1(a, b, \dots)}{F_2(a, b, \dots)},$$

which shows that the ratio of  $F_1$  to  $F_2$  is an absolute invariant. Again, if  $F_1(a, b, \dots)$  and  $F_2(a, b, \dots)$  are of different degrees  $i_1, i_2$ , let  $k$  be the L. C. M. of  $i_1$  and  $i_2$ . Then  $F_1^{\frac{k}{i_1}}$  and

$F_2^{\frac{k}{2}}$  are two distinct invariants of the same degree  $k$ , and their ratio  $F_1^{\frac{k}{2}} \div F_2^{\frac{k}{2}}$  is an absolute invariant.

For instance, we have seen (§ 7, Ex. 5) that

$$I \equiv ae - 4bcd + 3c^2$$

is an invariant of the binary quartic  $(a, b, c, d, e) (x, y)^4$ . Its degree is 2 and its weight 4, which is rightly equal to  $\frac{2 \cdot 4}{2}$ .

We have also seen (§ 17, Ex. 19) that the same quartic has another invariant

$$J \equiv ace + 2bcd - ad^2 - b^2e - c^3$$

of degree 3 and weight 6.  $I^3$  and  $J^2$  are then both of degree 6 and weight 12, and are distinct from one another. If then  $I'$  and  $J'$  are the same functions of the coefficients in the quartic obtained from the given quartic by a linear substitution for  $x$  and  $y$  as  $I$  and  $J$  are of the coefficients in the given quartic,

$$\frac{I'^3}{J'^2} = \frac{M^{12} I^3}{M^{12} J^2} = \frac{I^3}{J^2},$$

so that  $I^3 J^{-2}$  is an absolute invariant of the binary quartic.

30.] **Limit to the number of independent invariants.** A binary  $p$ -ic has  $p-3$  independent absolute invariants, if  $p$  exceed 3, and none if  $p$  do not exceed 3. The first part of this statement is one which cannot well be proved at the present stage; but it may be seen as follows that  $p-3$  is a superior limit which the number of independent absolute invariants cannot exceed.

Let  $(A_0, A_1, A_2, \dots, A_p) (X, Y)^p$  be the transformed quantic obtained from  $(a_0, a_1, a_2, \dots, a_p) (x, y)^p$  by the linear substitution

$$x = lX + mY, y = l'X + m'Y.$$

Its coefficients  $A_0, A_1, A_2, \dots, A_p$  are at once expressed as  $p+1$  functions of  $a_0, a_1, a_2, \dots, a_p$  and the four letters  $l, m, l', m'$ . If  $p$  do not exceed 3 it is impossible to eliminate  $l, m, l', m'$ , and obtain a relation connecting  $A_0, A_1, \dots, A_p$  with  $a_0, a_1, \dots, a_p$  alone. If, however,  $p$  exceed 3 it is possible, by elimination

of  $l, m, l', m'$ , to obtain  $p-3$  independent relations which must subsist between  $A_0, A_1, A_2, \dots, A_p$  and  $a_0, a_1, a_2, \dots, a_p$ , but no more. If, as is in fact the case, these  $p-3$  relations can be thrown into such a form as to express  $p-3$  equalities of functions of  $a_0, a_1, a_2, \dots, a_p$  to the same functions respectively of  $A_0, A_1, A_2, \dots, A_p$ , those  $p-3$  functions are absolute invariants; but there cannot be more than that number which are independent.

It now follows that if  $p$  do not exceed 3 there cannot be two independent invariants which are not absolute, and that if  $p$  exceed 3 there cannot be more than  $p-2$  which are independent. For, as seen in the preceding article, any two independent invariants determine an absolute invariant, so that two, or more than  $p-2$ , independent invariants would determine one, or more than  $p-3$ , independent absolute invariants.

We must not, however, form the erroneous conclusion that, when  $p-2$  independent rational integral invariants have been discovered, every other rational integral invariant can be expressed as a *rational integral* function of these  $p-2$ . The system of  $p-2$  invariants is *algebraically* complete, but another may be a function of them, as it must be, without being a rational integral function of them. For binary quantics of the first four orders there are, as a matter of fact, algebraically complete systems, 0, 1, 1, 2 in number, in terms of which all other invariants can be rationally and integrally expressed, but for the fifth, sixth, &c., orders there is no corresponding simplicity. For instance, the binary quintic has 3 ( $= p-2$ ) independent invariants of degrees 4, 8, 12, and these are the invariants of lowest degrees which it possesses. They form an algebraically complete system. But there is another invariant of the quintic of degree 18. This must be a function of the three first, but it is perfectly clear that it cannot be a rational integral function of them, for the degree 18, which is not divisible by 4, cannot be expressed as a sum of multiples of degrees chosen from 4, 8, 12, which are all divisible by 4. It is found to be the square root of a rational integral function of the three. Because it cannot be expressed rationally and integrally in terms of *irreducible* invariants of lower degrees, it is said to be itself *irreducible*.

That the number of irreducible invariants of a binary  $p$ -ic is finite for all values of  $p$  is a proposition of some difficulty which was first established by Gordan. The number, though finite, is not known to follow any simple law for all values of  $p$ . A proof of the finiteness due to Hilbert will be given in a later chapter.

31.] **Invariants of two or more quantics.** So far in this chapter we have been dealing with invariants of a single quantic only. With regard to invariants of a system consisting of two or more quantics in the same variables the methods of §§ 22 to 28 establish with equal ease the following theorems.

(1) In any invariant of  $r$  quantics of orders  $p_1, p_2, \dots, p_r$  in the same  $q$  variables, the sum

$$\Sigma (ip) = i_1 p_1 + i_2 p_2 + \dots + i_r p_r$$

is constant for all terms,  $i_1, i_2, \dots, i_r$  being the degrees of any term in the coefficients of the various quantics respectively.

This is established as in § 22.

(2) The factor, depending on the constants of the transformation only, by which the invariant has to be multiplied to make it equal to the same function of the coefficients in the transformed quantics, is  $M^c$ , where  $M$  is the modulus, and

$$w = \frac{1}{q} \Sigma (ip).$$

This is established as in § 23.

(3) The whole weight, i.e. the sum of the  $r$  weights in the sets of coefficients of the  $r$  quantics, is the same for every term of the invariant, and equal to  $w$  the index of the power of  $M$  in (2).

This is established as in §§ 26, 28.

It also follows that, for a rational integral invariant, the sum  $\Sigma(ip)$  is necessarily divisible by  $q$ ; for the weight, a sum of integers, must be integral.

32.] It will be observed that there is nothing in these conclusions to prevent our contemplating the existence of invariants of two or more quantics, which, though isobaric (i.e. of constant weight throughout), are not homogeneous,

either in the sets of coefficients of the various quantic separately, or on the whole. Nothing in the above indicates that  $i_1, i_2, \dots, i_r$  are constant throughout the invariant, or even that  $\Sigma i$  is so.

To contemplate such non-homogeneous invariants is, however, unnecessary, for the different parts of such an invariant, which are homogeneous on the whole and also separately in the coefficients of every quantic of the system, are separately invariants.

The proof of this may with ease be stated generally. It will perhaps be made all the clearer by considering an example only.

Suppose the fact to have been noticed that

$$\{ab'^3 - 3ba'b'^2 + 3ca'^2b' - db'^3\}^4 + \{(ac - b^2)b'^2 - (ad - bc)a'b' + (bd - c^2)a'^2\}^3$$

is an invariant of the binary cubic and linear forms

$$ax^3 + 3bx^2y + 3cxy^2 + dy^3, \\ a'x + b'y,$$

i.e. that, denoting as usual coefficients in the transformed quantities by capitals,

$$\{AB'^3 - \dots\}^4 + \{(AC - B^2)B'^2 - \dots\}^3 \\ = M^{12}[\{ab'^3 - \dots\}^4 + \{(ac - b^2)b'^2 - \dots\}^3]. \quad \dots(1)$$

The invariant consists of a part of degree 4 in the coefficients of the cubic and 12 in those of the linear form, and a part of degree 6 in the coefficients of each form.

Now  $A, B, C, D$  are of the first degree in  $a, b, c, d$ , and  $A', B'$  of the first degree in  $a', b'$ , involving besides, in each case, the constants  $l, m, l', m'$  of the transformation only. The left-hand member of (1) contains then like the right terms of partial degrees 4, 12, and terms of partial degrees 6, 6. Consequently, the equality being an identity holding whatever  $a, b, c, d, a', b'$  are, the terms of partial degrees 4, 12 on the left and right must be equal, and also those of partial degrees 6, 6. In other words,

$$\{ab'^3 - 3ba'b'^2 + 3ca'^2b' - da'^3\}^4 \\ \text{and} \quad \{(ac - b^2)b'^2 - (ad - bc)a'b' + (bd - c^2)a'^2\}^3$$

are invariants separately.

A single quantic has, we know (§ 22), homogeneous invariants only.

33.] We lose then no completeness by considering only those invariants of two or more quantics which are homogeneous in the different sets of coefficients separately as fundamental. Non-homogeneous invariants are linear functions of such homogeneous invariants as have the same whole weight. Thus with regard to a complete system of invariants of two or more quantics we have the conclusions:—

(1) That they are homogeneous in the coefficients of every quantic of the system separately, so that also, if  $i_1, i_2, \dots, i_r$  be the degrees of any invariant in these sets of coefficients, the whole degree is constant, viz.

$$i = i_1 + i_2 + \dots + i_r;$$

(2) that they are isobaric *on the whole*, any one being of weight

$$w = \frac{1}{q} (i_1 p_1 + i_2 p_2 + \dots + i_r p_r).$$

(N.B.—There is no reason to expect them to be isobaric in the coefficients of the quantics separately.)

(3) That the factor which has to multiply an invariant to produce the same function of the coefficients in the linearly transformed quantics is  $M^w$ .

## CHAPTER III.

### ESSENTIAL QUALITIES OF COVARIANTS.

34.] IN accordance with the remark in § 4, the consideration of covariants which are rational and integral both in the coefficients and variables is fundamental. By the word 'covariant' we, as a rule, mean 'rational integral covariant.' The conclusions which follow apply for the most part also to covariants which are irrational or fractional, but this will be stated where it is important to observe that it is the case.

It is well in the first place to see that we may confine attention to covariants which are homogeneous in the variables—to covariant quantics, in fact.

35.] A covariant which is not homogeneous in the variables is a sum of other covariants which are homogeneous in them.

For in the relation  
 $f(A, B, \dots, X, Y, \dots) = \phi(l, m, \dots, l', m', \dots)f(a, b, \dots, x, y, \dots)$ ,  
which expresses that  $f(a, b, \dots, x, y, \dots)$  is a covariant, the terms of order  $\varpi$  in  $x, y, \dots$  on the right can produce, upon putting  $x = lX + mY + \dots$ ,  $y = l'X + m'Y + \dots$ , ..., terms of order  $\varpi$  only in  $X, Y, \dots$ ; and no other terms on the right can produce terms of order  $\varpi$  in  $X, Y, \dots$ . Consequently, the relation being an identity, these terms must be identical with the terms of order  $\varpi$  in  $X, Y, \dots$  on the left. In other words, if the covariant  $f$  is not homogeneous in  $x, y, \dots$ , its various parts of different orders in  $x, y, \dots$  are separately covariants.

This applies also to irrational and fractional covariants, which by expansion can be expressed as sums of parts, not necessarily finite in number, arranged according to their orders in the variables.

The proof deals equally with covariants of one and covariants of several quantics. In the next few articles for



greater clearness covariants of a single quantic are alone first considered.

36.] **Homogeneity in the coefficients.** By the *order* of a covariant, now regarded as homogeneous in the variables, is meant its order or degree in those variables. By *degree* is meant, as in the preceding chapter, degree in the coefficients<sup>1</sup>.

If possible let the covariant  $f(a, b, \dots, x, y, \dots)$ , of the same order  $\varpi$  throughout, be a sum of parts of different degrees  $i_1, i_2, i_3, \dots$ . Apply the identity expressive of the covariancy to the case of the particular linear substitution  $x = \lambda X, y = \lambda Y, \dots$ . As in § 22, the coefficients  $A, B, \dots$  in the transformed quantic are in this case  $\lambda^p a, \lambda^p b, \dots$ , while the variables  $X, Y, \dots$  in the transformed quantic are  $\lambda^{-1} x, \lambda^{-1} y, \dots$ . Thus if  $H_r$  be the aggregate of those terms in  $f(a, b, \dots, x, y, \dots)$  which are of degree  $i_r$ , and of order  $\varpi$ , the corresponding terms in  $f(A, B, \dots, X, Y, \dots)$  are  $\lambda^{i_r p - \varpi} H_r$ . Hence, by exactly the same argument as in § 22, if  $\psi(\lambda)$  be what  $\phi(l, m, \dots l', m', \dots)$  becomes for the particular values of  $l, m, \dots l', m', \dots$  which we are considering,  $\psi(\lambda)$  must be equal separately to  $\lambda^{i_1 p - \varpi}, \lambda^{i_2 p - \varpi}, \lambda^{i_3 p - \varpi}, \dots$ . The assumption that  $i_1, i_2, i_3, \dots$  are different is then untenable.

Thus, while we lose no real generality by requiring a covariant to be of constant order throughout, we are compelled also to require a covariant of a single quantic whose order is the same throughout to be of the same degree throughout.

Were we to prefer to deal with a covariant having parts of different orders  $\varpi_1, \varpi_2, \varpi_3, \dots$  as a single covariant, rather than as a sum of covariants of orders  $\varpi_1, \varpi_2, \varpi_3, \dots$ , our conclusion come to as above would be that the degrees  $i_1, i_2, i_3, \dots$  of those parts respectively are connected with their orders by the equalities

$$i_1 p - \varpi_1 = i_2 p - \varpi_2 = i_3 p - \varpi_3 = \dots$$

These conclusions apply to irrational and fractional covariants.

37.] **The factor a power of the modulus.**

The proof that the factor  $\phi(l, m, \dots l', m', \dots)$ , in the rela-

<sup>1</sup> See the footnote to § 22.

tion (§ 35) which expresses the fact of covariancy of a covariant, is a power of the modulus  $M$  proceeds exactly as in § 23. If  $\varpi$  be the order and  $i$  the degree of the covariant  $f(a, b, \dots x, y, \dots)$ , the power is the  $\frac{ip - \varpi}{q}$ th,  $ip - \varpi$  being now the degree of the left-hand side  $f(A, B, \dots, X, Y, \dots)$  in the constants of transformation  $l, m, \dots l', m', \dots$ , when it is expressed explicitly in terms of those constants and

$$a, b, \dots, x, y, \dots$$

Thus, if we adopt the notation  $K(a, b, \dots)^i (x, y, \dots)^\varpi$  to denote a covariant of degree  $i$  and order  $\varpi$ , the fact of its being a covariant is expressed by

$$K(A, B, \dots)^i (X, Y, \dots)^\varpi = M^{\frac{ip - \varpi}{q}} K(a, b, \dots)^i (x, y, \dots)^\varpi.$$

All this applies as well to irrational and fractional covariants as to those which are rational and integral.

If the covariant be rational and integral we can at once draw the conclusion, as in § 24, that the index  $\frac{ip - \varpi}{q}$  cannot be fractional. It is perhaps well, however, to adopt a different order, and by introduction of the idea of *weight* to ascertain first the import of the integer, or zero, to which it is equal.

38.] **Weight in the case of a binary quantic.** As in § 25 the weight of a coefficient in the binary  $p$ -ic

$$(a_0, a_1, a_2, \dots a_p) (x, y)^p$$

is its suffix. For present purposes we do best to say further that  $x$  and  $y$  have weights 1 and 0 respectively. This is in accordance with the idea developed in § 25 that weight measures dimensions in a suppositious kind of quantity of which  $\frac{x}{y}$  contains  $\frac{x}{y}$  units, and in which  $\frac{a_1}{a_0}, \frac{a_2}{a_0}, \dots \frac{a_p}{a_0}$ , being of 1, 2, ...  $p$  dimensions in the values of  $\frac{x}{y}$  which make the quantic vanish, are of 1, 2, ...  $p$  dimensions respectively.

With this enlarged conception of weight we may see as follows that  $K(a_0, a_1, \dots a_p)^i (x, y)^\varpi$ , a covariant of the binary  $p$ -ic, is of constant weight  $\frac{1}{2}(ip + \varpi)$  throughout.

As in § 26 take for scheme of linear substitution the particular one

$$x = X, y = \lambda Y,$$

of which the modulus  $M$  is  $\lambda$ . If

$$(A_0, A_1, \dots, A_p)(X, Y)^p$$

be the transformed quantic, the values of  $A_0, A_1, \dots, A_p$  are now  $a_0, a_1\lambda, \dots, a_p\lambda^p$ , and, as in § 26, every product of powers of  $A_0, A_1, \dots, A_p$  is the same product of powers of  $a_0, a_1, \dots, a_p$  multiplied by  $\lambda$  raised to a power whose index is the weight of the product. Moreover every product  $X^r Y^{\varpi-r}$  of powers of  $X$  and  $Y$  is equal to  $\lambda^{-\varpi+r} x^r y^{\varpi-r}$ , i.e. to the corresponding product of powers  $x^r y^{\varpi-r}$  multiplied by a power of  $\lambda$  whose index is the weight of the product diminished by  $\varpi$  its order. Thus in the identity

$$K(A_0, A_1, \dots, A_p)^i (X, Y)^\varpi = \lambda^{\frac{i\varpi - \varpi}{2}} K(a_0, a_1, \dots, a_p)^i (x, y)^\varpi,$$

every term on the left is, for this substitution, the corresponding term in  $K(a_0, a_1, \dots, a_p)^i (x, y)^\varpi$  multiplied by  $\lambda^{w-\varpi}$ , where  $w$  is the weight of the term. The identity then tells us that for every term

$$\lambda^{w-\varpi} = \lambda^{\frac{1}{2}(i\varpi - \varpi)},$$

so that  $w = \frac{1}{2}(i\varpi + \varpi)$  for all terms. A covariant is then *isobaric*.

So far this applies to irrational and fractional as well as to rational integral covariants.

39.] For rational integral covariants the weight is a sum of positive integers, and is therefore itself a positive integer.

Thus  $\frac{1}{2}(i\varpi + \varpi)$  is necessarily a positive integer.

It follows that the index of the power of  $M$  in the equality expressive of the covariancy of a rational integral covariant is integral, or zero, for it is

$$\frac{1}{2}(i\varpi - \varpi) = \frac{1}{2}(i\varpi + \varpi) - \varpi = w - \varpi,$$

i.e. is the excess of one positive integer over another.

Moreover it cannot be a negative integer. For,  $w$  being the weight of the covariant,  $w - \varpi$  is the weight of the coefficient of  $x^\varpi$  in the covariant, and this coefficient being

a rational integral function of  $a_0, a_1, a_2, \dots, a_p$ , whose weights are zero and positive, cannot have a negative weight.

This assumes however that in a covariant of order  $\varpi$  the term in  $x^\varpi$  must necessarily occur. This is the case. Were it otherwise the covariant would have  $y$  for a factor. Now were it possible for  $yF(a, b, \dots, x, y)$  to be a covariant we should have, for any linear substitution whatever,

$$YF(A, B, \dots, X, Y) = (lm' - l'm)^{\varpi} y F(a, b, \dots, x, y),$$

which would necessitate that the covariant  $yF(a, b, \dots, x, y)$  have  $Y$ , i.e.  $\frac{m'x - my}{lm' - l'm}$ , for a factor, whatever  $m', m$  and  $lm' - l'm$  be. Now this is an absurdity, for  $yF(a, b, \dots, x, y)$  has only  $\varpi$  linear factors.

From the fact that  $\frac{1}{2}(ip - \varpi)$  is integral, or zero, we draw at once the conclusion that  $ip$  and  $\varpi$  must be either both odd or both even. Hence arise the following theorems.

(1) *No binary quantic of even order  $p$  can have a covariant of odd order  $\varpi$ .*

(2) *No covariant of a binary quantic can be of even degree  $i$  (in the coefficients) and of odd order  $\varpi$  (in the variables).*

(3) *No covariant of a binary quantic of odd order  $p$  can be of odd degree  $i$  and even order  $\varpi$ .*

In particular, from (1) and (2) no covariant linear in the variables can belong to a binary quantic of even order, or be of even degree in the coefficients.

Ex. 1. Every term in every coefficient of any covariant of a binary  $p$ -ic must contain one or more of the first  $r$  coefficients  $a_0, a_1, \dots, a_{r-1}$  of the  $p$ -ic as a factor if  $2ir > ip + \varpi$ .

Ex. 2. Every covariant of degree  $i$  and order  $\varpi$  must vanish for a binary  $p$ -ic which has a linear factor raised to the  $r$ th power if  $2ir > ip + \varpi$ .

*Ans.* Take the factor for  $Y$ .

Ex. 3. Every term in the coefficients of  $x^\varpi, x^{\varpi-1}y, \dots, x^{\varpi-\rho+1}y^{\rho-1}$  in a covariant of order  $\varpi$  and degree  $i$  of a binary  $p$ -ic must contain at least one of  $a_0, a_1, a_2, \dots, a_{r-1}$  as a factor if  $ir - \rho < \frac{1}{2}(ip - \varpi)$ .

Ex. 4. If the coefficients in a binary  $p$ -ic have such values that the  $p$ -ic has a linear factor raised to the  $r$ th power, a covariant of degree  $i$  and order  $\varpi$  must have that factor to the  $\rho$ th power, where  $\rho = ir - \frac{1}{2}(ip - \varpi)$ . (*Cayley*.)

**Ex. 5.** If the degree  $i$  and order  $\varpi$  of a covariant of a binary  $p$ -ic be connected by the relation  $ip - \varpi = 0$ , show that the covariant can only be the  $i$ th power of the  $p$ -ic, or a numerical multiple of that  $i$ th power.

*Ans.* The coefficient of  $x^\varpi$ , i.e.  $x^{ip}$ , must be  $a^i$ , for its weight must be zero. Also by Ex. 4 the  $i$ th power of every linear factor of the  $p$ -ic must be a factor of the covariant.

Or thus. The  $i$ th power of the  $p$ -ic is a covariant; and there cannot be another covariant with  $a^i x^{ip}$  for its first term, as otherwise by subtraction a covariant with  $y$  for a factor could be formed.

**Ex. 6.** If the coefficient of the highest power of  $x$  in a covariant of the general binary  $p$ -ic is known, the order  $\varpi$  is determinate, and the covariant unique.

40.] **Weight in general.** With regard to a quantic in  $q$  variables  $x, y, z, \dots, \omega$  the estimation of weight explained in § 27 requires the supplementary idea that  $x, y, z, \dots$ , all the variables except the last one  $\omega$ , have weight unity, while  $\omega$  is of weight zero. This being so the weight of the  $q$ -ary  $p$ -ic is  $p$  throughout. The examination for weight of a covariant of degree  $i$  and order  $\varpi$  proceeds exactly as in § 38, by the method of § 28. The conclusion is that the weight  $w$  is constant throughout the covariant, being given by

$$\lambda^{w-\varpi} = \lambda^{\frac{ip-\varpi}{q}},$$

so that

$$w = \frac{ip + (q-1)\varpi}{q}.$$

This applies to covariants which are not rational and integral as well as to those which are. For rational integral covariants we have the further fact that  $w$  is a positive integer, and consequently that

$$w - \varpi = \frac{ip - \varpi}{q}$$

is an integer or zero. Moreover that it cannot be a negative integer is proved exactly as in § 39, by showing that the terms free from  $\omega$  in a covariant cannot all be absent.

**Ex. 7.** If the terms free from  $\omega$  in a covariant are known, the covariant is unique.

*Ans.* Otherwise a covariant with  $\omega$  for a factor could be formed.

Ex. 8. If  $\varpi = ip$ , so that  $w = \varpi$ , the coefficients of the terms free from  $\omega$  in a covariant involve only the coefficients of the terms free from  $\omega$  in the  $p$ -ic.

Ex. 9. In this case of  $\varpi = ip$ , the terms free from  $\omega$  in a covariant of a  $q$ -ary  $p$ -ic constitute a covariant of the  $(q-1)$ ary  $p$ -ic, obtained by replacing  $\omega$  by zero in that  $q$ -ary  $p$ -ic.

*Ans.* Apply a linear substitution which leaves  $\omega$  unaltered and expresses the other variables  $x, y, z, \dots, \psi$  linearly in terms of  $X, Y, Z, \dots, \Psi$ . The terms free from  $\omega$  are then transformed by a  $(q-1)$ ary substitution.

Ex. 10. Hence, by passing in succession to  $(q-2)$ ary,  $(q-3)$ ary, ... binary,  $p$ -ics, deduce from § 39, Ex. 5, that a covariant of the  $q$ -ary  $p$ -ic for which  $\varpi = ip$  has for its term in  $x$  alone  $\alpha^i x^{ip}$ , or a numerical multiple of this.

Ex. 11. Hence, by returning in succession from a binary, to a ternary, a quaternary, ... and at length a  $q$ -ary,  $p$ -ic, show from Ex. 5 and Ex. 7 that a covariant, of a  $q$ -ary  $p$ -ic, for which  $\varpi = ip$ , can be only the  $i$ th power of that  $q$ -ary  $p$ -ic, affected at most by a numerical multiplier.

41.] **Absolute covariants.** An *absolute* covariant is one which is exactly equal, without any factor which is even a power of  $M$ , to the same function of the coefficients and variables in the linearly transformed quantic. Thus, if the function  $K$  be an absolute covariant, we must have, in the identity

$$K(A, B, \dots)^i (X, Y, \dots)^\varpi = M^{\frac{ip-\varpi}{q}} K(a, b, \dots)^i (x, y, \dots)^\varpi,$$

$$\frac{ip-\varpi}{q} = 0, \quad \text{i.e.} \quad w - \varpi = 0.$$

Now  $w - \varpi$  is the weight of those coefficients in the covariant which multiply products of the variables whose weight is  $\varpi$ , i.e. products into which the last variable  $\omega$  does not enter. The only rational integral absolute covariants are then those in which the coefficients of products of the variables into which the last  $\omega$  does not enter are of zero weight. In particular, for a binary quantic, the coefficient of  $x^\varpi$  must be a function of zero weight of  $a_0, a_1, a_2, \dots, a_p$ , and so must be a mere power of  $a_0$ , or a numerical multiple of such a power. In § 39, Ex. 5, it has been seen that such a covariant can only be a numerical multiple of a power of the binary quantic of

which it is a covariant. And in § 40, Ex. 11, the corresponding fact has been given for quantics in general. Thus powers of quantics are the only rational integral absolute covariants of those quantics. Further light will be thrown on this fact in future chapters.

Fractional absolute covariants may, however, be seen to exist, as were fractional absolute invariants in § 29, whenever we have two or more distinct integral covariants, powers of the same ~~invariant~~ <sup>covariant</sup> not being regarded as distinct, for each of which  $ip - \varpi$  does not vanish. If, for instance,  $K$  and  $K'$  be two covariants of a  $q$ -ary  $p$ -ic, whose degrees are  $i, i'$  and orders  $\varpi, \varpi'$  respectively, and if  $\mu$  be the least common multiple of the integers  $\frac{ip - \varpi}{q}, \frac{i'p - \varpi'}{q}$ , then the ratio of

$$K^{\frac{\mu q}{ip - \varpi}} \text{ to } K'^{\frac{\mu q}{i'p - \varpi'}}$$

is an absolute covariant.

For example, it will be seen later (§ 45, Ex. 13) that the binary cubic

$$(a, b, c, d)(x, y)^3$$

has, besides its quadratic covariant (§ 7, Ex. 8)

$$(ac - b^2)x^2 + (ad - bc)xy + (bd - c^2)y^2,$$

a cubic covariant

$$(a^2d - 3abc + 2b^3)x^3 + \dots$$

For these two covariants  $\frac{1}{2}(ip - \varpi)$  has the values 2, 3 respectively. The cube of the first divided by the square of the second is then a fractional absolute invariant.

42.] **Limit to the number of independent covariants.** A limit to the possible number of independent covariants of a binary quantic may be found as follows.

In the equations of linear substitution

$$x = lX + mY,$$

$$y = l'X + m'Y,$$

in the expression for the modulus

$$M = lm' - l'm,$$

and in the  $p+1$  equations, in the two sets of coefficients and  $l, m, l', m'$ , which are obtained by expressing the identity of

$$(A_0, A_1, \dots, A_p) (X, Y)^p \text{ with } (a_0, a_1, \dots, a_p) (x, y)^p,$$

i.e. with  $(a_0, a_1, \dots, a_p) (lX + mY, l'X + m'Y)^p,$

we have altogether  $p+4$  relations connecting the old and new coefficients, the old and new variables, the modulus  $M$ , and  $l, m, l', m'$ . The elimination of these last four leaves exactly  $p$  independent relations as all that can connect only the old and new coefficients and variables and  $M$ .

For instance, the first three equations

$$x = lX + mY, y = l'X + m'Y, M = lm' - l'm,$$

suffice to determine three of  $l, m, l', m'$ , the last three say, in terms of the fourth  $l$  and  $x, y, X, Y, M$ , and lead to no relation free from  $l, m, l', m'$ . The expressions for  $m, l', m'$  inserted in the remaining  $p+1$  equations, produce from them  $p+1$  equations involving one unknown  $l$ , the old and new coefficients and variables, and  $M$ . By elimination of  $l$  from these, exactly  $p$  independent relations in coefficients and variables and  $M$  follow.

Now if there were more than  $p$  independent covariants, including the quantic itself, there would be more than  $p$  independent relations in coefficients and variables, old and new, and  $M$ ; viz. the more than  $p$  equalities of the several covariants, multiplied by proper powers of  $M$ , to the same functions of the new coefficients and variables. The number  $p$  is then a superior limit to the possible number of independent covariants of a binary  $p$ -ic.

In fact, a little more than this is true. The number  $p$  is, as the same reasoning shows, a superior limit to the possible number of independent invariants and covariants together, the quantic being regarded as a covariant of itself.

As a matter of fact  $p$  is not only a superior limit to the number of algebraically independent covariants and invariants, but the exact number of a complete system. The present however is not the stage at which to prove this important fact.

The warning of the latter part of § 30 should be repeated. When  $p$  covariants and invariants, algebraically independent of one another, are known, any other covariant or invariant



is a function of them. But this does not imply that, when  $p$  independent rational integral covariants and invariants are known, all others can be expressed as rational integral functions of them. There may be others that are *irreducible* in the sense of not being expressible as rational integral functions of simpler irreducible covariants and invariants; and except for the values 1 and 2 of  $p$  this is in fact the case. Thus for the binary cubic  $p = 3$ , but, when the three independent covariants and invariants, all covariants in fact, of lowest degrees have been found, there proves to be a fourth, an invariant, which, though of course a function of them, is irreducible in that it cannot be expressed rationally and integrally in terms of them. So too for the binary quartic  $p = 4$ , but there prove to be five irreducible covariants and invariants. For the quintic,  $p = 5$ , the facts are even more striking. All covariants and invariants are functions of the five independent ones of lowest degrees. But there prove to be as many as eighteen other covariants and invariants which are irreducible, in that they are not rational integral functions of the five, or of those five and others of as low degrees as themselves among the eighteen.

43]. We have here for clearness adopted a different order of reasoning from that applied in § 30 to invariants alone. There we first found a limit to the number of independent absolute invariants, and deduced conclusions as to the number of independent invariants not necessarily absolute. Here the idea of absolute covariants and invariants is made the subsequent one. In all cases there is one absolute covariant, namely the quantic itself. We have also seen (§§ 41, 29) that there is no other rational integral absolute covariant or invariant. For the *linear* quantic,  $p = 1$ , there is no other independent covariant or invariant whatever, and consequently no other that is absolute. For higher binary quantics,  $p > 1$ , there cannot be more than  $p - 1$  independent absolute covariants and invariants. Otherwise a complete system of  $p$  independent covariants and invariants would be absolute, and consequently all covariants and invariants would be absolute. But for any value of  $p$  exceeding unity there is (§ 15) a non-absolute invariant, the discriminant.

44]. **Covariants of two or more Quantics.** With regard to covariants of two or more quantics in the same variables, the methods of the earlier articles of this chapter yield, in a manner analogous to that of §§ 31 to 33, conclusions of which a summary follows.

Such a covariant is, as in § 35, either homogeneous in the variables or a sum of covariants which are homogeneous in them. Those which are homogeneous in the variables—of the same *order* throughout—form a complete system.

A covariant homogeneous in the variables may or may not be homogeneous in the coefficients of the various quantics severally and collectively. If, however, it be not so homogeneous, it is a sum of covariants every one of which is homogeneous separately in the coefficients of each quantic, and of course therefore in the coefficients of all the quantics collectively. Covariants, then, which are throughout of constant partial *degrees* in the various sets of coefficients, and therefore of constant total *degree* in all the coefficients, form a complete system. This is seen as in § 33.

If  $p_1, p_2, \dots, p_r$  be the orders of  $r$   $q$ -ary ~~quatics~~ *quatics*, the factor by which a covariant of order  $\varpi$  and partial degrees  $i_1, i_2, \dots, i_r$  in their coefficients respectively has to be multiplied to be made equal to the same function of the variables and coefficients in the linearly transformed quantics is

$$M^{\frac{1}{q}(\Sigma \cdot ip - \varpi)},$$

where  $M$  is the modulus of the linear substitution, and

$$\Sigma \cdot ip = i_1 p_1 + i_2 p_2 + \dots + i_r p_r.$$

The whole *weight* of the covariant is constant throughout, and exceeds the index of this power of  $M$  by  $\varpi$ , i. e. is

$$\frac{1}{q} \{ \Sigma \cdot ip + (q-1) \varpi \}.$$

If the covariant is rational and integral this weight must be a positive integer, and consequently the index

$$\frac{1}{q} \{ \Sigma \cdot ip - \varpi \} (= w - \varpi)$$

is not a fraction. It is, moreover, not negative, being the

weight of those coefficients in the covariant which multiply products of the variables in which the last  $\omega$  does not occur, which coefficients cannot all be absent, as no covariant can have  $\omega$  for a factor when the coefficients in the quantics are general.

45.] **Covariants productive of other covariants and invariants.** At this point it may be well to prove an important fact, which, stated for the moment without complete generality, is that any invariant or covariant of a covariant of a quantic is an invariant or covariant, as the case may be, of that quantic itself.

Let  $(a, b, \dots)(x, y, \dots)^p$  be the quantic  $u$ , in any number of variables, in its untransformed shape, and let  $(A, B, \dots)(X, Y, \dots)^p$  be its linearly transformed shape. Also let  $(a', b', \dots)(x, y, \dots)^\omega$  be a covariant of  $u$ , so that  $a', b', \dots$  are functions, of degree  $i$  say, of the coefficients  $a, b, \dots$ , and let  $A', B', \dots$  be the same functions respectively of  $A, B, \dots$ . We have simultaneously the identities

$$(A, B, \dots)(X, Y, \dots)^p = (a, b, \dots)(x, y, \dots)^p,$$

$$(A', B', \dots)(X, Y, \dots)^\omega = M^{\frac{1}{q}(ip-\omega)}(a', b', \dots)(x, y, \dots)^\omega.$$

If then  $K(a', b', \dots)^i(x, y, \dots)^\omega$  be a covariant, or invariant in case  $\omega = 0$ , of the covariant  $(a', b', \dots)(x, y, \dots)^\omega$ , we have

$$\begin{aligned} & K(A', B', \dots)^i(X, Y, \dots)^\omega \\ &= M^{\frac{1}{q}(i'\omega-\omega')} K(M^{\frac{1}{q}(ip-\omega)} a', M^{\frac{1}{q}(ip-\omega)} b', \dots)^i(x, y, \dots)^\omega, \end{aligned}$$

and consequently, in virtue of the homogeneity of the covariant  $K$ ,

$$\begin{aligned} & K(A', B', \dots)^i(X, Y, \dots)^\omega \\ &= M^{\frac{1}{q}(i'\omega-\omega')+\frac{i'}{q}(ip-\omega)} K(a', b', \dots)^i(x, y, \dots)^\omega \\ &= M^{\frac{1}{q}(i'ip-\omega')} K(a', b', \dots)^i(x, y, \dots)^\omega, \end{aligned}$$

which, since  $a', b', \dots$  are functions, of degree  $i$ , of  $a, b, \dots$ , and  $A', B', \dots$  are the same functions respectively of  $A, B, \dots$ , shows that  $K(a', b', \dots)^i(x, y, \dots)^\omega$  is a function, of degree  $i'i$

and order  $\omega'$  of  $a, b, \dots, x, y, \dots$  which, when multiplied by the  $\frac{1}{q}(i' i - \omega')$ th power of the modulus  $M$ , becomes the same function of  $A, B, \dots, X, Y, \dots$ . It is then a covariant of  $u$ , or, in particular if  $\omega' = 0$ , an invariant.

It will be at once seen that only brevity of writing has been secured by attending to but one covariant  $(a', b', \dots)(x, y, \dots)^p$  of but one quantic  $u$ . The argument would have been exactly the same if we had been dealing with more given covariants than one of a quantic, or a given covariant or covariants of more quantics than one in the same variables. We may state in fact the general conclusion, to which the method leads us, as follows.

*Any covariant, or in particular invariant, of any covariant, or system of covariants, of any quantic, or system of quantics in the same variables, is a covariant, or in particular invariant, of that quantic or system of quantics.*

Ex. 12. The binary cubic  $(a, b, c, d)(x, y)^3$  has the covariant (§ 7, Ex. 8), its Hessian,

$$(ac - b^2)x^2 + (ad - bc)xy + (bd - c^2)y^2,$$

which has the invariant (§ 7, Ex. 1),

$$(ad - bc)^2 - 4(ac - b^2)(bd - c^2).$$

This then is an invariant of the cubic. It is its discriminant.

Ex. 13. Find a covariant of degree 3 and order 3, the *cubicovariant*, of the binary cubic.

$$\text{Ans. } (a^2d - 3abc + 2b^3, abd - 2ac^2 + b^2c, -acd + 2b^2d - bc^2, -ad^2 + 3bcd - 2c^3)(x, y)^3,$$

the Jacobian of the cubic and its Hessian.

Ex. 14. Show that the binary quintic  $(a, b, c, d, e, f)(x, y)^5$  has an invariant of the twelfth degree.

Ans. The discriminant of the canonizant. (Cf. § 17, Ex. 20, and Ex. 12 above.)

## CHAPTER IV.

### COGREDIENT AND CONTRAGREDIENT QUANTITIES.

46.] BEFORE proceeding to the further definitions and principles on which most of the propositions of this chapter are to rest, we here first investigate a fruitful method, whose connexion with them will be seen later, for the derivation of invariants and covariants of *binary* quantities, and binary quantities only.

The linear transformation of two variables,

$$\begin{aligned} x &= lX + mY, \\ y &= l'X + m'Y, \end{aligned} \quad \dots (1)$$

leads, as has been seen in § 10, to the equalities of differential operators

$$\left. \begin{aligned} \frac{d}{dX} &= l \frac{d}{dx} + l' \frac{d}{dy}, \\ \frac{d}{dY} &= m \frac{d}{dx} + m' \frac{d}{dy}, \end{aligned} \right\} \quad \dots (2)$$

where on the right the operation is on any function of  $x$  and  $y$ , and on the left it is on the function of  $X$  and  $Y$ , which is equivalent to that function of  $x$  and  $y$  in virtue of (1).

Now the equalities (2) may be written

$$\left. \begin{aligned} (lm' - l'm) \frac{d}{dy} &= l \frac{d}{dY} + m \left( -\frac{d}{dX} \right), \\ (lm' - l'm) \left( -\frac{d}{dx} \right) &= l' \frac{d}{dY} + m' \left( -\frac{d}{dX} \right), \end{aligned} \right\} \quad \dots (3).$$

Thus, except for the factor  $lm' - l'm$ , i.e.  $M$ , the symbols of operation  $\frac{d}{dy}$ ,  $-\frac{d}{dx}$  are transformed by the same scheme of linear substitution as are the variables  $x, y$ .

Thus if  $f(x, y)$  be any homogeneous function, of order  $\varpi$  say, of  $x$  and  $y$ , and if  $F(X, Y)$  be what this becomes when the substitutions (1) are made for  $x$  and  $y$  in it, we have not only

$$F(X, Y) = f(x, y),$$

but also

$$F\left(\frac{d}{dY}, -\frac{d}{dX}\right) = M^{\varpi} f\left(\frac{d}{dy}, -\frac{d}{dx}\right),$$

where the operations on the right and left are on any function of  $x$  and  $y$ , with or without other arguments independent of  $x$  and  $y$ , and on its equivalent in terms of  $X$  and  $Y$  obtained from it by means of (1), respectively.

47.] Let us now apply this fact to covariants of one or more binary quantics. If  $\phi(a, b, \dots, x, y)$  and  $\psi(a, b, \dots, x, y)$  be any two covariants of the quantic or quantics—either or both of them may be in particular the quantic itself, or one of the quantics—of orders  $\varpi, \varpi'$  in  $x$  and  $y$ , and of weights  $w, w'$  respectively, we have, §§ 37, 39,

$$\phi(A, B, \dots, X, Y) = M^{w-\varpi} \phi(a, b, \dots, x, y),$$

and 
$$\psi(A, B, \dots, X, Y) = M^{w'-\varpi'} \psi(a, b, \dots, x, y).$$

We have consequently also, by the preceding article,

$$\phi\left(A, B, \dots, \frac{d}{dY}, -\frac{d}{dX}\right) = M^w \phi\left(a, b, \dots, \frac{d}{dy}, -\frac{d}{dx}\right),$$

and 
$$\psi\left(A, B, \dots, \frac{d}{dY}, -\frac{d}{dX}\right) = M^{w'} \psi\left(a, b, \dots, \frac{d}{dy}, -\frac{d}{dx}\right);$$

whence it follows that by operating with either one of this last pair on either one of the immediately preceding pair, left on left and right on right, we get a covariant identity.

All the four conclusions are really contained in the one

$$\begin{aligned} \phi\left(A, B, \dots, \frac{d}{dY}, -\frac{d}{dX}\right) \psi(A, B, \dots, X, Y) \\ = M^{w+w'-\varpi'} \phi\left(a, b, \dots, \frac{d}{dy}, -\frac{d}{dx}\right) \psi(a, b, \dots, x, y), \end{aligned}$$

for  $\phi$  and  $\psi$  may be interchanged, or may be identical.

Thus the result of operating in this way with any covariant,

one of the quantics, on any covariant, or one of the quantics, is a covariant, or invariant, unless it vanishes. It will certainly vanish if  $\varpi$  the order of  $\phi$  exceeds  $\varpi'$  that of  $\psi$ . It will be an invariant, unless it vanishes, if  $\varpi = \varpi'$ . It will be a covariant, unless it vanishes, if  $\varpi$  is less than  $\varpi'$ .

The exact powers of  $M$  in the above are not essential to the argument. It is of interest, however, to verify that the power in the conclusion is what it should be in accordance with §§ 37, 39. In the operating factor on the right the weight of the coefficient of  $(\frac{d}{dy})^\varpi$  is  $w - \varpi$ , while in the factor operated on that of the coefficient of  $y^{\varpi'}$  is  $w'$ . Also the order is  $\varpi' - \varpi$ . Thus the index of the power of  $M$  should be (§ 39)

$$w - \varpi + w' - (\varpi' - \varpi),$$

i. e.  $w + w' - \varpi'$ , as is the case.

48.] **Invariants of the second degree.** One of the most interesting conclusions from the above is that every binary quantic of even order has an invariant of the second degree. For operate on the binary  $p$ -ic

$$(a_0, a_1, a_2, \dots, a_p)(x, y)^p$$

with the result of putting  $\frac{d}{dy}$ ,  $-\frac{d}{dx}$  for  $x, y$  in itself, i. e. with

$$(a_0, a_1, a_2, \dots, a_p) \left( \frac{d}{dy}, -\frac{d}{dx} \right)^p.$$

We thus get, after division by  $p!$ , that

$$a_0 a_p - p a_1 a_{p-1} + \frac{p(p-1)}{1 \cdot 2} a_2 a_{p-2} - \dots + (-1)^{p-1} p a_{p-1} a_1 + (-1)^p a_p a_0$$

is an invariant unless it vanishes.

It vanishes if  $p$  is odd, as the first and last, second and last but one, &c., terms in that case cancel. If, however,  $p$  is even it does not vanish, but the last term is a repetition of the first, the last but one of the second, and so on till the middle term which stands alone. Thus, halving, and replacing  $p$  by  $2n$ ,

we see that if  $\binom{2n}{r}$  denote the number of combinations of  $2n$  things  $r$  together,

$$a_0 a_{2n} - 2n a_1 a_{2n-1} + \binom{2n}{2} a_2 a_{2n-2} - \binom{2n}{3} a_3 a_{2n-3} + \dots \\ + (-1)^{n-1} \binom{2n}{n-1} a_{n-1} a_{n+1} + (-1)^n \frac{1}{2} \binom{2n}{n} a_n^2$$

is an invariant of the binary  $2n$ -ic

$$(a_0, a_1, a_2, \dots, a_{2n}) (x, y)^{2n}.$$

In particular the binary quadratic, quartic, sextic, &c.

$(a, b, c) (x, y)^2$ ,  $(a, b, c, d, e) (x, y)^4$ ,  $(a, b, c, d, e, f, g) (x, y)^6$ , &c.

have the invariants of the second degree

$$ac - b^2, \\ ae - 4bd + 3c^2, \\ ag - 6bf + 15ce - 10d^2, \\ \&c.,$$

of which the first two have been obtained earlier.

49.] Two different binary quantics of the same order have in all cases, whether their order be even or odd, an invariant of the first degree in the coefficients of each quantic, and so of the second degree on the whole. If the two quantics be

$$(a_0, a_1, a_2, \dots, a_p) (x, y)^p,$$

and

$$(b_0, b_1, b_2, \dots, b_p) (x, y)^p,$$

this joint invariant is in fact

$$(a_0, a_1, a_2, \dots, a_p) \left( \frac{d}{dy}, -\frac{d}{dx} \right)^p (b_0, b_1, b_2, \dots, b_p) (x, y)^p,$$

which, divided by  $p!$ , is seen to be

$$a_0 b_p - p a_1 b_{p-1} + \frac{p(p-1)}{1 \cdot 2} a_2 b_{p-2} - \dots \\ + (-1)^{p-1} p a_{p-1} b_1 + (-1)^p a_p b_0.$$



This is called the *lineo-linear* invariant of the two binary  $p$ -ics.

Of the result Exx. 2, 4 of § 7 are particular cases.

We notice that the results of the preceding article are correctly given from this one by making the  $b$ 's the same as the  $a$ 's, i. e. by making the quantics the same.

We also notice that for an even order  $p$  the joint invariant obtained here for two  $p$ -ics is the intermediate (§ 19) between the invariants of the second degree of the two  $p$ -ics.

These two observations illustrate the fact that we can either pass from invariants of one quantic to those of two of the same kind and order, or from those of two quantics to those of one, but that the information given by two quantics as to one is complete, while that given by one as to two is not so.

Ex. 1. Employ § 47 to find the invariant  $ace + 2bcd - ad^2 - b^2e - c^3$  of a binary quartic by aid of the quartic and its Hessian (§ 11, Ex. 16).

Ex. 2. Find the invariant of degree 4, the discriminant, of a binary cubic by operating with the Hessian on itself, or again by operating on the cubic with its cubicovariant (§ 45, Ex. 13).

Ex. 3. Prove that

$$(a, b, c, d) \left( \frac{d}{dy}, -\frac{d}{dx} \right)^3 \cdot \{(a, b, c, d) (x, y)^3\}^2$$

is  $-108$  times the cubicovariant of the binary cubic.

Ex. 4. The invariants of the second degree

$$ac - b^2, ae - 4bd + 3c^2, ag - 6bf + 15ce - 10d^2, \dots$$

of binary quantics of even order are linear functions of determinants chosen from among

$$\left\| \begin{array}{cccccc} a, & b, & c, & d, & e, & \dots \\ b, & c, & d, & e, & f, & \dots \end{array} \right\|. \quad (\text{Cayley.})$$

Ex. 5. The lineo-linear invariant of the  $x$ - and  $y$ - first differential coefficients of a binary quantic  $u$  of even order is the invariant of the second degree of  $u$ . (Cayley.)

(N.B.—The function obtained in the same way from a binary quantic of odd order is not an invariant.)

Ex. 6. Two binary quantics of different orders  $p, p'$ , ( $p > p'$ ), have a covariant of order  $p - p'$  whose coefficients are lineo-linear.

*Ans.* The result of operating with the second on the first.

Ex. 7. In particular two binary quantics of orders  $p, p - 1$ , have a linear covariant, in the variables, which is also linear in the coefficients of each quantic.

50.] Another result of the close resemblance in form between the schemes (1) and (3) of § 46 is obtained by making (3) operate on any binary quantic  $u$ . We thus get that, when formulae give  $x$  and  $y$  linearly in terms of  $X$  and  $Y$ , the same formulae give  $M \frac{du}{dy}$  and  $-M \frac{du}{dx}$  in terms of  $\frac{du}{dY}$  and  $-\frac{du}{dX}$ . It follows that if in any covariant of a binary quantic  $u$ , homogeneous as usual in the variables,  $\frac{du}{dy}$  and  $-\frac{du}{dx}$  are substituted for  $x$  and  $y$  another covariant of  $u$  is obtained. This theorem is Sylvester's, having been overlooked by Boole who had given the more far-reaching kindred theorem of § 47.

Ex. 8. If in a binary quantic  $u$  we replace  $x$  and  $y$  by

$$\frac{du}{dy} \quad \text{and} \quad -\frac{du}{dx},$$

we obtain the product of  $u$  and a covariant. (*Salmon.*)

*Ans.* That  $u$  is a factor we may see as follows. The values of  $x, y$  which make  $u = 0$  make

$$x \frac{du}{dx} + y \frac{du}{dy} = 0, \text{ i. e. make } \frac{du}{dy} : -\frac{du}{dx} = x : y,$$

so that  $u$ , a homogeneous function  $f(x, y)$ , is a factor of

$$f\left(\frac{du}{dy}, -\frac{du}{dx}\right).$$

Ex. 9. Hence obtain the cubicovariant of a binary cubic.

51.] **Cogredient quantities.** If two equally numerous sets of quantities  $x, y, z, \dots$  and  $x', y', z', \dots$  are such that, whenever one set  $x, y, z, \dots$  are expressed in terms of new quantities  $X, Y, Z, \dots$  by any scheme of linear substitution,

the second set  $x', y', z', \dots$  are expressed in terms of other new quantities  $X', Y', Z', \dots$  by the same scheme of linear substitution, the two sets are said to be sets of *cogredient* quantities.

For instance, the coordinates of two points in a plane, or in space, are cogredient sets of three, or four, quantities.

Again in § 46 it has been shown that, but for the factor  $M$ ,  $\frac{d}{dy}$  and  $-\frac{d}{dx}$  are cogredient with  $x$  and  $y$ .

Once more, if the binary  $p$ -ic

$$(a_0, a_1, a_2, \dots, a_p)(x, y)^p$$

be regarded as a product of  $p$  factors

$$(xy_1 - x_1y)(xy_2 - x_2y) \dots (xy_p - x_py),$$

so that  $\frac{x_1}{y_1}, \frac{x_2}{y_2}, \dots, \frac{x_p}{y_p}$  are the roots, and  $y_1, y_2, \dots, y_p$  may in fact be chosen arbitrarily subject to  $y_1y_2\dots y_p = a_0$ , and if the quantic be linearly transformed by taking

$$x = lX + mY, \quad y = l'X + m'Y,$$

into one of which  $\frac{X_1}{Y_1}, \frac{X_2}{Y_2}, \dots, \frac{X_p}{Y_p}$ , say, are the roots, we have

$$\frac{x_1}{y_1} = \frac{l\frac{X_1}{Y_1} + m}{l'\frac{X_1}{Y_1} + m'} = \frac{lX_1 + mY_1}{l'X_1 + m'Y_1},$$

so that without impropriety we take

$$x_1 = lX_1 + mY_1, \quad y_1 = l'X_1 + m'Y_1,$$

and similarly for other suffixes 2, 3,  $\dots$ ,  $p$ . We have then, in the language of the present article,  $x_1, y_1; x_2, y_2; \dots; x_p, y_p$  cogredient with  $x, y$ .

52.] **Emanants.** Some functions have the covariant property with regard to a quantic or set of quantics, though they involve, not only the coefficients and variables in the quantic or quantics, but also a set or sets of quantities cogredient with those variables. Allowing ourselves some freedom of expression, when no confusion can arise, we may designate

such functions covariants. We proceed to the consideration of a very important class of covariants of this kind.

Let  $u$  be a  $p$ -ic in the  $q$  variables  $x, y, z, \dots$ . The functions

$$\left(x' \frac{d}{dx} + y' \frac{d}{dy} + z' \frac{d}{dz} + \dots\right)^r u,$$

for values of the positive integer  $r$  from 1 to  $p$  inclusive, are defined as the first, second, ...,  $p$ th *emanants* of  $u$ . There would be some convenience in defining the  $r$ th emanant rather as the above expression multiplied by the numerical factor  $\frac{(p-r)!}{p!}$ , but there is no real importance in this, as a numerical multiple of a covariant is of course a covariant not distinct from it, and as we have as yet introduced no convention as to the best numerical multiple of a function, found to have the property of an invariant or covariant of any quantic, to denote by a letter and speak of as that invariant or covariant. Moreover the simplest form is given to general conclusions by use of emanants as written above. Inconvenient numerical factors in any conclusions with regard to quantics of particular orders can be rejected when the end is reached.

The  $p$ th emanant is  $p!$  times the quantic  $u$  itself with  $x, y, z, \dots$  replaced by  $x', y', z', \dots$ . For values of  $r$  exceeding  $p$  there are no emanants, as  $(p+1)$ th differential coefficients of  $u$  vanish.

That the emanants of  $u$  are absolute covariants in the extended sense is readily seen. If we have

$$\begin{aligned} x &= lX + mY + nZ + \dots, \\ y &= l'X + m'Y + n'Z + \dots, \\ z &= l''X + m''Y + n''Z + \dots, \\ &\dots \dots \dots \end{aligned}$$

and

$$\begin{aligned} x' &= lX' + mY' + nZ' + \dots, \\ y' &= l'X' + m'Y' + n'Z' + \dots, \\ z' &= l''X' + m''Y' + n''Z' + \dots, \\ &\dots \dots \dots \end{aligned}$$

then, since

$$\frac{d}{dX} = l \frac{d}{dx} + l' \frac{d}{dy} + l'' \frac{d}{dz} + \dots,$$

$$\frac{d}{dY} = m \frac{d}{dx} + m' \frac{d}{dy} + m'' \frac{d}{dz} + \dots,$$

$$\frac{d}{dZ} = n \frac{d}{dx} + n' \frac{d}{dy} + n'' \frac{d}{dz} + \dots,$$

. . . . .

where on the right the operations are upon any function of  $x, y, z, \dots$ , with or without  $x', y', z', \dots$ , and on the left they are upon the equivalent of that function expressed in terms of  $X, Y, Z, \dots$ , with or without  $X', Y', Z', \dots$ , we have

$$\begin{aligned} X' \frac{d}{dX} + Y' \frac{d}{dY} + Z' \frac{d}{dZ} + \dots &= (lX' + mY' + nZ' + \dots) \frac{d}{dx} \\ + (l'X' + m'Y' + n'Z' + \dots) \frac{d}{dy} &+ (l''X' + m''Y' + n''Z' + \dots) \frac{d}{dz} + \dots \\ &= x' \frac{d}{dx} + y' \frac{d}{dy} + z' \frac{d}{dz} + \dots \end{aligned}$$

Hence by successive operations on  $u$ , any quantic in  $x, y, z, \dots$ , or indeed any function of those variables, the operations on the right and left being upon its original and transformed forms respectively,

$$(X' \frac{d}{dX} + Y' \frac{d}{dY} + Z' \frac{d}{dZ} + \dots) u = (x' \frac{d}{dx} + y' \frac{d}{dy} + z' \frac{d}{dz} + \dots) u,$$

$$(X' \frac{d}{dX} + Y' \frac{d}{dY} + Z' \frac{d}{dZ} + \dots)^2 u = (x' \frac{d}{dx} + y' \frac{d}{dy} + z' \frac{d}{dz} + \dots)^2 u,$$

. . . . .

$$(X' \frac{d}{dX} + Y' \frac{d}{dY} + Z' \frac{d}{dZ} + \dots)^r u = (x' \frac{d}{dx} + y' \frac{d}{dy} + z' \frac{d}{dz} + \dots)^r u,$$

&c., &c.

Thus the emanants are all absolute covariants.

It may be noticed that the emanants may be otherwise expressed. Thus

$$\begin{aligned} \frac{1}{r!} (x' \frac{d}{dx} + y' \frac{d}{dy} + z' \frac{d}{dz} + \dots)^r u \\ = \frac{1}{(p-r)!} (x \frac{d}{dx'} + y \frac{d}{dy'} + z \frac{d}{dz'} + \dots)^{p-r} u', \end{aligned}$$

where  $u'$  is what  $u$  becomes when in it  $x', y', z', \dots$  are put for  $x, y, z, \dots$ . This follows at once from the fact that either side is the coefficient of  $t^r$  in the expansion in powers of  $t$  of  $f(x + tx', y + ty', z + tz', \dots)$ , where  $f(x, y, z, \dots)$  is  $u$ .

Ex. 10. Prove that the emanants are absolute covariants by identifying the results of replacing in  $u$

$$x, y, z, \dots \text{ by } x + tx', y + ty', z + tz', \dots,$$

and  $X, Y, Z, \dots$  by  $X + tX', Y + tY', Z + tZ', \dots$ .

53.] **Geometrical aspect of emanants.** The process of finding emanants is sometimes called the polar process. The student of geometry will notice that the theory of emanants, with regard to ternary and quaternary systems, is that of polar curves and surfaces.

Thus if the ternary  $p$ -ic  $u$  be taken as representing a curve, when equated to zero, its first emanant equated to zero represents the first polar curve of a point  $x', y', z'$  with regard to  $u$ , i.e. a certain curve of order  $p-1$  which possesses the property, among others, of determining by its intersections with  $u$  all the points of contact of tangents from  $x', y', z'$ . The second emanant is in like manner the criterion of the second polar curve of  $x', y', z'$ , i.e. of the first polar curve with regard to the first polar curve; &c., &c.

That the emanants are covariants is the expression of the fact that the various polar curves of a point with regard to a curve are the same, for the same point and the same curve, in whatever system of point-coordinates the curve and point are taken as expressed, and to whatever axes or triangle of reference they are referred.

In like manner, with regard to quaternary quantities, the fact that the emanants are covariants is the fact that the polar surfaces of a point with regard to a surface are the same surfaces whatever be the reference.

54.] **Geometry of binary systems.** The occasion is a good one for a geometrical consideration of binary systems. Their geometry may be regarded either as that of ranges of points on a line or of pencils of lines through a point. To begin with we adopt the former aspect.

Let  $a$  and  $b$  be two fixed points of reference on a straight line,  $P$  any point on that line. Let  $x$  and  $y$  denote  $\lambda aP$  and  $\mu bP$  respectively, where  $\lambda$  and  $\mu$  are constants. Take  $A$  and  $B$  two new fixed points of reference on the same line, and let  $X$  and  $Y$  denote  $\lambda' AP$  and  $\mu' BP$  respectively, where  $\lambda', \mu'$  are new constants. Suppose that  $a$  divides  $AB$  in the ratio  $r : s$  and that  $b$  divides it in the ratio  $\rho : \sigma$ . Then

$$(r+s)aP = sAP + rBP,$$

$$(\rho + \sigma)bP = \sigma AP + \rho BP,$$

so that

$$x = \frac{\lambda}{r+s} \left( \frac{s}{\lambda'} X + \frac{r}{\mu'} Y \right),$$

$$y = \frac{\mu}{\rho + \sigma} \left( \frac{\sigma}{\lambda'} X + \frac{\rho}{\mu'} Y \right).$$

Now these may be identified with

$$x = lX + mY,$$

$$y = l'X + m'Y,$$

by proper choice of  $\frac{r}{s}, \frac{r'}{s'}, \lambda', \mu'$  in terms of  $l, m, l', m'$  and  $\lambda, \mu$ , provided that  $lm' - l'm$  does not vanish.

Thus the most general linear substitution for  $x$  and  $y$  is equivalent to the change of the reference of points  $(x, y)$  to new fixed base points, and the adoption for the new coordinates  $(X, Y)$  of new constant multiples of the distances from those new base points.

A binary  $p$ -ic represents a range of  $p$  points  $P$ .  $(x', y')$  is an additional point  $P'$  on the line of reference. The first, second, &c. emanants are first, second, &c. polar ranges of  $p-1, p-2, \&c.$  points on that line. The property of covariancy belonging to the emanants is the expression of the fact that the polar systems of points are systems of points determined by strictly geometrical connexions with the  $p$  points and the additional point  $P'$ , so that their equations have the same relation to the point  $x', y'$  and the  $p$ -ic whatever

be the base points of reference or the constant multiples. The geometrical connexions are with the  $p+1$  points only, and have no reference, for instance, to the point at infinity on the line, so that they are not metrical connexions.

55.] Or we may adopt a strictly correlative geometrical representation. We may regard a binary  $p$ -ic, equated to zero, as representing  $p$  straight lines through an origin, taking the  $x, y$  of any line through the origin as given constant multiples of the sines of the angles which that line makes with two fixed lines. We may take as new lines of reference any other pair of lines through the origin, and adopt for the  $X, Y$  of the line  $x, y$  any new constant multiples of the sines of the angles which it makes with the new lines of reference. The substitution for  $x, y$  in terms of  $X, Y$  is readily seen to be the most general linear substitution, in virtue of the two degrees of arbitrariness involved in the choice of the new lines of reference, and the two degrees of arbitrariness involved in the choice of the multiples.

A property of covariancy of a function with regard to the  $p$ -ic is expressive of the fact that the pencil of lines obtained by equating the covariant function to zero is a fixed pencil of lines, whatever be the lines of reference or the multiples, the  $p$ -ic equated to zero being a given pencil of lines. In other words, the relation of the pencils of lines is one of strictly geometrical connexion, of a nature entirely uninfluenced by the geometry of other pencils, such as the pencil to the circular points at infinity. If the cogredient  $x', y'$  enter, as is the case with emanants, there is no difference, except that the geometrical connexion of the covariant pencil of lines is with the  $p$ -ic pencil and the line  $(x', y')$ .

It will be noticed that the aspect of the geometry of covariants sketched in § 6 differs from that here developed. There we looked upon a linear substitution as replacing a pencil of lines by a projectively corresponding pencil, retaining the same reference. Here we look upon the substitution as changing the reference, retaining the same pencil. There is a corresponding choice when, as in the last article, we regard the geometry of binary systems as that of ranges of points on a line.



56.] **Covariants derived from emanants.** From the emanants of  $u$ , themselves, as has been seen, covariants in an extended sense, can be derived covariants of  $u$  in the ordinary sense, i.e. covariants free from the quantities  $x', y', z', \dots$  which are cogredient with the variables. The basis of this fact is the following theorem.

*If any of the emanants of  $u$  be expanded and arranged as a quantic in  $x', y', z', \dots$ , any invariant of that quantic is a covariant of  $u$ .*

Considered as a quantic in  $x', y', z', \dots$ , the  $r$ th emanant

$$\left(x' \frac{d}{dx} + y' \frac{d}{dy} + z' \frac{d}{dz} + \dots\right)^r u$$

may be written

$$\left(\frac{d^r u}{dx^r}, \frac{d^r u}{dx^{r-1} dy}, \dots, \frac{d^r u}{dz^r}, \dots\right) (x', y', z', \dots)^r,$$

its coefficients being all the  $r$ th partial derivatives of  $u$ , and so functions of  $x, y, z, \dots$  for values of  $r$  less than  $p$ . Its transformed form is, as has been seen in § 52, similar, so that it may be written

$$\left(\frac{d^r u}{dX^r}, \frac{d^r u}{dX^{r-1} dY}, \dots, \frac{d^r u}{dZ^r}, \dots\right) (X', Y', Z', \dots)^r.$$

Now let  $F(a, b, \dots, k, \dots)$  be an invariant of the quantic  $(a, b, \dots, k, \dots) (x', y', z', \dots)^r$ , so that, for some value of  $\mu$ ,

$$F(A, B, \dots, K, \dots) = M^\mu F(a, b, \dots, k, \dots),$$

where  $(A, B, \dots, K, \dots) (X', Y', Z', \dots)^r$  is the transformed quantic. We conclude that

$$\begin{aligned} F\left(\frac{d^r u}{dX^r}, \frac{d^r u}{dX^{r-1} dY}, \dots, \frac{d^r u}{dZ^r}, \dots\right) \\ = M^\mu F\left(\frac{d^r u}{dx^r}, \frac{d^r u}{dx^{r-1} dy}, \dots, \frac{d^r u}{dz^r}, \dots\right). \end{aligned}$$

But the function  $F$  on the left is the same function of the coefficients and variables in the transformed  $u$  as the function

$F$  on the right is of the coefficients and variables in the untransformed  $u$ ; for each differential coefficient which occurs on the left is the same function of the new variables and coefficients as the corresponding differential coefficient on the right is of the old ones. Thus

$$F\left(\frac{d^r u}{dx^r}, \frac{d^r u}{dx^{r-1} dy}, \dots, \frac{d^r u}{dz^r}, \dots\right)$$

is a function of the coefficients and variables which obeys the definition of covariants.

We see from this theorem that every invariant of a  $q$ -ary  $r$ -ic gives a covariant of any  $q$ -ary quantic of order  $p$  higher than  $r$ , by taking for the  $r$ -ic the  $r$ th emanant of the  $p$ -ic.

Moreover the identity expressive of the fact of covariancy, for any covariant thus derived, involves as its factor  $M^\mu$  exactly the same power of the modulus  $M$  as does the identity which expresses the invariancy of the invariant from which it is derived. In other words, the weight of the coefficients of terms free from the last variable  $\omega$  in the covariant is exactly the weight of the invariant. The degree (in the coefficients) of the covariant is moreover equal to the degree of the invariant.

57.] For an example take the second emanant

$$\left(x' \frac{d}{dx} + y' \frac{d}{dy} + z' \frac{d}{dz} + \dots\right)^2 u.$$

Written as a quantic in  $x', y', z', \dots$  this is

$$\begin{aligned} \frac{d^2 u}{dx^2} x'^2 + 2 \frac{d^2 u}{dx dy} x' y' + 2 \frac{d^2 u}{dx dz} x' z' + \dots \\ + \frac{d^2 u}{dy^2} y'^2 + 2 \frac{d^2 u}{dy dz} y' z' + \dots \\ + \frac{d^2 u}{dz^2} z'^2 + \dots \\ + \dots, \end{aligned}$$

and of this the discriminant, which is (§ 15) an invariant, is

$$\begin{vmatrix} \frac{d^2u}{dx^2} & \frac{d^2u}{dxdy} & \frac{d^2u}{dxdz} & \dots \\ \frac{d^2u}{dxdy} & \frac{d^2u}{dy^2} & \frac{d^2u}{dydz} & \dots \\ \frac{d^2u}{dxdz} & \frac{d^2u}{dydz} & \frac{d^2u}{dz^2} & \dots \\ \dots & \dots & \dots & \dots \end{vmatrix},$$

the Hessian of  $u$ . Another proof that the Hessian is a covariant (§ 11) is thus afforded.

In § 12 we saw that a knowledge that Hessians are covariants told us in particular that discriminants of quadratics are invariants. We now see that the order of reasoning may be reversed. Discriminants are invariants by § 15, and therefore Hessians are covariants.

The geometrical aspect of the fact that the Hessian of a ternary quantic is a covariant may be mentioned. In works on geometry the Hessian of a curve of order  $p$  is found as a curve of order  $3(p-2)$  which has the property of determining the points of inflexion on the first curve by its intersections with it. The covariant property tells us that the curve found by expressing this fact is the same curve whatever be the system of point coordinates or the triangle of reference, i.e. that we do not, when employing different references, obtain different curves with the one property of determining points of inflexion in common to them, but identically the same curve.

Ex. 11. Prove that

$$\frac{d^4u}{dx^4} \frac{d^4u}{dy^4} - 4 \frac{d^4u}{dx^3dy} \frac{d^4u}{dxdy^3} + 3 \left( \frac{d^4u}{dx^2dy^2} \right)^2$$

is a covariant of a binary quantic  $u$  of order exceeding 4.

*Ans.* Factor  $M^4$ . Use § 7, Ex. 5.

Ex. 12. The invariant of the second degree of a binary  $2n$ -ic gives a covariant of any binary quantic of order exceeding  $2n$ .

Ex. 13. Deduce the covariant of a binary quantic found in § 17 from the catalecticant of a quartic (§ 49, Ex. 1).

Ex. 14. Write down from Ex. 11 a quadratic covariant of a binary quintic, and a quartic covariant of a binary sextic.

Ex. 15. Every binary quantic of odd order  $2n + 1$  has a covariant of the second order and second degree.

*Ans.* The invariant of the second degree (§ 48) of its  $2n$ th emanant.

Ex. 16. Every binary quantic of odd order  $2n + 1$  exceeding 3, has at least one linear covariant, obtained by operating on it, as in § 47, with the  $n$ th power of its quadro-quadric covariant (Ex. 15). (*Hermite.*)

*Ans.* For order 3 the result vanishes. For higher orders it does not. To see this let the substitution be adopted which reduces the quadro-quadric covariant to the form  $kXY$ , and the quantic to

$$(A_0, A_1, A_2, \dots, A_{2n+1}) (X, Y)^{2n+1},$$

where  $A_0, A_1, A_2, \dots, A_{2n+1}$  have consequently to satisfy only

$$A_0 A_{2n} - 2n A_1 A_{2n-1} + \binom{2n}{2} A_2 A_{2n-2} - \dots = 0$$

and

$$A_1 A_{2n+1} - 2n A_2 A_{2n} + \binom{2n}{2} A_3 A_{2n-1} - \dots = 0.$$

The linear covariant derived becomes a numerical multiple of

$$A_n X + A_{n+1} Y,$$

and  $A_n = 0, A_{n+1} = 0$  do not follow from the two conditions when  $n > 1$ .

Ex. 17. For the binary quintic this linear covariant is of degree 5. Show that there is another of degree 7, and, assuming as suggested by § 48 and proved hereafter that a binary quantic of odd order has no invariant of degree 2, that it must be distinct from the former.

*Ans.* The Jacobian of the quadro-quadric covariant (Ex. 14) and the linear covariant of Ex. 16.

58]. Precisely as in § 56 we see that if we take more emanants than one of  $u$ , or if we take any emanant or emanants of a covariant of  $u$ , or if we take any emanants of two or more quantics  $u, v, w, \dots$  in the same variables, or of covariants of two or more quantics, taking in all of course the same cogredient quantities  $x', y', z', \dots$ , and if we arrange them as quantics in  $x', y', z', \dots$ , and write down any invariant

of the system of quantics, in  $x', y', z', \dots$  thus obtained, we have a covariant of  $u$ , or of the quantics  $u, v, w, \dots$  jointly.

Ex. 18. From § 9 deduce § 10.

Ex. 19. Employ § 49 to obtain covariants of two binary quantics.

59.] Symbolical representation of covariants and invariants. For full information as to the system of invariants and covariants of a single quantic, it proves to be necessary to have recourse to more quantics than one; and not to quantics in one and the same set of variables only, but to quantics in different cogredient sets of variables.

We here consider binary quantics only.

Let  $x_1, y_1$  and  $x_2, y_2$  be two cogredient pairs of variables, so that simultaneously

$$\begin{aligned} x_1 &= lX_1 + mY_1, & x_2 &= lX_2 + mY_2, \\ y_1 &= l'X_1 + m'Y_1, & y_2 &= l'X_2 + m'Y_2. \end{aligned}$$

We notice that

$$x_1y_2 - x_2y_1 = M(X_1Y_2 - X_2Y_1),$$

where  $M$  is the modulus  $lm' - l'm$ .

Hence, by § 46,  ~~$\frac{d}{dx_1} \frac{d}{dy_2} - \frac{d}{dy_1} \frac{d}{dx_2} = \left[ \frac{d}{dx_1} \frac{d}{dy_2} - \frac{d}{dy_1} \frac{d}{dx_2} \right]$~~

$$\frac{d}{dX_1} \frac{d}{dY_2} - \frac{d}{dY_1} \frac{d}{dX_2} = M \left( \frac{d}{dx_1} \frac{d}{dy_2} - \frac{d}{dy_1} \frac{d}{dx_2} \right),$$

so that  $\frac{d}{dx_1} \frac{d}{dy_2} - \frac{d}{dy_1} \frac{d}{dx_2}$  is what may be called an invariant symbol of operation.

Now let  $u, v$  be any two binary quantics, and let them be called  $u_1, v_1$  when the variables in them are  $x_1, y_1$ , and  $u_2, v_2$  when they are  $x_2, y_2$ . Also let  $U, V; U_1, V_1; U_2, V_2$  denote their linearly transformed forms.

We deduce from the above that, for any positive integral value of  $r$ ,

$$\left( \frac{d}{dX_1} \frac{d}{dY_2} - \frac{d}{dY_1} \frac{d}{dX_2} \right)^r (U_1 V_2) = M^r \left( \frac{d}{dx_1} \frac{d}{dy_2} - \frac{d}{dy_1} \frac{d}{dx_2} \right)^r (u_1 v_2),$$

i. e. 
$$\frac{d^r U_1}{dX_1^r} \cdot \frac{d^r V_2}{dY_2^r} - r \frac{d^r U_1}{dX_1^{r-1} dY_1} \cdot \frac{d^r V_2}{dX_2 dY_2^{r-1}} + \dots$$

$$= M^r \left\{ \frac{d^r u_1}{dx_1^r} \cdot \frac{d^r v_2}{dy_2^r} - r \frac{d^r u_1}{dx_1^{r-1} dy_1} \cdot \frac{d^r v_2}{dx_2 dy_2^{r-1}} + \dots \right\}.$$

In this  $x_1, y_1$  and  $x_2, y_2$  are any cogredient pairs. We may in the expanded result obtained make them the same  $x, y$ . Thus

$$\begin{aligned} \frac{d^r U}{dX^r} \cdot \frac{d^r V}{dY^r} - r \frac{d^r U}{dX^{r-1} dY} \cdot \frac{d^r V}{dX dY^{r-1}} + \dots \\ = M^r \left\{ \frac{d^r u}{dx^r} \cdot \frac{d^r v}{dy^r} - r \frac{d^r u}{dx^{r-1} dy} \cdot \frac{d^r v}{dx dy^{r-1}} + \dots \right\}. \end{aligned}$$

We accordingly have a system of covariants of two binary quantics  $u, v$ . They have already been obtained in § 58, Ex. 19, as the lineo-linear invariants (§ 49) of the  $r$ th emanants of  $u$  and  $v$ . In particular if  $p$  is the order both of  $u$  and  $v$ , the value  $p$  of  $r$  gives the lineo-linear invariant itself, multiplied by  $(p!)^2$ .

Again we may in the expanded result make  $u$  and  $v$  the same quantic, and thus get that

$$\begin{aligned} \frac{d^r u}{dx^r} \cdot \frac{d^r u}{dy^r} - r \frac{d^r u}{dx^{r-1} dy} \cdot \frac{d^r u}{dx dy^{r-1}} \\ + \frac{r(r-1)}{1 \cdot 2} \cdot \frac{d^r u}{dx^{r-2} dy^2} \cdot \frac{d^r u}{dx^2 dy^{r-2}} - \dots \end{aligned}$$

is a covariant, or invariant if  $r = p$  the order of  $u$ , of factor  $M^r$ . For odd values of  $r$  the result is nugatory, in that its first and last, its second and last but one, &c. terms cancel against one another. For even values of  $r$  the last term repeats the first, the last but one repeats the second, and so on till the middle term which occurs once only.

These covariants of  $u$  have already been obtained (§ 57, Ex. 12) as the invariants of the second degree of the emanants of even order of  $u$ . For  $r = p$ , the order of  $u$ , we have in particular a numerical multiple of the invariant of the second degree (§ 48) itself.

So far then the method is only another one for determining results already known in other ways. It as yet gives us only the covariants and invariants of  $u$  which are of the second degree in the coefficients. Its convenience is that it suggests an expressive symbolization for covariants and invariants, and paves the way to a systematic examination of all forms which can be covariants or invariants.

60.] **Hyperdeterminants.** The operator  $\frac{d}{dx_1} \frac{d}{dy_2} - \frac{d}{dy_1} \frac{d}{dx_2}$  is denoted by the brief symbol  $\overline{12}$ . More briefly still the covariant, or invariant, of  $u$  found above by operating  $r$  times with this symbol  $\overline{12}$  on the product  $u_1 u_2$ , and removing all suffixes in the expanded result when the operations have been performed, is called the covariant or invariant  $\overline{12}^r$  of  $u$ .

To get covariants and invariants of the third degree in the coefficients of  $u$  we may consider the product of three quantities  $u_1, v_2, w_3$ , whose suffixes imply that their variables are three cogredient sets, and operate on the product with

$$\left(\frac{d}{dx_1} \frac{d}{dy_2} - \frac{d}{dy_1} \frac{d}{dx_2}\right)^r \left(\frac{d}{dx_2} \frac{d}{dy_3} - \frac{d}{dy_2} \frac{d}{dx_3}\right)^s \left(\frac{d}{dx_3} \frac{d}{dy_1} - \frac{d}{dy_3} \frac{d}{dx_1}\right)^t,$$

thus getting, for positive integral and zero values of  $r, s, t$ , functions seen as in § 59 to have the covariant or invariant property. In the result, after giving it its fully expanded form, we may replace all three sets of cogredient variables  $x_1, y_1; x_2, y_2; x_3, y_3$  by the same set  $x, y$ , and also make  $u, v$  and  $w$  all the same quantic  $u$ . We thus get a system of covariants and invariants of  $u$  which may be symbolically written

$$\overline{12}^r \cdot \overline{23}^s \cdot \overline{31}^t,$$

for different positive integral and zero values of  $r, s$  and  $t$ .

These covariants and invariants are all of the third degree in the coefficients. It is easy to see the necessary connexions of  $r, s, t$  and  $p$ , the order of  $u$ , that they may be invariants. Any term in the covariant or invariant is a product of differential coefficients of the three factors  $u u u$ , the first being differentiated as many times as the figure 1 occurs in the symbolic product, the second as many times as the figure 2 occurs, and the third as many times as 3 occurs. Now if the expansion be an invariant each one must be differentiated  $p$  times, where  $p$  is the order of  $u$ . For an invariant then the conditions are

$$r + s = s + t = t + u = p.$$

To get covariants and invariants of the fourth degree we have in like manner to operate on products of four quantities.

The symbolical form of such covariants and invariants is

$$\overline{12}r_{12} \overline{13}r_{13} \overline{14}r_{14} \overline{23}r_{23} \overline{24}r_{24} \overline{34}r_{34},$$

and the conditions for an invariant are

$$r_{12} + r_{13} + r_{14} = r_{12} + r_{23} + r_{24} = r_{13} + r_{23} + r_{34} = r_{14} + r_{24} + r_{34} = p.$$

In like manner covariants and invariants of the fifth, sixth, &c. degrees have symbolical expressions involving five, six, &c. letters, derived from products of five, six, &c. quantics, eventually made the same. For invariants every figure must occur in the same number of symbolical factors, and that number must be the order of the quantic of which they are invariants.

The method is one by which Cayley made great advances in the systematic exhibition of covariant and invariant forms. (See, for instance, his collected works, Vol. I, pp. 95-112.) To develop it is outside the limits of the present introductory work. For the examination of what symbolical expressions do not vanish, and the reduction of irreducible systems, the student is referred to Cayley's original memoirs, and to Salmon's *Higher Algebra*, Lesson XIV, &c. The method, which is spoken of as that of 'hyperdeterminants,' did not, in its originator's form, succeed in establishing the finiteness of complete systems of irreducible covariants in general. That triumph was reserved for another symbolical method, having much in common with it, which will be referred to in the following article.

There is a corresponding theory for ternary and higher quantics, which will not be entered into. The student will have no difficulty in seeing that, acting upon a product of three ternary quantics,

$$\left| \begin{array}{ccc} \frac{d}{dx_1}, & \frac{d}{dy_1}, & \frac{d}{dz_1} \\ \frac{d}{dx_2}, & \frac{d}{dy_2}, & \frac{d}{dz_2} \\ \frac{d}{dx_3}, & \frac{d}{dy_3}, & \frac{d}{dz_3} \end{array} \right|,$$

which may be called  $\overline{123}$ , is an invariant operator.

61.] **Transvectants.** The invariant or covariant  $\overline{12}^r (uv)$ ,



or rather this multiplied by  $\frac{(p-r)! (p'-r)!}{p! p'!}$ , is called the  $r$ th *transvectant* of  $u$  and  $v$ , in German the  $r$ th 'Ueberschiebung von  $u$  über  $v$ .' The process of forming transvectants of  $u$  and  $v$  is called *transvection*.

In particular the covariant or invariant  $\overline{12}^r$  of  $u$ , i.e. the covariant or invariant  $\overline{12}^r (uu)$ , is  $\left\{ \frac{p!}{(p-r)!} \right\}^2$  times the  $r$ th transvectant of  $u$  and itself.

From two binary quantics  $u, v$ , whose orders are  $p, p'$ , of which  $p \neq p'$ , are derived  $p+1$  transvectants. For, besides the values  $1, 2, 3, \dots, p$  of  $r$  the value  $0$  is also admissible. The  $0$ th transvectant of  $u$  and  $v$  is the product  $uv$ . The other transvectants of  $u$  and  $v$  are the covariants, or covariants and invariant, which are of the first degree in the coefficients of  $u$  and also of the first degree in the coefficients of  $v$ , and so altogether of the second degree. As has been seen, one is the product  $uv$ , and the rest may be found from the first, second, ...,  $p$ th emanants of  $u$  and of  $v$  by writing down the linear invariants of corresponding pairs of those emanants as in § 49.

The  $0$ th transvectant of  $u$  and itself is  $u^2$ . The other transvectants of  $u$  and itself are the covariants of  $u$  of degree 2 in the coefficients, obtained by writing down the invariants of the second degree of the successive emanants of  $u$ . The first, third, fifth, &c. transvectants of  $u$  and itself vanish.

In the theory of transvectants symbolic products  $\overline{12}^r \cdot \overline{23}^s \dots$  involving powers of more than one symbolic factor have no place.

The Cayleyan notation  $\overline{12}^r (uv)$  for transvectants is the most concise and easily grasped for the presentation of a particular covariant, but another symbolical notation, which will not here be dealt with, is more frequently used, and is best for the purposes of research into the theory of complete covariant systems. It is symbolic *ab initio*, denoting a binary quantic  $(a_0, a_1, a_2, \dots, a_p) (x, y)^p$  by  $(ax + a'y)^p$ , where  $a^r a'^{p-r}$  means  $a_r$ , and  $a^r a'^s$  has no meaning unless  $r+s$  is a multiple of  $p$ , and denoting also a covariant  $(\theta_0, \theta_1, \theta_2, \dots, \theta_w) (x, y)^w$  by  $(\theta x + \theta' y)^w$  in like manner. It is identified with the names of Aronhold, Clebsch and Gordan.

The last named succeeded, in fact, by means of it, in establishing the finiteness of the complete system of irreducible covariants and invariants for any binary quantic or quantics.

With regard to a single binary quantic  $u$ , what researches with the improved notation of transvectants have succeeded in establishing is, that all concomitants (this term including both covariants and invariants) of  $u$  which are of the second degree in the coefficients are transvectants of  $u, u^2$  in particular being the 0th transvectant; that all of the third degree in the coefficients are linear functions of transvectants of  $u$  and concomitants of the second degree, products of  $u$  into  $u^2$  and other concomitants of the second degree being included as 0th transvectants; that all of the fourth degree are transvectants of  $u$  and concomitants of the third degree; and so on from degree to degree. Gordan has proved that this continued process ceases after a time to give new irreducible concomitants, so that the determination of a complete system of irreducible concomitants for any binary quantic is reduced to the examination for irreducibility of those which are obtained as transvectants up to a certain point. His proof is a somewhat difficult mathematical induction based upon showing that if the system is finite for the  $p$ -ic it is for the  $(p + 1)$ -ic.

And the finiteness of the complete system of irreducible concomitants of a number of binary quantics has also been established. In the case of two quantics  $u, v$  the complete system is comprised in the complete system of  $u$ , the complete system of  $v$ , and a terminating system of transvectants of the one complete system with the other complete system. In the case of three quantics  $u, v, w$ , the complete system of concomitants is comprised in the complete system of  $u, v$ , the complete system of  $w$ , and a terminating system of mutual transvectants of these complete systems; and so on for any number of quantics.

This grand theory should be studied in Clebsch's *Theorie der Binären Algebraischen Formen*, or in Gordan's *Vorlesungen über Invariantentheorie*. A brief treatment of it is also given in Salmon's *Higher Algebra*, Lesson XX. It lies beyond the scope of the present treatise.

Ex. 20. The Jacobian of  $u$  and  $v$  is their first mutual transvectant, a numerical factor discarded.

Ex. 21. The Hessian of a binary quantic  $u$  is the second transvectant of  $u$  and itself, but for a numerical factor.

Ex. 22. The quadratic invariant (§ 48) of a binary  $2n$ -ic is half its  $2n$ th transvectant with itself.

Ex. 23. The lineo-linear invariant (§ 49) of two binary  $p$ -ics is their  $p$ th mutual transvectant.

Ex. 24. The second transvectant of the cubic  $(a, b, c, d) (x, y)^3$  and itself is twice the quadratic covariant

$$(ac - b^2)x^2 + (ad - bc)xy + (bd - c^2)y^2.$$

Also the first and third transvectants vanish.

Ex. 25. The first transvectant of the cubic and its second transvectant (Ex. 24) is the cubicovariant (§ 45, Ex. 13)

$$(a^2d - 3abc + 2b^3)x^3 + \dots$$

The second mutual transvectant vanishes.

Ex. 26. The first transvectant of the cubic and its cubicovariant is minus twice the square of the quadratic covariant  $(ac - b^2)x^2 + \dots$ . Their second mutual transvectant vanishes. Their third is minus twice the discriminant

$$a^2d^2 - 6abcd + 4ac^3 + 4b^3d - 3b^2c^2 \equiv (ad - bc)^2 - 4(ac - b^2)(bd - c^2).$$

Ex. 27. The second transvectant of the quartic  $(a, b, c, d, e) (x, y)^4$  and itself is twice the quartic covariant (the Hessian simplified by omitting a numerical factor)

$$(ac - b^2)x^4 + 2(ad - bc)x^3y + (ae + 2bd - 3c^2)x^2y^2 + 2(be - cd)xy^3 + (ce - d^2)y^4;$$

and the fourth transvectant is twice the invariant

$$I \equiv ae - 4bd + 3c^2.$$

Ex. 28. The first transvectant of the quartic and the quartic covariant of Ex. 27 is half a sextic covariant beginning

$$(a^2d - 3abc + 2b^3)x^6 + \dots:$$

the second is one sixth of the product of  $I$  and the quartic; the third vanishes; and the fourth is three times the invariant

$$J \equiv ace + 2bcd - ad^2 - b^3e - c^3.$$

Ex. 29. The binary quintic  $(a, b, c, d, e, f) (x, y)^5$  has a covariant of the second order and the second degree.

$$\text{Ans. } (ae - 4bd + 3c^2)x^2 + (af - 3be + 2cd)xy + (bf - 4ce + 3d^2)y^2,$$

half the fourth transvectant of the quintic and itself. Cf. § 57, Ex. 16.

Ex. 30. The binary quintic has an invariant of the fourth degree.

*Ans.*  $(af - 3be + 2cd)^2 - 4(ae - 4bd + 3c^2)(bf - 4ce + 3d^2)$ ,  
found as minus twice the second transvectant of the covariant of Ex. 29 and itself.

Ex. 31. The binary quintic has a linear covariant of degree 5 in the coefficients.

*Ans.* The fourth transvectant of the quintic and the square of the covariant of Ex. 29. Cf. § 57, Ex. 17.

Ex. 32. Find a covariant of the fourth order and second degree of the binary sextic.

*Ans.*  $\frac{\overline{123}^4}{12}$ .

Ex. 33. A ternary quadratic has an invariant of the third degree whose symbol is  $\overline{123}^2$ ; and a ternary quartic one whose symbol is  $\overline{123}^4$ .

Ex. 34. A  $q$ -ary  $2n$ -ic has an invariant of degree  $q$ .

62.] **Contragredient quantities.** Two sets of quantities  $x, y, z, \dots; \xi, \eta, \zeta, \dots$  are said to be *contragredient* when formulæ of linear substitution for the first set

$$\begin{aligned} x &= lX + mY + nZ + \dots, \\ y &= l'X + m'Y + n'Z + \dots, \\ z &= l''X + m''Y + n''Z + \dots, \\ &\dots \dots \dots \end{aligned}$$

are necessarily accompanied by the associated but different formulæ of substitution for the second set

$$\begin{aligned} \Xi &= l\xi + l'\eta + l''\zeta + \dots, \\ H &= m\xi + m'\eta + m''\zeta + \dots, \\ Z &= n\xi + n'\eta + n''\zeta + \dots, \\ &\dots \dots \dots \end{aligned}$$

in which latter set it is to be noticed that the new quantities are expressed in terms of the old, and not vice versa. Reversed they are

$$\begin{aligned} \xi &= M^{-1}\{\lambda\Xi + \mu H + \nu Z + \dots\}, \\ \eta &= M^{-1}\{\lambda'\Xi + \mu'H + \nu'Z + \dots\}, \\ \zeta &= M^{-1}\{\lambda''\Xi + \mu''H + \nu''Z + \dots\}, \\ &\dots \dots \dots \end{aligned}$$

where  $M$  is as usual the modulus of the scheme of substitution

for  $x, y, z, \dots$ , and where  $\lambda, \mu, \dots, \lambda', \mu', \dots, \dots$  denote the minors

$$\frac{dM}{dl}, \frac{dM}{dm}, \dots, \frac{dM}{dl'}, \frac{dM}{dm'}, \dots, \dots$$

It is convenient to speak of contragredient quantities as being linearly transformed by schemes of substitution of which one is the *dual* of the other. The name is reasonable, as the duality or reciprocal connexion of the two substitutions is precise, it being possible, and a good simple exercise in determinant algebra, to show that the first substitution stands in precisely the same relation to the second as the second does to the first.

63.] **Geometrical contragrediency.** The duality of transformations of contragredient quantities has its counterpart and its application in the method of duality in geometry. Taking ternary systems, we know in fact that if  $x, y, z$  and  $\xi, \eta, \zeta$  be point- and line-coordinates of associated systems, so that  $\xi x + \eta y + \zeta z = 0$  is the condition that the point  $(x, y, z)$  lie on the line  $(\xi, \eta, \zeta)$ , or that the line  $(\xi, \eta, \zeta)$  pass through the point  $(x, y, z)$ , as, for instance, is the case when the coordinates of a point are areal and the coordinates of a line the perpendiculars upon it from the vertices of the triangle of reference, then the first scheme of linear substitution of § 62 applied to  $x, y, z, \dots$  reduces this condition to the form

$$\xi(lX + mY + nZ) + \eta(l'X + m'Y + n'Z) + \zeta(l''X + m''Y + n''Z) = 0,$$

$$\text{i. e. } (l\xi + l'\eta + l''\zeta)X + (m\xi + m'\eta + m''\zeta)Y + (n\xi + n'\eta + n''\zeta)Z = 0,$$

which is of the same form,

$$\Xi X + H Y + Z Z = 0,$$

as before if the formulae of substitution for  $\xi, \eta, \zeta$  be those of the second scheme in § 62.

Thus corresponding systems of point- and line-coordinates are transformed to corresponding systems by dual linear substitutions. In other words they are contragredient quantities.

64.] Another remark of great importance is that the

symbols  $\frac{d}{dx}, \frac{d}{dy}, \frac{d}{dz}, \dots$  are contragredient with the variables  $x, y, z, \dots$ . For when

$$\begin{aligned} x &= lX + mY + nZ + \dots, \\ y &= l'X + m'Y + n'Z + \dots, \\ z &= l''X + m''Y + n''Z + \dots, \\ &\dots \dots \dots \end{aligned}$$

we have, as used frequently already,

$$\begin{aligned} \frac{d}{dX} &= l \frac{d}{dx} + l' \frac{d}{dy} + l'' \frac{d}{dz} + \dots, \\ \frac{d}{dY} &= m \frac{d}{dx} + m' \frac{d}{dy} + m'' \frac{d}{dz} + \dots, \\ \frac{d}{dZ} &= n \frac{d}{dx} + n' \frac{d}{dy} + n'' \frac{d}{dz} + \dots, \\ &\dots \dots \dots \end{aligned}$$

and these accord with the two dual schemes of § 62.

65.] From the formulae of § 62 it follows at once that

$$\begin{aligned} \Xi X + \text{H} Y + \text{Z} Z + \dots &= (lX + mY + nZ + \dots) \xi \\ &+ (l'X + m'Y + n'Z + \dots) \eta + (l''X + m''Y + n''Z + \dots) \zeta + \dots \\ &= \xi x + \eta y + \zeta z + \dots; \end{aligned}$$

so that  $\xi x + \eta y + \zeta z + \dots$  obeys the absolute invariant law.

We might, in fact, have defined the contragrediency of  $x, y, z, \dots$  and  $\xi, \eta, \zeta, \dots$  by postulating that their corresponding schemes of linear substitution are such as to leave

$$\xi x + \eta y + \zeta z + \dots$$

unaltered. This has been illustrated by means of the geometrical contragrediency of § 63.

In the case of the contragrediency of § 64 this persistence in form of  $\xi x + \eta y + \zeta z + \dots$  means simply that

$$X \frac{d}{dX} + Y \frac{d}{dY} + Z \frac{d}{dZ} + \dots = x \frac{d}{dx} + y \frac{d}{dy} + z \frac{d}{dz} + \dots,$$

which, when we remember Euler's theorem of homogeneous functions, we see to be only the expression of the fact that a homogeneous function of any order in  $x, y, z, \dots$  becomes

upon linear transformation a homogeneous function of the same order in the new variables.

66.] **Contravariants and mixed concomitants.** If  $u$ , a quantic in  $x, y, z, \dots$ , be expressed in terms of new variables  $X, Y, Z, \dots$  by linear transformation, and if  $\xi, \eta, \zeta, \dots$  be quantities contragredient to  $x, y, z, \dots$ , and accordingly expressed in terms of new quantities  $\Xi, H, Z, \dots$  by the substitution dual to that giving  $x, y, z, \dots$  in terms of  $X, Y, Z, \dots$ , there are found to exist functions of  $\xi, \eta, \zeta, \dots$  and of the coefficients in  $u$ , which need at most to be multiplied by factors involving only the constants of the transformation, always, in fact, powers of the modulus, to be made equal to the same functions of  $\Xi, H, Z, \dots$  and of the coefficients in the transformed form of  $u$ . Such functions are called *contravariants* of  $u$ .

There also exist functions possessing the same property, which involve both  $x, y, z, \dots$  and  $\xi, \eta, \zeta, \dots$ , as well as the coefficients in  $u$ . Such functions are called *mixed concomitants* of  $u$ .

There also exist contravariants and mixed concomitants of systems of two or more quantics in the same variables  $x, y, z, \dots$ .

Invariants, covariants, contravariants, and mixed concomitants are all spoken of as *concomitants* of the quantic or quantics to which they belong.

In a certain sense  $\xi x + \eta y + \zeta z + \dots$  may be itself spoken of as a mixed concomitant. It has, however, no particular reference to any quantic or quantics, but is a function of persistent form of the two contragredient sets of quantities only. It is the *universal* mixed concomitant of all quantics in  $x, y, z, \dots$  or in  $\xi, \eta, \zeta, \dots$ .

In a better sense, contravariants and mixed concomitants of a quantic  $u$  in  $x, y, z, \dots$  are regarded as respectively invariants and covariants of the system consisting of  $u$  and the linear form  $\xi x + \eta y + \zeta z + \dots$ .

Ex. 35. If  $\phi(x, y, z, \dots)$  be a covariant and  $\psi(\xi, \eta, \zeta, \dots)$  a contravariant of  $u$ , then

$$\psi\left(\frac{d}{dx}, \frac{d}{dy}, \frac{d}{dz}, \dots\right)\phi(x, y, z, \dots)$$

is a covariant or invariant, and

$$\phi\left(\frac{d}{d\xi}, \frac{d}{d\eta}, \frac{d}{d\zeta}, \dots\right)\psi(\xi, \eta, \zeta, \dots)$$

is a contravariant or invariant. (*Sylvester.*)

67.] **Evectants.** It is from the last-mentioned point of view that contravariants and mixed concomitants of a quantic or quantics are most easily discovered.

The method of § 19 may, in fact, be applied to determine a series of contravariants from any invariant of a quantic  $u$ , or a series of mixed concomitants from any covariant. We have only in any invariant or covariant  $P$  to put for every coefficient in  $u$  the corresponding coefficient in

$$u + k(\xi x + \eta y + \zeta z + \dots)^p,$$

where  $p$  is the order of  $u$ , and to take separately the coefficients of  $k, k^2, \dots$  in the expanded result. These separately are (§ 19) invariants or covariants, as the case may be, of  $u$  and  $(\xi x + \eta y + \zeta z + \dots)^p$ , and consequently of  $u$  and

$$\xi x + \eta y + \zeta z + \dots$$

In other words, they are, as the case may be, contravariants or mixed concomitants of  $u$ .

The method has, it will be remembered, been already used for binary quantics in § 20, Ex. 33.

If the quantic  $u$  be

$$ax^p + pbx^{p-1}y + pb'x^{p-1}z + \dots \\ + \frac{p(p-1)}{1.2}cx^{p-2}y^2 + \frac{p(p-1)}{1.2}c'x^{p-2}z^2 + \dots,$$

where the numerical factors of the various coefficients are the corresponding coefficients in the expansion of the multinomial  $(x + y + z + \dots)^p$ , the  $r$ th of these contravariants or mixed concomitants is

$$\left(\xi^p \frac{d}{da} + \xi^{p-1}\eta \frac{d}{db} + \xi^{p-1}\zeta \frac{d}{db'} + \dots \right. \\ \left. + \xi^{p-2}\eta^2 \frac{d}{dc} + \xi^{p-2}\zeta^2 \frac{d}{dc'} + \dots\right)^r P.$$

The contravariants obtained from any invariant  $P$  of  $u$  in this way are called the first, second, ...  $r$ th, ... *evectants* of  $P$ .



The same method applies for the determination of contravariants and mixed concomitants of two or more quantities in the same variables. The same operator as before

$$\xi^p \frac{d}{da} + \xi^{p-1} \eta \frac{d}{db} + \xi^{p-1} \zeta \frac{d}{db'} + \dots \\ + \xi^{p-2} \eta^2 \frac{d}{dc} + \xi^{p-2} \zeta^2 \frac{d}{dc'} + \dots,$$

in which  $p$  and  $a, b, b', \dots, c, c', \dots$  refer to one of the quantities only, suffices to derive series of contravariants and mixed concomitants from invariants and covariants of the system.

The general theory of contravariants is Sylvester's. Evectants are due to Hermite.

Ex. 36. Show that

$$e\xi^4 - 4d\xi^3\eta + 6c\xi^2\eta^2 - 4b\xi\eta^3 + a\eta^4,$$

and

$$(ce - d^2)\xi^4 - 2(be - cd)\xi^3\eta + (ae + 2bd - 3c^2)\xi^2\eta^2 \\ - 2(ad - bc)\xi\eta^3 + (ac - b^2)\eta^4,$$

are contravariants of the binary quartic  $(a, b, c, d, e)$   $(x, y)^4$ .

*Ans.* The first evectants of  $I$  and  $J$ .

Ex. 37. Find a cubic contravariant of a binary cubic as an evectant of its discriminant.

Ex. 38. Use the method of evectants to show that the left-hand side of the tangential equation

$$(bc - f^2)\xi^2 + \dots + 2(gh - af)\eta\xi + \dots = 0$$

of the conic

$$u \equiv ax^2 + by^2 + cz^2 + 2fyz + 2gzx + 2hxy = 0$$

is a contravariant of  $u$ .

Ex. 39. From the invariant

$$\Theta \equiv a'(bc - f^2) + \dots + 2f'(gh - af) + \dots$$

of two ternary quadratics obtain a contravariant of

$$ax^2 + \dots + 2fyz + \dots, a'x^2 + \dots + 2f'yz + \dots$$

in which both sets of coefficients occur.

68.] **Contravariants of binary quantities not distinct from covariants.** In the case of binary systems there is a connexion between contragrediency and cogrediency which has nothing analogous to it in the cases of ternary and higher systems.

Let  $x, y$  and  $\xi, \eta$  be pairs of contragredient variables, so that with the formulae of linear substitution

$$x = lX + mY, \quad y = l'X + m'Y \quad \dots (1)$$

go as companions the formulae

$$\Xi = l\xi + l'\eta, \quad \text{H} = m\xi + m'\eta. \quad \dots (2)$$

These latter may be written

$$M\eta = l\text{H} - m\Xi, \quad -M\xi = l'\text{H} - m'\Xi, \quad \dots (3)$$

where  $M = lm' - l'm$ ; and these differ only by the presence of the factor  $M$  on the left of each from the results of putting  $\eta, -\xi, \text{H}, -\Xi$  for  $x, y, X, Y$  in (1). This has already been encountered in the particular case of § 46.

In an extended sense, then, we may say that  $\eta$  and  $-\xi$  are *cogredient* with  $x$  and  $y$ . The factor  $M$  will not affect the legitimacy of their use as cogredient variables with  $x$  and  $y$ , so long as only homogeneous functions are dealt with, provided, of course, that we pay proper attention to the alteration of the power of  $M$  which occurs as a factor on a side of any equality we are dealing with.

We may equally say that, when  $x'$  and  $y'$  are cogredient with  $x$  and  $y$ , then  $-y'$  and  $x'$  are contragredient with  $x$  and  $y$ , but for a factor which is immaterial so long as homogeneous functions are dealt with. In particular we may say that  $-y$  and  $x$  are, with this reservation, contragredient with  $x$  and  $y$ .

If, then, in any contravariant of a binary quantic  $u$ , or of several binary quantics, we replace  $\xi$  and  $\eta$  by  $-y$  and  $x$ , we obtain a function of  $x, y$  and the coefficients of  $u$ , or of  $u$  and the other quantics, which persists in form but for a factor involving only the constants of transformation, a power of  $M$ , after any linear transformation. In other words, we obtain a covariant of  $u$ , or of  $u$  and the other quantics.

In accordance with what has been said, however, it is clear that the power of  $M$ , which occurs as a factor in the relation expressive of the covariancy of the derived covariant, is different from that which occurs in the relation expressive of the contravariancy of the contravariant from which it is obtained.

Ex. 40. Apply this conclusion to § 67, Ex. 36.

Ex. 41. Obtain the cubicovariant (§ 45, Ex. 13) of a binary cubic by means of the first evectant of the discriminant. (Cf. § 67, Ex. 37.)

69.] **Covariants of  $u$  are invariants of  $u$  and a linear form.** It is a proposition closely associated with the remark of the preceding article that all invariants, of a complete system (§ 33), of a binary quantic or quantics  $u, v, w, \dots$  and the linear form  $xy' - x'y$  are, when in them  $x', y'$  are replaced by  $x, y$ , covariants of the quantic or quantics  $u, v, w, \dots$ ; and that conversely all covariants of  $u, v, w, \dots$  are, when in them  $x, y$  are replaced by  $x', y'$ , invariants of the system consisting of  $u, v, w, \dots$  and the linear form  $xy' - x'y$ .

This is easy to see; for, if  $x', y'$  are cogredient with  $x, y$ ,

$$xy' - x'y = M(XY' - X'Y).$$

Now a complete system of invariants of  $u, v, w, \dots$  and  $xy' - x'y$  involve  $x', y'$  homogeneously (§ 33), so that to insert  $MX, MY$  for  $x', y'$  in an invariant is the same thing as to insert  $X, Y$  for them and multiply by a power of  $M$ . If, then, in a supposed invariant of  $u, v, w, \dots$  and  $xy' - x'y$ , we put  $x, y$  for  $x', y'$ , we get a function of the coefficients in  $u, v, w, \dots$  and of  $x, y$ , which persists in form, but for a power of  $M$  as factor, after linear transformation. In other words, we get a covariant of  $u, v, w, \dots$ . And conversely, if in any covariant of  $u, v, w, \dots$  we put  $x', y'$  for  $x, y$ , we get a function of the sets of coefficients in  $u, v, w, \dots$  and  $xy' - x'y$  which again persists in form, but for a power of  $M$  as factor, after linear transformation. In other words, we get an invariant of  $u, v, w, \dots$  and  $xy' - x'y$ .

A fact closely related to this, and, indeed, a particular case of the first part of the theorem, is that all invariants of the binary  $(p+1)$ -ic

$$(xy' - x'y)u$$

are, when  $x', y'$  are replaced by  $x, y$ , covariants of the binary  $p$ -ic  $u$ .

## CHAPTER V.

### BINARY QUANTICS. INVARIANTS, ETC., AS FUNCTIONS OF DIFFERENCES.

70.] IN most of the chapters which follow binary quantics will alone be considered, except where otherwise stated. Special methods may be with advantage adopted for the discovery and examination of their concomitants. Moreover, it will be seen later that from invariants and covariants of binary quantics there is a means of passing to those of a ternary quantic, that, in fact, invariants and covariants of a ternary quantic are a class of invariants and covariants of a system of binary quantics. From ternary quantics there is a like passage to quaternary; and so on. Thus there is more than simplicity of treatment in favour of a close examination of binary quantics alone in the first place.

71.] **Convention as to numerical multiples of concomitants.** If  $I$  is an invariant, so is  $\mu I$ , where  $\mu$  is any numerical constant. If  $K$  is a covariant, so is  $\mu K$ . The invariants  $I$  and  $\mu I$  are not of course regarded as distinct invariants, nor the covariants  $K, \mu K$  distinct covariants. It will be well now to adopt some convention which will relieve us from any ambiguity as to numerical multipliers when we speak of any invariant or covariant  $I$  or  $K$ . The following is probably the best convention as to invariants of a single binary quantic

$$(a_0, a_1, a_2, \dots a_p)(x, y)^p.$$

Take the term or terms in the invariant which involve  $a_0$  to the highest power. If there are more such terms than one, suppose that  $a_r$  is the next earliest coefficient which occurs in any of them. Choose among them the term or terms which involve  $a_r$  to the highest power. If there are more than one of these terms, let  $a_s$  be the next earliest coefficient which

occurs in any of them, and take that term or those terms among them which involve  $a$ , to the highest power; and so on continually till we get but a single term. Now divide or multiply the invariant by such a numerical quantity as will give this term the coefficient +1. The invariant thus prepared is what we nearly always henceforth mean when we speak of the invariant as a precise function.

And as to covariants the convention is similar. Take the coefficient of the term free from  $y$  in a covariant which has been found. Among the terms of which this coefficient consists single out one by the same rule as above, and apply to the covariant the numerical factor which will reduce the numerical coefficient of this term to +1. By the covariant we henceforth mean the covariant thus numerically prepared.

The rule may be more briefly stated if we call the coefficients in the quantic  $a, b, c, d, e, \dots$  instead of  $a_0, a_1, a_2, a_3, a_4, \dots$ . Suppose the factors of every term in the invariant, or in the coefficient of  $x^m$  in the covariant, written from left to right in alphabetical order. Among all the terms choose the one which comes first in alphabetical order, i. e. the one which would stand first in a dictionary. Make the coefficient of this term +1.

Thus, in invariants and covariants of various quantic which we have already met with,

$$\begin{aligned} ac - b^2, \\ (ad - bc)^2 - 4(ac - b^2)(bd - c^2), \\ (a^2d - 3abc + 2b^3)x^3 + \dots, \\ ae - 4bd + 3c^2, \\ ace + 2bcd - ad^2 - b^2e - c^3, \end{aligned}$$

the coefficient +1 is given to the terms  $ac, a^2d^2, a^2dx^3, ae, ace$ , respectively, by the above rule.

72.] **Covariancy of the factors of a binary quantic.** A method already touched upon in § 51 will be now more fully considered.

Let  $\frac{x_1}{y_1}, \frac{x_2}{y_2}, \dots, \frac{x_p}{y_p}$  be the  $p$  roots of the general binary  $p$ -ic

$$(a_0, a_1, a_2, \dots, a_p)(x, y)^p,$$

i. e. let them be the roots of the equation in  $x:y$  obtained by equating the  $p$ -ic to zero. Moreover let the denominators  $y_1, y_2, \dots, y_p$ , which are of course arbitrary, be so chosen that

$$y_1 y_2 \dots y_p = a_0.$$

Another expression for the  $p$ -ic must then be

$$(xy_1 - x_1y)(xy_2 - x_2y) \dots (xy_p - x_py).$$

As explained in § 51, we may with propriety say that the pairs  $x_1, y_1; x_2, y_2; \dots; x_p, y_p$  are cogredient with  $x$  and  $y$ ; that every suffixed  $x$  and the corresponding suffixed  $y$  are particular corresponding values of  $x$  and  $y$ . Thus, going with the formulae of linear substitution

$$x = lX + mY, \quad y = l'X + m'Y,$$

we have, for every value of  $r$  from 1 to  $p$  inclusive,

$$x_r = lX_r + mY_r, \quad y_r = l'X_r + m'Y_r,$$

so that

$$xy_r - x_r y = M(XY_r - X_r Y), \quad \dots (1)$$

and similarly,  $r$  and  $s$  being two distinct numbers not exceeding  $p$ ,

$$x_r y_s - x_s y_r = M(X_r Y_s - X_s Y_r). \quad \dots (2)$$

By means of (1) we have that

$$\begin{aligned} (xy_1 - x_1y)(xy_2 - x_2y) \dots (xy_p - x_py) \\ = M^p (XY_1 - X_1Y)(XY_2 - X_2Y) \dots (XY_p - X_pY) \end{aligned}$$

is an equivalent way of writing the identity

$$(a_0, a_1, a_2, \dots, a_p)(x, y)^p = (A_0, A_1, A_2, \dots, A_p)(X, Y)^p;$$

and we are consequently told that,  $a_r$ , any coefficient in the original form of our  $p$ -ic, being a function of  $x_1, x_2, \dots, x_p$  and  $y_1, y_2, \dots, y_p$ , the corresponding coefficient  $A_r$  in the transformed form of the  $p$ -ic is  $M^p$  times that same function of  $X_1, X_2, \dots, X_p$  and  $Y_1, Y_2, \dots, Y_p$ .

In particular we have

$$M^p Y_1 Y_2 \dots Y_p = A_0.$$

73.] Now take  $H_w$  any homogeneous function of degree  $w$  in the differences such as  $x_r y_s - x_s y_r$ , and let  $H'_w$  be the same

function of the corresponding differences  $X_r Y_s - X_s Y_r$ . By the equalities (2) above we see that

$$H_w = M^w H'_w.$$

Should it then be possible to express  $H_w$  as a homogeneous function, of degree  $i$  say, of  $a_0, a_1, a_2, \dots a_p$ , in which case  $H'_w$  would be the same function of  $M^{-p}$  times the corresponding new coefficients  $A_0, A_1, A_2, \dots A_p$ , and so, in virtue of the homogeneity, would be  $M^{-ip}$  times the same function of  $A_0, A_1, A_2, \dots A_p$ , that function will be an invariant of the  $p$ -ic.

Such expression is not always possible. We shall see, however, that there is no invariant which cannot be given in this manner.

74.] Invariants functions of differences of roots. *Every invariant of*

$$(a_0, a_1, a_2, \dots a_p)(x, y)^p$$

*can be expressed as a function of the differences between pairs of roots, symmetric in the roots, multiplied by a power of  $a_0$ , with or without a purely numerical factor.*

It may of course be expressed as a product of  $a_0^i$ , where  $i$  is its degree, into a symmetric function of the roots; for it is (§ 22) a homogeneous function of  $a_0, a_1, a_2, \dots a_p$ , and  $\frac{a_1}{a_0}, \frac{a_2}{a_0}, \dots \frac{a_p}{a_0}$  are symmetric functions of the roots.

To see that only differences between pairs of roots need occur in the expression for it, it will suffice to prove that it is unaltered when all the roots are increased or diminished by any the same quantity.

This is made clear by expressing that the invariant is one for the particular substitution

$$x = X + mY, y = Y,$$

of which the modulus is unity, so that by it the invariant is absolutely unaltered. Now this substitution is that of putting  $\frac{x}{y} = \frac{X}{Y} + m$ , i.e. is that by which the equation given by equating the  $p$ -ic to zero is altered into one whose roots are all less by  $m$  than its own roots.

may be  
proved  
582

An invariant of degree  $i$  is then the product of  $a_0^i$  and a function of differences between pairs of roots which is symmetric in the roots. By 'invariant' we of course here mean 'rational integral invariant.' An irrational invariant need not be symmetric in the roots. For instance, if  $a, \beta, \gamma, \delta$  be the roots of a quartic, the anharmonic ratio  $\frac{(a-\gamma)(\beta-\delta)}{(a-\delta)(\beta-\gamma)}$ , when expressed in terms of the coefficients, is an irrational invariant. Throughout the following our reasoning is with regard to rational integral invariants.

The converse of the proposition of this article is not true. Every invariant is a product of a power of  $a_0$  and a function of differences between roots, but it is not every function of differences between roots, symmetric in the roots, which, when expressed in terms of the coefficients and made integral in them by a power of  $a_0$  as factor, gives an invariant.

In the following article functions of the differences between roots, symmetric in the roots, are examined in the light of §§ 72, 73 to ascertain what classes of them are and what are not productive of invariants.

Ex. 1. Show that every invariant of a binary  $p$ -ic must vanish for the special  $p$ -ic  $(x+y)^p$ ; and hence that the sum of the numerical coefficients of the terms in an invariant must vanish.

75.] What functions of differences produce invariants? We have to consider such functions as

$$a_0^i \times \text{a sum of products of differences like } \frac{x_r}{y_r} - \frac{x_s}{y_s},$$

the sum involving all the roots  $\frac{x_1}{y_1}, \frac{x_2}{y_2}, \dots, \frac{x_p}{y_p}$  symmetrically.

Note that the idea of this symmetry is not quite identical with that of symmetry in differences like  $\frac{x_r}{y_r} - \frac{x_s}{y_s}$ . We do best to avoid the expression 'symmetric function of the differences.'

For the function, when expressed in terms of the coefficients, to be integral,  $i$  must be not less than the index of the highest power to which any root occurs in the symmetric function; for,  $\frac{a_1}{a_0}, \frac{a_2}{a_0}, \dots, \frac{a_p}{a_0}$  being all of the first degree in any par-



icular root, a symmetric function of the roots whose degree in any particular one is  $i'$  cannot be equal to a function of degree less than  $i'$  in  $\frac{a_1}{a_0}, \frac{a_2}{a_0}, \dots, \frac{a_p}{a_0}$ , so that to make it integral the factor  $a_0^{i'}$  at least is necessary.

Referring to § 72 we may now write the function in the form  $(y_1 y_2 \dots y_p)^i \Sigma \left\{ \text{product of factors like } \frac{x_r y_s - x_s y_r}{y_r y_s} \right\}$ ,

which may also be written

$$\Sigma \{ y_1^{\lambda_1} y_2^{\lambda_2} \dots y_p^{\lambda_p} \times \text{product of factors like } x_r y_s - x_s y_r \},$$

where  $\lambda_1, \lambda_2, \dots, \lambda_p$  are in general not all zero; but such as are not zero are positive.

The indices  $\lambda_1, \lambda_2, \dots, \lambda_p$  will, however, be all zero if in every product of differences in the summation every root occurs in the same number  $i$  of factors. In such a case the product takes the form

$$\Sigma \{ \text{product of } \frac{1}{2} ip \text{ factors like } x_r y_s - x_s y_r \},$$

and is consequently such a function as the  $H_w$  of § 73, for the value  $\frac{1}{2} ip$  of  $w$ . Under such circumstances the chosen function is an invariant by that article.

And this is the only class of cases in which the chosen function is an invariant. In other cases the integers which we have called  $\lambda_1, \lambda_2, \dots, \lambda_p$  are none of them negative and do not all vanish. Now if to

$$\Sigma \{ y_1^{\lambda_1} y_2^{\lambda_2} \dots y_p^{\lambda_p} \times \text{product of factors like } x_r y_s - x_s y_r \}$$

we apply the substitution  $x = -Y, y = X$ , of which the modulus is unity, we obtain, remembering the cogrediency with  $x$  and  $y$  of every suffixed  $x$  and  $y$ ,

$$\Sigma \{ X_1^{\lambda_1} X_2^{\lambda_2} \dots X_p^{\lambda_p} \times \text{same product of factors like } X_r Y_s - X_s Y_r \},$$

and this cannot be the same function of the coefficients in

$$(X Y_1 - X_1 Y) (X Y_2 - X_2 Y) \dots (X Y_p - X_p Y)$$

as the form above in small letters is of the coefficients in

$$(x y_1 - x_1 y) (x y_2 - x_2 y) \dots (x y_p - x_p y)$$

unless the  $\lambda$ 's all vanish; as otherwise we should have an identity

$$\begin{aligned} \Sigma \{y_1^{\lambda_1} y_2^{\lambda_2} \dots y_p^{\lambda_p} \times \text{product of factors like } x_r y_s - x_s y_r\} \\ = \Sigma \{x_1^{\lambda_1} x_2^{\lambda_2} \dots x_p^{\lambda_p} \times \text{same product of factors}\}. \end{aligned}$$

Now this is an impossibility; for, since  $\lambda_1 + \lambda_2 + \dots + \lambda_p$  is positive, the dimensions of the left in the  $y$ 's collectively exceed those of the right, and the dimensions of the right in the  $x$ 's collectively exceed those of the left.

Putting then together the results of this and the preceding article, we have the following complete theorem.

*Any homogeneous function of weight  $w$  of the differences between pairs of roots of a binary quantic, which is symmetric in the roots, and such that in all products of differences of which it consists every root is involved in the same number  $\frac{2w}{p}$ , or  $i$ , of factors, is, when made integral in the coefficients by the factor  $a_0^i$ , an invariant of that quantic: other functions of the differences, though symmetric in the roots, do not however produce invariants: nor is there any invariant which cannot be thus expressed.*

We may, however, contemplate the possibility of invariants which are sums of numerical multiples of different invariants obtained, as above, from different simple symmetric sums of differences for which  $w$  and  $i$  are the same. In other words, there may be different invariants of the same weight and degree. This does not conflict with the above general statement.

By 'invariant' in the theorem we mean 'rational integral invariant.' Irrational invariants are given by functions of differences having all the same properties except that of symmetry in the roots.

Ex. 2. If  $a, \beta, \gamma, \delta$  be the roots of a binary quartic

$$a_0^2 \Sigma \{(a-\beta)^2 (\gamma-\delta)^2\}, \quad \text{i. e.} \quad \Sigma \{(x_1 y_2 - x_2 y_1)^2 (x_3 y_4 - x_4 y_3)^2\}$$

is an invariant.

On the other hand  $\Sigma \{(a-\beta)^4 (\gamma-\delta)^2\},$

$$\text{i. e.} \quad \Sigma \{y_1^{-4} y_2^{-4} y_3^{-2} y_4^{-2} (x_1 y_2 - x_2 y_1)^4 (x_3 y_4 - x_4 y_3)^2\}$$

is not productive of an invariant.

76.] The identity expressive of the invariancy of an invariant written down as above is deduced from the result

$$H_w = M^w H'_w$$

of § 73 as follows.

$H_w$  is of  $w$  dimensions in  $y_1, y_2, \dots, y_p$ . In case it produces an invariant, what we have is that

$$\begin{aligned} H_w &= (y_1 y_2 \dots y_p)^{\frac{2w}{p}} \times \text{function of weight } w \text{ of differences of} \\ &\quad \text{roots, symmetric in the roots,} \\ &= a_0^{\frac{2w}{p}} F\left(\frac{a_1}{a_0}, \frac{a_2}{a_0}, \dots, \frac{a_p}{a_0}\right) \\ &= I_w^{\frac{2w}{p}}(a_0, a_1, a_2, \dots, a_p), \text{ say,} \end{aligned}$$

where  $w$  denotes the weight, and  $\frac{2w}{p}$  the degree in

$$a_0, a_1, a_2, \dots, a_p.$$

Similarly

$$\begin{aligned} H'_w &= \left(\frac{A_0}{M^p}\right)^{\frac{2w}{p}} F\left(\frac{A_1}{A_0}, \frac{A_2}{A_0}, \dots, \frac{A_p}{A_0}\right) \\ &= M^{-2w} I_w^{\frac{2w}{p}}(A_0, A_1, A_2, \dots, A_p). \end{aligned}$$

Thus the equality  $H_w = M^w H'_w$  becomes

$$I_w^{\frac{2w}{p}}(A_0, A_1, A_2, \dots, A_p) = M^w I_w^{\frac{2w}{p}}(a_0, a_1, a_2, \dots, a_p),$$

which is in accordance with the result obtained earlier (§§ 23, 26) that the index of the power of  $M$  in the identity expressive of invariancy is the weight of the invariant, or, writing  $i$  for  $\frac{2w}{p}$  the degree of the invariant, is  $\frac{1}{2} ip$ .

77.] **Discriminants.** The product of the squares of the differences between roots of a binary  $p$ -ic is a single product, and involves all roots in equal numbers of its factors, viz. every root in  $2(p-1)$  factors. It belongs then to the class of symmetric functions which according to § 75 produce invariants. Now  $a_0^{2(p-1)}$  times this product is the discriminant, or rather (cf. § 71) is a numerical multiple of the discriminant. We have then a direct proof that the discriminant of any binary quantic is an invariant, as has been otherwise seen earlier (§ 15) for all quantics.

The weight of the discriminant is  $p(p-1)$ , and its degree is  $2(p-1)$ . Thus, in accordance with the general theory,

$$w = \frac{1}{2} ip.$$

78.] Invariants of quadratics and of cubics. Binary *quadratics* and *cubics* have no invariants but their discriminants and powers of those discriminants.

For the *quadratic* this is obvious. For there is only one difference  $a \sim \beta$  between two roots  $a, \beta$ , and no function, of a single weight, of this difference can be symmetric in  $a$  and  $\beta$  unless it be an even power of  $a - \beta$ .

For the *cubic* § 30 tells us that there cannot be two independent invariants. The discriminant, then, being one, it follows that there is no other which is not a function of that discriminant, and consequently, as invariants are of one weight throughout, none that is not a mere power of the discriminant.

We may also reason as follows. Let  $a, \beta, \gamma$  be the roots of the cubic. Any invariant must, by § 75, but for a possible numerical factor, be of the form

$$\alpha_0^i \Sigma \{(\beta - \gamma)^r (\gamma - a)^s (a - \beta)^t\},$$

and must be such that every root occurs in the same number of factors of  $(\beta - \gamma)^r (\gamma - a)^s (a - \beta)^t$ . Consequently

$$s + t = t + r = r + s; \cdot$$

whence

$$r = s = t.$$

Moreover,  $r + s + t$  must be constant, and equal to the weight  $\frac{1}{2} ip = \frac{3}{2} i$ , for all products under  $\Sigma$ . There can then be only one product repeated or not. The invariant is in fact necessarily

$$\alpha_0^i \{(\beta - \gamma)(\gamma - a)(a - \beta)\}^{\frac{3}{2}i}.$$

Again, the index  $\frac{1}{2} i$  of the power of the product must be not only integral, but even. For an odd power of

$$(\beta - \gamma)(\gamma - a)(a - \beta)$$

is not symmetric in the roots, being altered in sign when two roots,  $\beta$  and  $\gamma$  say, are interchanged. Thus every invariant of the cubic is a power of

$$\alpha_0^i \{(\beta - \gamma)(\gamma - a)(a - \beta)\}^2$$

or a numerical multiple of such a power; and this function is a numerical multiple of the discriminant.

79.] **Discriminants freed from inconvenient numerical factors.** The right numerical multiples of  $a_0^2(a-\beta)^2$  and  $a_0^4\{(\beta-\gamma)(\gamma-a)(a-\beta)\}^2$  to speak of as the discriminants of the binary quadratic and cubic, are decided by the convention of § 71.

By elementary processes of the theory of equations it can be proved, taking for convenience  $(a, b, c)(x, y)^2$  and  $(a, b, c, d)(x, y)^3$  to be the quadratic and cubic, that for the two cases respectively

$$a^2(a-\beta)^2 = -4(ac-b^2),$$

$$\text{and } a^4\{(\beta-\gamma)(\gamma-a)(a-\beta)\}^2 \\ = -27\{(ad-bc)^2 - 4(ac-b^2)(bd-c^2)\};$$

and it is to the expressions in brackets on the right that, in accordance with § 71, the name of discriminants is properly given, and not to  $-4$  and  $-27$  times those expressions respectively.

In Cayley's fourth memoir on quantics the corresponding multiple to the  $-4$  and  $-27$  above has been found in the case of the discriminant of any binary quantic. Consider the binary  $p$ -ic  $(a_0, a_1, a_2, \dots, a_p)(x, y)^p$ . The product of the squares of differences between pairs of its roots is of weight  $p(p-1)$ , being the product of  $\frac{1}{2}p(p-1)$  factors of two dimensions in the roots, and consequently, being an invariant, is of degree  $\frac{2w}{p} = 2(p-1)$ . That this is the degree also follows from the fact that any particular root enters in  $p-1$  squared factors, and so to the degree  $2(p-1)$ . For the degree in the coefficients  $a_1, a_2, \dots, a_p$  is the degree in any particular root. We have, then, to consider the expression for

$$a_0^{2(p-1)} \Pi (a_r - a_s)^2$$

in terms of the coefficients.

Since  $p(p-1)$  is the weight, the term  $a_0^{p-1} a_p^{p-1}$ , if it actually occurs, is the one term in the discriminant into which  $a_0$  enters to the highest power. This, then, if it occurs, is the term in the discriminant to which the coef-

ficient + 1 is given in accordance with the convention of § 71. Now the term must occur; for it is the only one which does not vanish for the special  $p$ -ic  $a_0 x^p + a_p y^p$ , and the discriminant of this special  $p$ -ic does not vanish, since no two roots of the equation  $a_0 z^p + a_p = 0$  are equal when neither  $a_0$  nor  $a_p$  vanishes.

Now consider the yet more special  $p$ -ic  $x^p + y^p$ . Its roots are those of  $z^p + 1 = 0$ , i.e. the  $p$   $p$ th roots of  $-1$ . Denote these by  $\rho_1, \rho_2, \dots, \rho_p$ . We know that if  $\rho_1, \rho_2, \dots, \rho_p$  are the roots of  $f(z) = 0$ , then

$$f'(z) = \frac{f(z)}{z - \rho_1} + \frac{f(z)}{z - \rho_2} + \dots + \frac{f(z)}{z - \rho_p},$$

so that  $f'(\rho_1) = \left[ \frac{f(z)}{z - \rho_1} \right]_{z = \rho_1} = (\rho_1 - \rho_2)(\rho_1 - \rho_3) \dots (\rho_1 - \rho_p)$ ,

and so for other roots  $\rho_2, \rho_3, \dots, \rho_p$ . It follows, by multiplication of the  $p$  right-hand and  $p$  left-hand members, that

$$f'(\rho_1) f'(\rho_2) \dots f'(\rho_p) = (-1)^{\frac{1}{2} p(p-1)} \Pi (\rho_r - \rho_s)^2,$$

the sign being as stated because each of the  $\frac{1}{2} p(p-1)$  differences  $\rho_r \sim \rho_s$  occurs once in the product as  $\rho_r - \rho_s$  and once as  $\rho_s - \rho_r$ . In the present case, then, of the equation  $z^p + 1 = 0$ ,

$$p \rho_1^{p-1} \cdot p \rho_2^{p-1} \cdot \dots \cdot p \rho_p^{p-1} = (-1)^{\frac{1}{2} p(p-1)} \Pi (\rho_r - \rho_s)^2,$$

so that  $\Pi (\rho_r - \rho_s)^2 = (-1)^{-\frac{1}{2} p(p-1)} p^p (\rho_1 \rho_2, \dots, \rho_p)^{p-1}$   
 $= (-1)^{-\frac{1}{2} p(p-1)} p^p \{(-1)^p\}^{p-1}$   
 $= (-1)^{\frac{1}{2} p(p-1)} p^p.$

Now if, for the general  $p$ -ic  $(a_0, a_1, a_2, \dots, a_p)(x, y)^p$ ,

$$a_0^{2(p-1)} \Pi (a_r - a_s)^2 = k \{a_0^{p-1} a_p^{p-1} + \dots\},$$

we get as a particular case of this

$$\Pi (\rho_r - \rho_s)^2 = k.$$

Consequently  $k = (-1)^{\frac{1}{2} p(p-1)} p^p$ .

Thus the product  $a_0^{2(p-1)} \Pi (a_r - a_s)^2$  for the binary  $p$ -ic is properly spoken of, not as the discriminant, but as

$$(-1)^{\frac{1}{2} p(p-1)} p^p$$

times the discriminant.

The multipliers  $-4$ ,  $-27$  for the cases of the quadratic and cubic accord with this general result.

80.] **The binary quartic.** The binary quartic  $(a, b, c, d, e)(x, y)^4$  has not more than two independent invariants (§ 30). Now the discriminant is one (§§ 15, 77). There are, however, two of lower degrees than this. It is preferable to regard them as the two fundamental invariants, and the discriminant as consequently a function of them. The two are the  $I$  and  $J$  of § 29.

In fact, 
$$a^2 \Sigma \{(\beta - \gamma)^2 (a - \delta)^2\}$$
 and 
$$a^3 \Sigma \{(\beta - \gamma) (a - \delta) (\gamma - a)^2 (\beta - \delta)^2\}$$
 are invariants according to the criterion of § 75.

A remark as to the number of terms covered in these and such like summations will be here in place. There is generally clearness gained by the wider of two possible interpretations. In fact we do best to bear always in mind that for purposes of expression in terms of the coefficients it is symmetry in the roots rather than in the differences which is fundamental. Thus, since we may take a first root  $a$  in four ways, then a second  $\beta$  in three ways, and then a third  $\gamma$  in two ways, we regard each of the above sums as a sum of twenty-four terms, even though these are, in the first, three terms eight times repeated, and, in the second, six terms four times repeated. The student is recommended to give the close attention necessary to convince him of this second fact.

Let the four roots be separated into a triad  $a, \beta, \gamma$  and the fourth  $\delta$ , and let  $(\beta - \gamma) (a - \delta)$ ,  $(\gamma - a) (\beta - \delta)$ ,  $(a - \beta) (\gamma - \delta)$  be denoted by  $u, v, w$ . Then he will see that the first sum is eight times

$$a^2 (u^2 + v^2 + w^2), \quad \dots (1)$$

and the second four times

$$a^3 \{u (v^2 - w^2) + v (w^2 - u^2) + w (u^2 - v^2)\}, \quad \dots (2)$$

which latter may, by elementary algebra, equally be written

$$-a^3 \{u^2 (v - w) + v^2 (w - u) + w^2 (u - v)\},$$

or 
$$a^3 (v - w) (w - u) (u - v),$$

or again 
$$\frac{1}{3} a^3 \{(v - w)^3 + (w - u)^3 + (u - v)^3\}.$$

The values of the invariants (1) and (2) in terms of the

coefficients may be calculated by the ordinary methods of symmetric functions. (See, for instance, Burnside and Panton's *Theory of Equations*, § 27, Exx. 16, 18.) The results are that

$$a^2(u^2 + v^2 + w^2) = 24(ae - 4bd + 3c^2) = 24I$$

$$\begin{aligned} \text{and} \quad a^3 \{u(v^2 - w^2) + v(w^2 - u^2) + w(u^2 - v^2)\} \\ = -432(ace + 2bcd - ad^2 - b^2e - c^3) = -432J. \end{aligned}$$

This direct process is however unnecessary. For it will be seen in the next chapter that  $I$  and  $J$  are the only invariants of degrees 2 and 3 which the quartic possesses. The invariants (1) and (2) must then be numerical multiples  $kI, k'J$  of  $I$  and  $J$  respectively. This being granted, that  $k$  and  $k'$  have the values 24 and  $-432$  respectively may be seen by considering a particular case. Take, for instance, the particular case of the quartic whose roots are  $\pm 1, \pm 2$ .

It will be noticed that in  $I$  and  $J$  the alphabetically leading terms  $ae, ace$  have the coefficient  $+1$  according to the convention of § 71.

The proof that all rational integral invariants of the quartic can be rationally and integrally expressed in terms of  $I$  and  $J$  is reserved till a later chapter.

81.] **Discriminant of quartic.** The discriminant, an invariant, must, as explained at the opening of the last article, be a function of  $I$  and  $J$ . We proceed to see what function.

It is an invariant of degree 6 and weight 12 (§ 77), which vanishes when the quartic has a square factor, and consequently when it has the square factor  $y^2$ , i.e. when  $a = 0$  and  $b = 0$ .

Now in this case  $I$  and  $J$  become  $3c^2$  and  $-c^3$  respectively. Of these no rational integral function vanishes except

$$(3c^2)^3 - 27(-c^3)^2$$

and powers of  $(3c^2)^3 - 27(-c^3)^2$ .

Consequently

$$I^3 - 27J^2 \equiv (ae - 4bd + 3c^2)^3 - 27(ace + 2bcd - ad^2 - b^2e - c^3)^2,$$

whose degree and weight are right, is the discriminant of the quartic.

The coefficient of its alphabetically leading term  $a^3e^3$  is correctly  $+1$ .



By the general proposition of § 79, the invariant

$$a^6(a-\beta)^2(a-\gamma)^2(a-\delta)^2(\beta-\gamma)^2(\beta-\delta)^2(\gamma-\delta)^2,$$

or, in the notation of the preceding article,  $a^6 u^2 v^2 w^2$ , which must be a numerical multiple of the discriminant, is equal to

$$4^4 (I^3 - 27J^2) = 256 (I^3 - 27J^2).$$

Ex. 3. Prove this also by showing that  $v-w$ ,  $w-u$ ,  $u-v$  are the roots of the cubic  $z^3 - 36a^{-2}Iz + 432a^{-3}J = 0$ , and that  $27^2 u^2 v^2 w^2$  is the product of the squares of differences between roots of this cubic.

Ex. 4. Obtain the same result by showing that  $432^2 a^{-6} J^2$  is the product of the squares of differences between roots of the cubic

$$t^3 - 12a^{-2}It - uvw = 0,$$

whose roots are  $u$ ,  $v$ ,  $w$ .

Ex. 5. The products  $au$ ,  $av$ ,  $aw$  are irrational invariants of the binary quartic.

*Ans.* Cf. § 75, or Ex. 4 above.

Ex. 6. The six ratios of  $u$ ,  $v$ ,  $w$  to one another, which are respectively minus the six anharmonic ratios of the factors of the quartic, are irrational fractional invariants.

Ex. 7. Any anharmonic ratio  $\frac{(a_1 - a_3)(a_2 - a_4)}{(a_2 - a_3)(a_1 - a_4)}$  of any four factors of a binary  $p$ -ic is an irrational invariant of the  $p$ -ic.

Ex. 8. All invariants of a binary quartic can be expressed as functions of the discriminant and any single anharmonic ratio

$$\frac{(a-\gamma)(\beta-\delta)}{(a-\delta)(\beta-\gamma)}$$

of the factors.

Ex. 9. All invariants of a binary  $p$ -ic are functions of the discriminant and the  $p-3$  anharmonic ratios of four factors

$$\frac{(\beta-a)(\gamma-a_4)}{(\gamma-a)(\beta-a_4)}, \quad \frac{(\beta-a)(\gamma-a_5)}{(\gamma-a)(\beta-a_5)}, \quad \dots \quad \frac{(\beta-a)(\gamma-a_p)}{(\gamma-a)(\beta-a_p)},$$

where  $a$ ,  $\beta$ ,  $\gamma$  are three roots, and  $a_4, a_5, \dots, a_p$  the rest. (*Cayley.*)

*Ans.* These are  $p-2$  independent invariants.

Ex. 10. In a binary quartic for which  $I = 0$  the six anharmonic ratios of the four factors are equal in sets of three to the two imaginary cube roots of  $-1$ . Geometrically the pencil or range of

four elements which the quartic denotes is said to be 'equi-anharmonic.'

*Ans.* From  $u^2 + v^2 + w^2 = 0$  and  $u + v + w = 0$

we have 
$$\left(\frac{u}{v}\right)^2 + \frac{u}{v} + 1 = 0. \quad \text{Hence, \&c.}$$

Ex. 11. In a binary quartic for which  $J = 0$  the six anharmonic ratios are  $-1, -1, 2, 2, \frac{1}{2}, \frac{1}{2}$ ; and the pencil or range is harmonic.

82.] **Covariants as functions of differences.** We now proceed to notice briefly the facts as to covariants of a binary quantic which are analogous to, and in reality include, the facts as to invariants dealt with in §§ 72 to 76.

Using the results (1) and (2) of § 72 we can at once write down the analogue of the first statement in § 73; viz. that, if  $G_w$  be a homogeneous function of degree  $w$  in the two sets of differences whose types are  $xy_r - x_r y$  and  $x_r y_s - x_s y_r$ , where

$\frac{x_1}{y_1}, \frac{x_2}{y_2}, \dots, \frac{x_p}{y_p}$  are the roots of

$$(a_0, a_1, a_2, \dots, a_p)(x, y)^p,$$

and if  $G'_w$  be the same homogeneous function of the corresponding differences  $XY_r - X_r Y$ , &c., and  $X_r Y_s - X_s Y_r$ , &c. with reference to the roots of the transformed quantic

$$(A_0, A_1, A_2, \dots, A_p)(X, Y)^p,$$

then

$$G_w = M^w G'_w.$$

Should it then be possible to express  $G_w$  in terms of the variables  $x, y$  and the coefficients  $a_0, a_1, a_2, \dots, a_p$ , and  $G'_w$  as the same function, divided by a power of the modulus, of  $X, Y$  and  $A_0, A_1, A_2, \dots, A_p$ , such a function  $G_w$  when so expressed will be a covariant.

Notice that any covariant so obtained must be *homogeneous in the coefficients*  $a_0, a_1, a_2, \dots, a_p$ . For if the expression for  $G_w$  be

$$K(a_0, a_1, a_2, \dots, a_p; x, y),$$

that for  $G'_w$  is

$$K\left(\frac{A_0}{M^p}, \frac{A_1}{M^p}, \frac{A_2}{M^p}, \dots, \frac{A_p}{M^p}; X, Y\right);$$

and this must be homogeneous in the fractions with denomi-

nator  $M^p$  if the equality  $G_w = M^w G'_w$  take the form

$$K(A_c, A_1, A_2, \dots, A_p; X, Y) = M^w K(a_0, a_1, a_2, \dots, a_p; x, y).$$

It is to be concluded also that  $G_w$ , or  $K$ , is *homogeneous in  $x$  and  $y$* , i. e. that in every product of differences which is a part of  $G_w$  there must occur the same number of differences  $xy_r - x_r y$ ,  $xy_s - x_s y$ , ... of the first type. For, the coefficients  $a_0, a_1, a_2, \dots, a_p$  being all homogeneous in  $x_1, x_2, \dots, x_p, y_1, y_2, \dots, y_p$ , the covariant  $K$ , or  $G_w$ , must be homogeneous in these quantities. Now any term in  $G_w$  which is a product of  $\omega$  differences of the type  $xy_r - x_r y$  and, consequently,  $w - \omega$  differences of the other type  $x_r y_s - x_s y_r$ , is of dimensions  $\omega + 2(w - \omega)$ , i. e.  $2w - \omega$ , in  $x_1, x_2, \dots, x_p, y_1, y_2, \dots, y_p$ . This then having to be constant for all products of which  $G_w$  consists,  $\omega$  must be the same for all.

Covariants thus produced are then necessarily of one degree in the coefficients and one order in the variables throughout.

83.] Now any covariant of degree  $i$  and order  $\omega$  of a binary

quantic is necessarily  $a_0^i y^\omega$  times a function of the differences between roots and of the differences between  $\frac{x}{y}$  and roots.

For it must be unaltered in form by the linear substitution

$$x = X + mY, y = Y$$

of which the modulus is 1; that is to say, it must be unchanged when  $\frac{x}{y}$  and all the roots are diminished by any the same quantity  $m$ .

The particular functions of the two sets of differences which, when prepared by such factors  $a_0^i y^\omega$ , are covariants, are exactly those which when so prepared become functions such as  $G_w$  above.

Such functions of the differences must, of course, for the covariants to be rational and integral, be symmetric in the roots, as otherwise they could not be expressed rationally in terms of the coefficients and  $x, y$  at all. They must also be

such sums of products of differences that in every product  $\frac{x}{y}$  occurs in a constant number  $\omega$  of factors, and  $\frac{x_1}{y_1}, \frac{x_2}{y_2}, \dots, \frac{x_p}{y_p}$

each in a constant number  $i$  of factors, the same number  $i$  for all. The prepared symmetric function is, in fact,

$$a_0^i y^\varpi \Sigma \left\{ \begin{array}{l} \text{product of } \varpi \text{ differences like } \frac{x}{y} - \frac{x_r}{y_r} \\ \times \text{product of } w - \varpi \text{ differences like } \frac{x_r}{y_r} - \frac{x_s}{y_s} \end{array} \right\},$$

i. e.

$$(y_1 y_2 \dots y_p)^i y^\varpi \Sigma \left\{ \begin{array}{l} \text{product of } \varpi \text{ factors like } \frac{x y_r - x_r y}{y y_r} \\ \times \text{product of } w - \varpi \text{ factors like } \frac{x_r y_s - x_s y_r}{y_r y_s} \end{array} \right\},$$

which, as we see by consideration of the denominators, can only assume the form  $G_w$ , or

$$\Sigma \left\{ \begin{array}{l} \text{product of } \varpi \text{ factors like } x y_r - x_r y \\ \times \text{product of } w - \varpi \text{ factors like } x_r y_s - x_s y_r \end{array} \right\},$$

if, in every product of differences under the  $\Sigma$ ,  $y$  occurs in  $\varpi$  factors and  $y_1, y_2, \dots, y_p$  each in  $i$  factors, where  $p, i, w$  and  $\varpi$  are connected by the relation  $ip + \varpi = 2w$ , the left being the degree of  $(y_1 y_2 \dots y_p)^i y^\varpi$  in its arguments, and the right the degree of each product of denominators under the  $\Sigma$  in the same.

That products of the two sets of differences which cannot be prepared by a factor so as to assume the form  $G_w$  cannot produce covariants is established exactly as in § 75.

There may be functions which are sums of parts like the above, which when prepared by factors  $(y_1 y_2 \dots y_p)^i y^\varpi$  for different values of  $i$  and  $\varpi$  which satisfy  $ip + \varpi = 2w$ , produce sums of functions  $G_w$  for which  $w$  is the same. Such non-homogeneous covariants are, however, as explained in §§ 35, 36, with advantage considered as sums of covariants rather than as single covariants. Meaning then by covariant a rational integral one which is not a sum of simpler covariant parts, we may summarize as follows the conclusions at which we have arrived.

$$\begin{aligned} \text{If } u &\equiv (a_0, a_1, a_2, \dots, a_p) (x, y)^p \\ &\equiv a_0(x - a_1 y)(x - a_2 y) \dots (x - a_p y) \end{aligned}$$

be a binary  $p$ -ic of which  $a_1, a_2, \dots, a_p$  are the roots, then a function of the differences  $a_1 - a_2$ , &c. between pairs of roots,

and of the  $p$  differences  $x - a_1 y, x - a_2 y, \dots x - a_p y$ , which is homogeneous in both sets of differences and symmetric in the roots, will, when expressed in terms of  $x, y$  and the coefficients, and made integral in these latter by multiplication by the lowest necessary power of  $a_0$ , be a covariant, if and only if every one of the products of differences of which it consists involves all roots  $a$  in equal numbers of its factors. Moreover all covariants of  $u$  are given in this way.

Ex. 12. All covariants, except powers of the  $p$ -ic itself, vanish for the special  $p$ -ic  $(x + y)^p$ .

Ex. 13. The sums of the numerical coefficients of the products of

$$a_0, a_1, a_2, \dots a_p$$

which occur in  $K_0, K_1, K_2, \dots K_\varpi$ , where  $(K_0, K_1, K_2, \dots K_\varpi)(x, y)^\varpi$  is a covariant of  $(a_0, a_1, a_2, \dots a_p)(x, y)^p$ , all vanish, unless the covariant is a mere power of  $(a_0, a_1, a_2, \dots a_p)(x, y)^p$ .

Ex. 14. Every term in the summation which gives a covariant of a binary  $p$ -ic must involve at least one difference between a pair of  $r$  chosen roots if  $2ir > ip + \varpi$ .

Ans. Cf. § 39, Ex. 2.

Ex. 15. Every term in the summation which gives an invariant must have this property if  $2r > p$ .

Ans. Cf. § 26, Ex. 2.

84.] In the preceding article it is clear that  $\varpi$  is the order,  $i$  the degree, and  $w$  the weight, reckoning  $x, y$  as of weights 1, 0 respectively, of the covariant. That the facts here exhibited with regard to the order, degree and weight of a covariant, and with regard to the index in the equality expressive of covariancy, accord with the results of chap. iii may be readily seen. The relation  $ip + \varpi = 2w$  of § 83 is in fact that of § 38.

As to the index of the power of  $M$  in the equality expressive of covariancy which is derived from

$$G_w = M^w G'_w,$$

we notice that

$$\begin{aligned} G_w &= a_0^i y^\varpi F\left(\frac{a_1}{a_0}, \frac{a_2}{a_0}, \dots \frac{a_p}{a_0}; \frac{x}{y}\right) \\ &= K(a_0, a_1, a_2, \dots a_p)^i (x, y)^\varpi, \end{aligned}$$

where the notation denotes a covariant of degree  $i$  and order

$\varpi$ , and that in like manner

$$\begin{aligned} G'_w &= \left(\frac{A_0}{M^p}\right)^i Y^\varpi F\left(\frac{A_1}{A_0}, \frac{A_2}{A_0}, \dots, \frac{A_p}{A_0}; \frac{X}{Y}\right) \\ &= M^{-ip} K(A_0, A_1, A_2, \dots, A_p)^i (X, Y)^\varpi. \end{aligned}$$

Thus  $G_w = M^w G'_w = M^{\frac{1}{2}(ip+\varpi)} G'_w$  gives

$$\begin{aligned} K(A_0, A_1, A_2, \dots, A_p)^i (X, Y)^\varpi \\ = M^{\frac{1}{2}(ip-\varpi)} K(a_0, a_1, a_2, \dots, a_p) (x, y)^\varpi, \end{aligned}$$

which is the result, for  $q = 2$ , of § 37.

The index  $\frac{1}{2}(ip-\varpi)$  is, it may be here repeated,  $w-\varpi$ , where  $w$  is the weight of the covariant, and is therefore the weight of the coefficient of  $x^\varpi$  in the covariant. The weights of the successive coefficients in the covariant forms an arithmetic progression of common difference unity.

85.] **The binary quadratic.** This can have no covariant distinct from itself and its one invariant the discriminant.

In fact, if  $\alpha, \beta$  be the roots of

$$ax^2 + 2bxy + cy^2 \equiv a(x-\alpha y)(x-\beta y),$$

and if

$$a^i \Sigma \{(a-\beta)^\lambda (x-\alpha y)^\mu (x-\beta y)^\nu\}$$

be a covariant, we have, by the results of § 83, that for every product under the  $\Sigma$

(1)  $\lambda = \text{constant}$ , by homogeneity in the difference between roots;

(2)  $\mu + \nu = \text{constant}$ , by homogeneity in  $x - \alpha y, x - \beta y$ ;

(3)  $\lambda + \mu = \lambda + \nu = i$ , since  $a$  must occur in as many factors as  $\beta$ , viz. in  $i$  factors.

Thus  $\mu = \nu = \text{constant}$ , and  $\lambda = \text{constant}$ . Consequently the covariant is

$$a^{\lambda+\mu} \Sigma \{(a-\beta)^\lambda [(x-\alpha y)(x-\beta y)]^\mu\},$$

the summation consisting of two terms, corresponding to  $\alpha, \beta$  and  $\beta, \alpha$ . These two cancel if  $\lambda$  is odd, but repeat one another if  $\lambda$  is even. Hence any covariant is, but for a numerical multiple, of the form

$$\{a^2(a-\beta)^2\}^{\lambda'} \{a(x-\alpha y)(x-\beta y)\}^\mu,$$

i.e.

$$\{-4(ac-b^2)\}^{\lambda'} \{ax^2 + 2bxy + cy^2\}^\mu.$$

There are then no covariants of the quadratic which are not, but for numerical multipliers, powers of the quadratic itself, or such powers multiplied by powers of the discriminant  $ac-b^2$ .

In other words, the binary quadratic has for its only irreducible concomitants one invariant, viz. the discriminant  $ac - b^2$ , and itself.

86.] **Covariants of the cubic.** If  $a, \beta, \gamma$  be the roots of the binary cubic

$$ax^3 + 3bx^2y + 3cxy^2 + dy^3 \equiv a(x - \alpha y)(x - \beta y)(x - \gamma y),$$

the cubic has two independent covariants, numerical multiples of

$$a^2 \Sigma \{(x - \alpha y)^2 (\beta - \gamma)^2\}$$

and

$$a^3 \Sigma \{(x - \alpha y)^2 (x - \beta y)(\beta - \gamma)^2 (\gamma - \alpha)\},$$

which, writing  $\theta, \phi, \psi$  for

$$(x - \alpha y)(\beta - \gamma), (x - \beta y)(\gamma - \alpha), (x - \gamma y)(\alpha - \beta),$$

are respectively twice and once

$$a^2 (\theta^2 + \phi^2 + \psi^2) = a^2 h, \text{ say,}$$

and  $a^3 \{\theta^2 (\phi - \psi) + \phi^2 (\psi - \theta) + \psi^2 (\theta - \phi)\} = a^3 g, \text{ say.}$

That the two obey the criteria of § 83, and are consequently covariants, is at once verified. That there cannot be more than two covariants, independent of one another and the cubic itself, is known from § 42. All other covariants, and invariants too, can then be expressed in terms of them and the cubic. The one irreducible invariant, the discriminant (§ 78), is of course not a rational integral function of the three—no rational integral function of them can be free from the variables. We reserve till a later chapter the proof that there is no other irreducible concomitant of the cubic, so that any other concomitant is a rational integral function of the cubic, the two covariants above, and the discriminant.

The expressions for  $a^2 h$  and  $a^3 g$  in terms of the coefficients and variables can be effected by elementary methods of the theory of equations. We know however (cf. § 45, Exx. 12, 13) two covariants of the degrees and orders of  $a^2 h$  and  $a^3 g$ , viz.

$$H = (ac - b^2)x^2 + (ad - bc)xy + (bd - c^2)y^2,$$

and

$$G = (a^2d - 3abc + 2b^3, abd - 2ac^2 + b^2c, \\ -acd + 2b^2d - bc^2, -ad^2 + 3bcd - 2c^3)(x, y)^3,$$

and we shall see later (see, for instance, § 113, Ex. 16) that these are the only covariants of the degrees and orders in

question. Hence, for some numerical values of  $k$  and  $k'$ , we must have

$$a^2h = kH,$$

and

$$a^3g = k'G.$$

This being known, we can find  $k$  and  $k'$  by consideration of the values of the covariants in a particular case. Take for instance the cubic  $x^3 - y^3$ , for which  $a, b, c, d, \alpha, \beta, \gamma$  have the values 1, 0, 0, -1, 1,  $\omega, \omega^2$ . The above equalities become

$$18xy = -kxy,$$

$$27(x^3 + y^3) = -k'(x^3 + y^3),$$

whence the values of  $k$  and  $k'$  are -18 and -27. Thus

$$a^2h = -18H,$$

and

$$a^3g = -27G.$$

In accordance with the convention of § 71 we speak of  $H$  and  $G$ , rather than of  $a^2h$  and  $a^3g$ , as the fundamental covariants.

87.] Syzygy among concomitants of cubic. The cubic  $u$ , its two covariants  $H$  and  $G$ , and its discriminant

$$\Delta = (ad - bc)^2 - 4(ac - b^2)(bd - c^2)$$

must, as we have seen, be connected by a relation. To find this relation we may consider the cubic in the form

$$u = ax^3 + dy^3$$

which is not special, but is, as we have seen in § 11, Ex. 14, one to which the general cubic can be reduced by linear substitution. For this form

$$H = adxy,$$

$$G = a^2dx^3 - ad^2y^3 = ad(ax^3 - dy^3),$$

$$\Delta = a^2d^2;$$

whence, by elimination of  $a, d, x, y$ , the connecting relation is seen to be

$$\Delta u^2 = G^2 + 4H^3,$$

which holds when the general expressions for  $u, G, H$  and  $\Delta$  are substituted.

Ex. 16. Prove the same relation by showing that  $\theta^2\phi^2\psi^2$ , a determinate numerical multiple of the product  $\Delta u^2$ , is  $3^{-6}$  times the product of the squares of the differences between roots of the equation  $z^3 - \frac{3}{2}hz + g = 0$  whose roots are  $\phi - \psi, \psi - \theta, \theta - \phi$ , and so a determinate numerical multiple of the discriminant of this cubic.



Ex. 17. If the roots of the cubic equation  $u = 0$  are  $a, \beta, \gamma$ , the roots of  $H = 0$  are

$$-\frac{\beta\gamma + \omega\gamma a + \omega^2 a\beta}{a + \omega\beta + \omega^2\gamma}, \quad -\frac{\beta\gamma + \omega^2\gamma a + \omega a\beta}{a + \omega^2\beta + \omega\gamma},$$

and those of  $G = 0$  are

$$\frac{2\beta\gamma - \gamma a - a\beta}{\beta + \gamma - 2a}, \quad \frac{2\gamma a - a\beta - \beta\gamma}{\gamma + a - 2\beta}, \quad \frac{2a\beta - \beta\gamma - \gamma a}{a + \beta - 2\gamma}. \quad (\text{Cayley.})$$

88.] The close similarity between the forms, in terms of differences, of the covariants of a cubic and the invariants of a quartic will not have escaped notice. It is not accidental, but is a result of the fact, to which attention has been called in § 69, that the invariants of

$$(xy' - x'y)u$$

are, when  $x'$  and  $y'$  are replaced by  $x$  and  $y$ , covariants of  $u$ .

Any consideration of covariants of binary quantics above the third order, and of invariants of binary quantics above the fourth, is postponed till later chapters.

89.] **Several binary quantics.** Into a full discussion of the expressions by means of differences of invariants and covariants of systems of more binary quantics than one it is not proposed here to enter; but the facts may be developed by the same methods as have been adopted in this chapter.

It will be found, for instance, with regard to *invariants* of *two* binary quantics, that functions of the roots which produce them must of course, for rationality, be symmetrical in the roots of each quantic separately, and will in general be functions of three classes of differences, viz. (1) differences between two roots of the first quantic, (2) differences between two roots of the second quantic, and (3) differences between a root of the first and one of the second. In order to produce invariants which are not more properly regarded as sums of simpler invariants, such functions must be homogeneous, not only on the whole, but in each of the three sets of differences singly. Any one must, moreover, be a sum of products of differences, in every one of which all roots of the first quantic occur in equal constant numbers of factors, and all roots of the second in equal constant numbers of factors, the numbers not being, however, necessarily or as a rule the same.

## CHAPTER VI.

### BINARY QUANTICS CONTINUED. ANNIHILATORS. SEMINVARIANTS.

90.] **Annihilators of invariants.** For the calculation of invariants it is a matter of great importance that  $I$  any invariant of  $(a_0, a_1, a_2, \dots, a_p)(x, y)^p$  must satisfy the two differential equations

$$a_0 \frac{dI}{da_0} + 2a_1 \frac{dI}{da_2} + 3a_2 \frac{dI}{da_3} + \dots + pa_{p-1} \frac{dI}{da_p} = 0,$$

$$pa_1 \frac{dI}{da_0} + (p-1)a_2 \frac{dI}{da_1} + (p-2)a_3 \frac{dI}{da_2} + \dots + a_p \frac{dI}{da_{p-1}} = 0.$$

Professor Sylvester, to whom and to Cayley the theory is due, though the idea had also presented itself to Aronhold, expresses this fact by saying that any invariant  $I$  has two *annihilators*, called  $\Omega$  and  $O$ , viz.

$$\Omega \equiv a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + 3a_2 \frac{d}{da_3} + \dots + pa_{p-1} \frac{d}{da_p},$$

$$O \equiv pa_1 \frac{d}{da_0} + (p-1)a_2 \frac{d}{da_1} + (p-2)a_3 \frac{d}{da_2} + \dots + a_p \frac{d}{da_{p-1}}.$$

The language is a convenient one for expressing that  $\Omega I = 0$ , and  $OI = 0$ .

We proceed to prove these facts of annihilation.

91.] **The annihilator  $\Omega$ .** The property of having  $\Omega$  for an annihilator is one that invariants possess in common with other functions of the coefficients which, when expressed in terms of  $a_0$  and the roots, involve only differences of these latter.

This may be proved by seeing, as we shall later, that the

operation with  $\Omega$  on a function of the coefficients is equivalent to the operation with

$$-\left(\frac{d}{da_1} + \frac{d}{da_2} + \dots + \frac{d}{da_p}\right)$$

on the equal function of  $a_0$  and the roots  $a_1, a_2, \dots, a_p$ . We adopt here, however, a different method.

Functions of the coefficients which are equivalent to functions of  $a_0$  and differences between roots are, in fact, equal to the same functions of the altered coefficients when the quantic is transformed by the substitution of  $X + mY$  and  $Y$  for  $x$  and  $y$ , that is to say, when the roots are diminished by the same (positive or negative) quantity  $m$ . Now, this being so for all values of  $m$ , let  $m$  be taken as very small, so that its square and higher powers may be neglected in comparison with any finite multiple of itself. The quantic

$$a_0x^p + pa_1x^{p-1}y + \frac{p(p-1)}{1 \cdot 2}a_2x^{p-2}y^2 + \dots + a_px^py^p$$

becomes in this case, after the substitution,

$$a_0X^p + p(a_1 + ma_0)X^{p-1}Y + \frac{p(p-1)}{1 \cdot 2}(a_2 + 2ma_1)X^{p-2}Y^2 + \dots + (a_p + pma_{p-1})Y^p,$$

so that the new coefficients are the old ones altered by the increments

$$\delta a_0 = 0, \delta a_1 = ma_0, \delta a_2 = 2ma_1, \dots, \delta a_p = pma_{p-1}.$$

Now  $I$  our supposed invariant, or other function of  $a_0$  and the differences between roots, becomes, by Taylor's theorem

$$I + \left(\delta a_0 \frac{d}{da_0} + \delta a_1 \frac{d}{da_1} + \delta a_2 \frac{d}{da_2} + \dots + \delta a_p \frac{d}{da_p}\right) I,$$

in which quadratic, &c. terms in the  $\delta a$ 's, i. e. in  $m$ , are omitted as vanishing in comparison with the increment retained.

Thus a necessary result of  $I$  being unaltered is that

$$\left(\delta a_0 \frac{d}{da_0} + \delta a_1 \frac{d}{da_1} + \delta a_2 \frac{d}{da_2} + \dots + \delta a_p \frac{d}{da_p}\right) I = 0,$$

i. e. that  $m(a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + \dots + pa_{p-1} \frac{d}{da_p}) I = 0,$

i. e. that  $\Omega I = 0.$

92.] It will be well to give another proof of this, both because of the convenient symbolical form of results to which it will lead, and in order to convince ourselves that  $\Omega I = 0$  is a sufficient condition to ensure that  $I$  is a function equal to the same function of the coefficients in the quantic obtained by putting  $X + mY$ ,  $Y$  for  $x$  and  $y$  in the given quantic, whatever constant  $m$  be, as well as a necessary consequence if this persistence is a fact.

If

$(A_0, A_1, A_2, \dots, A_p)(X, Y)^p = (a_0, a_1, a_2, \dots, a_p)(X + mY, Y)^p$ ,  
 where  $m$  is not now necessarily very small, the expressions for the new coefficients are easily seen to be, by use of Taylor's theorem for the expansion of a function of  $\frac{X}{Y} + m$  in powers of  $\frac{X}{Y}$ ,

$$A_0 = a_0,$$

$$A_1 = a_1 + a_0 m,$$

$$A_2 = a_2 + 2a_1 m + a_0 m^2,$$

$$\dots \dots \dots$$

$$A_p = a_p + p a_{p-1} m + \frac{p(p-1)}{1 \cdot 2} a_{p-2} m^2 + \dots + a_0 m^p,$$

where we notice that

$$\frac{dA_0}{dm} = 0, \quad \frac{dA_1}{dm} = A_0, \quad \frac{dA_2}{dm} = 2A_1, \quad \frac{dA_3}{dm} = 3A_2, \dots, \quad \frac{dA_p}{dm} = pA_{p-1}.$$

We draw the conclusion that, if  $F(A_0, A_1, A_2, \dots, A_p)$  be any function of the new coefficients,

$$\begin{aligned} & \frac{d}{dm} F(A_0, A_1, A_2, \dots, A_p) \\ &= \frac{dF}{dA_0} \cdot \frac{dA_0}{dm} + \frac{dF}{dA_1} \cdot \frac{dA_1}{dm} + \frac{dF}{dA_2} \cdot \frac{dA_2}{dm} + \dots + \frac{dF}{dA_p} \cdot \frac{dA_p}{dm} \\ &= A_0 \frac{dF}{dA_1} + 2A_1 \frac{dF}{dA_2} + 3A_2 \frac{dF}{dA_3} + \dots + pA_{p-1} \frac{dF}{dA_p}. \end{aligned}$$

Now for  $\frac{d}{dm} F(A_0, A_1, A_2, \dots, A_p)$  to vanish, whatever  $m$  be, is the necessary and sufficient condition that

$$F(A_0, A_1, A_2, \dots, A_p)$$

be independent of  $m$ , and so equal to  $F(a_0, a_1, a_2, \dots a_p)$  which is its value when  $m = 0$ . Thus the condition, both sufficient and necessary, that

$$F(A_0, A_1, A_2, \dots A_p) = F(a_0, a_1, a_2, \dots a_p)$$

$$\text{is } \left( A_0 \frac{d}{dA_1} + 2A_1 \frac{d}{dA_2} + 3A_2 \frac{d}{dA_3} + \dots + pA_{p-1} \frac{d}{dA_p} \right) F(A_0, A_1, A_2, \dots A_p) = 0,$$

or, replacing capital by small letters,

$$\Omega F(a_0, a_1, a_2, \dots a_p) = 0.$$

93.] We can by this method prove that, if  $F$  be any rational integral function,

$$F(A_0, A_1, A_2, \dots A_p) = \left( 1 + m\Omega + \frac{m^2}{1.2} \Omega^2 + \frac{m^3}{1.2.3} \Omega^3 + \dots \right) F(a_0, a_1, a_2, \dots a_p),$$

which may be written symbolically

$$= e^{m\Omega} F(a_0, a_1, a_2, \dots a_p).$$

In fact we have, by Maclaurin's theorem,

$$\begin{aligned} \phi(m') &= \phi(0) + m' \left[ \frac{d\phi}{dm} \right]_0 + \frac{m'^2}{1.2} \left[ \frac{d^2\phi}{dm^2} \right]_0 + \dots \\ &= \left[ e^{m' \frac{d}{dm}} \phi(m) \right]_{m=0}. \end{aligned}$$

Now  $F(A_0, A_1, A_2, \dots A_p)$  is a function of  $m$ . Hence, if  $A'_r$  denote the result of replacing  $m$  by  $m'$  in  $A_r$ , we have by the preceding article

$$\begin{aligned} &F(A'_0, A'_1, A'_2, \dots A'_p) \\ &= \left[ e^{m' \left( A_0 \frac{d}{dA_1} + 2A_1 \frac{d}{dA_2} + \dots + pA_{p-1} \frac{d}{dA_p} \right)} F(A_0, A_1, A_2, \dots A_p) \right]_{m=0}, \\ &= e^{m'\Omega} F(a_0, a_1, a_2, \dots a_p). \end{aligned}$$

This is proved, subject to considerations of convergency, for any function  $F$ . When  $F$  is a rational integral function no question of convergency arises. For we notice that  $F, \Omega F, \Omega^2 F, \dots$  are of weights regularly diminishing by unity, so that presently we get to a term  $\Omega^w F$  of zero weight, i. e. a function of  $a_0$ , and beyond this point  $\Omega^{w+1} F, \Omega^{w+2} F, \&c.$  all

vanish. The symbolic series practically consists therefore of only a finite number of terms.

We have thus another proof that, if a function  $I$  persists in form after the substitution of  $X + mY$ ,  $Y$  for  $x$  and  $y$  when  $m$  is infinitesimal, it does equally when  $m$  is finite. For if  $\Omega I = 0$ , the condition of § 91, then also  $\Omega^2 I = \Omega \Omega I = 0$ ,  $\Omega^3 I = 0$ , &c., &c.

Of course the student will recognize that  $\Omega^2 I$  denotes the full expression for the result of operating with  $\Omega$  on  $\Omega I$ , viz.

$$a_0^2 \frac{d^2 I}{da_1^2} + 4 a_0 a_1 \frac{d^2 I}{da_1 da_2} + 6 a_0 a_2 \frac{d^2 I}{da_1 da_3} + \dots + 4 a_1^2 \frac{d^2 I}{da_2^2} + \dots \\ + 1 \cdot 2 a_0 \frac{dI}{da_2} + 2 \cdot 3 a_1 \frac{dI}{da_3} + 3 \cdot 4 a_2 \frac{dI}{da_4} + \dots,$$

and not merely the first line of this expression; and so for  $\Omega^3 I$ ,  $\Omega^4 I$ , &c.

94.] The annihilator  $O$ . We have still to see that, if  $I$  be an invariant, the second operator  $O$  of § 90 is an annihilator of  $I$ , as well as the first  $\Omega$ .

This property invariants have in common with other functions of the coefficients which persist in form after the substitution

$$x = X, \quad y = \nu X + Y,$$

i. e. in common with all functions of the coefficients which can be expressed in terms of  $a_p$  and the differences between reciprocals of roots. The substitution is in fact one which transforms the quantic into another in which  $a_p$  is unaltered and the reciprocals of the roots differ by  $\nu$  from the reciprocals of the original roots.

The proof is exactly as before, the present substitution dealing with  $y$  and  $x$  exactly as that of the preceding articles has dealt with  $x$  and  $y$ , and consequently dealing with the quantic read backwards from its end  $a_p y^p$ , exactly as the former substitution dealt with it read forwards from its beginning  $a_0 x^p$ . It will be noticed that  $O$  exactly corresponds to  $\Omega$  in this reversed reading.

We have then that  $OF = 0$  is the necessary and sufficient condition that  $F$  persist in form when for  $x$  and  $y$  we make such substitutions as  $X, \nu X + Y$ .

95.] **Symmetry of an invariant. Skew invariants.** If in an invariant  $a_0$  and  $a_p$ ,  $a_1$  and  $a_{p-1}$ ,  $a_2$  and  $a_{p-2}$ , &c. be interchanged, the invariant is unaltered if its weight be even, and changed only in sign if its weight be odd.

For the substitution  $x = Y$ ,  $y = X$  has for its modulus  $-1$ . Now the effect of it is to interchange  $a_0$  and  $a_p$ ,  $a_1$  and  $a_{p-1}$ ,  $a_2$  and  $a_{p-2}$ , &c. in the quantic. Consequently, if

$$F(a_0, a_1, a_2, \dots, a_p)$$

be an invariant, we have (§§ 23, 26, 76)

$$F(a_p, a_{p-1}, a_{p-2}, \dots, a_0) = (-1)^w F(a_0, a_1, a_2, \dots, a_p).$$

We see then that there is an essential difference in character between invariants of even and invariants of odd weight. Those of odd weight are, because of this change of sign, known as *skew invariants*.

Skew invariants do not exist for the quadratic, cubic and quartic, and it came as a surprise upon mathematicians when Hermite discovered the first skew invariant of a higher quantic; viz. that of degree 18 and weight 45 of the quintic.

Invariants of odd weight cannot, it is clear, be rational integral functions of invariants of even weight. Thus when a binary quantic has one or more skew invariants one at least of them must be irreducible.

96.] One result of the symmetry to which attention has just been called is that when we have found a function of the coefficients which has  $\Omega$  for an annihilator it is unnecessary to test directly whether it is also annihilated by  $O$  in order to ascertain whether it is or is not an invariant. If it is altered in more than sign when the first coefficient and last in the quantic, the second and last but one, &c. are interchanged in pairs, it is not an invariant. If on the other hand it is not so altered in more than sign it is certainly annihilated by  $O$  as well as by  $\Omega$ , for  $O$  is what  $\Omega$  becomes when these interchanges are made.

We must prove, however, that any function which is of one order and one weight throughout, and which is annihilated both by  $\Omega$  and by  $O$ , is an invariant.

97.] A homogeneous isobaric function annihilated by  $\Omega$  and by  $O$  is necessarily an invariant.

Consider in succession the substitutions

$$\left. \begin{aligned} x &= \lambda x', \\ y &= \mu y', \end{aligned} \right\} \dots (1)$$

$$\left. \begin{aligned} x' &= X' + tY', \\ y' &= Y', \end{aligned} \right\} \dots (2)$$

$$\left. \begin{aligned} X' &= X, \\ Y' &= \tau X + Y. \end{aligned} \right\} \dots (3)$$

The result of the succession is that of the performance of the substitutions

$$\left. \begin{aligned} x &= \lambda X + \lambda t (\tau X + Y) \\ &= \lambda (1 + t\tau) X + \lambda t Y, \\ y &= \mu \tau X + \mu Y. \end{aligned} \right\} \dots (4)$$

Now these are the most general formulae of linear substitution; for,  $\lambda, \mu, t, \tau$  being arbitrary, so are the coefficients

$$\lambda (1 + t\tau), \lambda t, \mu\tau, \mu,$$

as is clear by taking them in reversed order. The modulus of the resultant substitution (4) is  $\lambda\mu$ .

Let the original form of a  $p$ -ic be

$$(a_0, a_1, a_2, \dots a_p) (x, y)^p,$$

and let the forms it successively takes be

$$\begin{aligned} &(a'_0, a'_1, a'_2, \dots a'_p) (x', y')^p, \\ &(A'_0, A'_1, A'_2, \dots A'_p) (X', Y')^p, \\ &(A_0, A_1, A_2, \dots A_p) (X, Y)^p. \end{aligned}$$

Take  $F(a_0, a_1, a_2, \dots a_p)$  a homogeneous isobaric function, of degree  $i$  and weight  $w$ , which is annihilated by  $\Omega$  and by  $O$ .

We have first that, for values of  $r$  from 0 to  $p$  inclusive,

$$a'_r = \lambda^{p-r} \mu^r a_r,$$

where the index of the power of  $\mu$  is the weight of  $a_r$ , and that of the power of  $\lambda$  is the excess of  $p$  over that weight. Accordingly, because  $F$  is homogeneous and isobaric,

$$F(a'_0, a'_1, a'_2, \dots a'_p) = \lambda^{ip-w} \mu^w F(a_0, a_1, a_2, \dots a_p).$$



Again, because  $\Omega$  annihilates  $F$ ,  $F$  persists in form after the substitution (2). Therefore

$$F(A_0', A_1', A_2', \dots, A_p') = F(a_0', a_1', a_2', \dots, a_p').$$

(2) is the subst  
where  $\Omega$   
derived  
3: is the subst  
where  $\Omega$

Once more, because  $O$  annihilates  $F$ ,

$$F(A_0, A_1, A_2, \dots, A_p) = F(A_0', A_1', A_2', \dots, A_p').$$

We see then, taking these three facts together, that

$$F(A_0, A_1, A_2, \dots, A_p) = \lambda^{ip-w} \mu^w F(a_0, a_1, a_2, \dots, a_p). \quad \dots (5)$$

In other words, we see that  $F$  is a function of the coefficients, which needs only to be multiplied by a factor involving only the constants in the general scheme of linear substitution (4) to be made equal to the same function of the coefficients in the quantic into which the given quantic is transformed by that substitution. By the definition then  $F$  is an invariant.

Moreover the fact (§ 26) that  $w = \frac{1}{2}ip$  follows. For, by § 23, the factor  $\lambda^{ip-w} \mu^w$  in (5) must be a power of the modulus, i. e. of  $\lambda\mu$ . Thus the indices of  $\lambda^{ip-w}$  and  $\mu^w$  must be equal. Therefore

$$ip - w = w,$$

i. e.  $w = \frac{1}{2}ip$ .

Another interesting proof that, when  $\Omega$  and  $O$  both annihilate a homogeneous isobaric function  $F$ , the weight and degree of  $F$  must be connected with  $p$  by the relation  $ip - 2w = 0$ , will be afforded when we have seen in the next chapter that

$$(\Omega O - O \Omega) F = (ip - 2w) F.$$

For the left-hand member vanishes when  $\Omega F = 0$  and  $OF = 0$ . So then must the right-hand member.

98.] A good proof in small compass of all the fundamental properties of invariants of a binary quantic is afforded by a method which will also be useful for other purposes.

We notice that, if

$$u = (a_0, a_1, a_2, \dots, a_p) (x, y)^p,$$

then  $\Omega u = y \frac{d}{dx} u$ ,  $\Omega^2 u = \left(y \frac{d}{dx}\right)^2 u, \dots$

and, generally,  $\Omega^r u = \left(y \frac{d}{dx}\right)^r u.$

Consequently

$$e^{t\Omega} (a_0, a_1, a_2, \dots, a_p) (x, y)^p = e^{ty} \frac{d}{dx} (a_0, a_1, a_2, \dots, a_p) (x, y)^p \\ = (a_0, a_1, a_2, \dots, a_p) (x + ty, y)^p,$$

by Taylor's theorem.

Similarly

$$Ou = x \frac{d}{dy} u, O^2u = \left(x \frac{d}{dy}\right)^2 u, \dots$$

and, generally,  $O^r u = \left(x \frac{d}{dy}\right)^r u;$

and therefore

$$e^{\tau O} (a_0, a_1, a_2, \dots, a_p) (x, y)^p = e^{\tau x} \frac{d}{dy} (a_0, a_1, a_2, \dots, a_p) (x, y)^p \\ = (a_0, a_1, a_2, \dots, a_p) (x, \tau x + y)^p.$$

Hence, performing one operation after the other,

$$e^{t\Omega} e^{\tau O} (a_0, a_1, a_2, \dots, a_p) (x, y)^p \\ = (a_0, a_1, a_2, \dots, a_p) (x + t(\tau x + y), \tau x + y)^p,$$

and, putting  $\lambda x, \mu y$  for  $x, y,$

$$e^{t\Omega} e^{\tau O} (a_0, a_1, a_2, \dots, a_p) (\lambda x, \mu y)^p \\ = (a_0, a_1, a_2, \dots, a_p) ((1 + t\tau)\lambda x + t\mu y, \tau\lambda x + \mu y)^p,$$

i.e., taking

$$(1 + t\tau)\lambda = l, t\mu = m, \tau\lambda = l', \mu = m',$$

so that

$$\mu = m', t = \frac{m}{m'}, \tau = \frac{l'm'}{lm' - l'm}, \lambda = \frac{lm' - l'm}{m'},$$

$$(a_0, a_1, a_2, \dots, a_p) (lx + my, l'x + m'y)^p \\ = e^{\frac{m}{m'}\Omega} e^{\frac{l'm'}{lm' - l'm}O} (a_0, a_1, a_2, \dots, a_p) \left(\frac{lm' - l'm}{m'}x, m'y\right)^p.$$

Thus the most general linear substitution of  $lx + my, l'x + m'y$  for  $x$  and  $y$  is effected by a substitution of the form  $\lambda x, \mu y$  followed by a complex differential operation.

This is an identity. If then the expanded left be

$$(A_0, A_1, A_2, \dots, A_p) (x, y)^p,$$

we have for all values of  $r$  from 0 to  $p$  inclusive

$$A_r = e^{\frac{m}{m'}\Omega} e^{\frac{l'm'}{lm' - l'm}O} \left(\frac{lm' - l'm}{m'}\right)^{p-r} m'^r a_r \\ = (lm' - l'm)^{p-r} m'^{2r-p} e^{\frac{m}{m'}\Omega} e^{\frac{l'm'}{lm' - l'm}O} a_r,$$

so that we have a formula for every new coefficient.

We can readily pass to products of coefficients. For, if  $P, Q$  be two functions of the coefficients,

$$e^{k0} PQ = e^{k(O_1+O_2)} PQ,$$

where  $O_1$  and  $O_2$  both mean the same as  $O$ , but the former and its repetitions act on  $P$  only, and the latter and its repetitions on  $Q$  only,

$$\begin{aligned} &= e^{kO_1} e^{kO_2} PQ \\ &= e^{kO_1} P \cdot e^{kO_2} Q \\ &= e^{k0} P \cdot e^{k0} Q; \end{aligned}$$

and, in like manner,

$$\begin{aligned} e^{k'\Omega} e^{k0} PQ &= e^{k'\Omega} (e^{k0} P \cdot e^{k0} Q) = \{ \begin{matrix} k'(\Omega, \Omega') \\ \{ \end{matrix} \} [ \{ \begin{matrix} k0 \\ \{ \end{matrix} \} P \cdot \{ \begin{matrix} k0 \\ \{ \end{matrix} \} Q ] \\ &= e^{k'\Omega} e^{k0} P \cdot e^{k'\Omega'} e^{k0} Q. = \{ \begin{matrix} k'a \\ \{ \end{matrix} \} \{ \begin{matrix} k0 \\ \{ \end{matrix} \} P \cdot \{ \begin{matrix} k'a \\ \{ \end{matrix} \} \{ \begin{matrix} k0 \\ \{ \end{matrix} \} Q \end{aligned}$$

Hence, if  $F(a_0, a_1, a_2, \dots a_p)$  be a product, or a sum of multiples of products of the same degree  $i$  and weight  $w$ , of coefficients chosen from among  $a_0, a_1, a_2, \dots a_p$ ,

$$\begin{aligned} &F(A_0, A_1, A_2, \dots A_p) \\ &= (lm' - l'm)^{ip-w} m'^{2w-ip} e^{\frac{m}{m'}\Omega} e^{\frac{l'm'}{lm'-l'm}O} F(a_0, a_1, a_2, \dots a_p), \end{aligned}$$

so that any rational integral homogeneous isobaric function of the new coefficients  $A$  is obtained from the same function of the old coefficients  $a$  by a complex differential operation and a multiplication.

By reversing the order of the  $\Omega$  and  $O$  operations we obtain in like manner a second expression for  $F(A_0, A_1, A_2, \dots A_p)$ ; viz.

$$\begin{aligned} &F(A_0, A_1, A_2, \dots A_p) \\ &= (lm' - l'm)^w l^{ip-2w} e^{\frac{l'}{l}O} e^{\frac{lm}{lm'-l'm}\Omega} F(a_0, a_1, a_2, \dots a_p). \end{aligned}$$

99.] All the fundamental facts as to invariants flow hence. These are that an invariant is annihilated by  $\Omega$  and by  $O$ , that its degree and weight are connected with  $p$ , the order of the quantic, by the relation  $ip - 2w = 0$ , and that the factor in the equality expressive of its invariancy is the  $w$ th power of the modulus  $lm' - l'm$ .

To see this, suppose that  $F(a_0, a_1, a_2, \dots a_p)$ , a homogeneous isobaric function, is an invariant, so that

$$F(A_0, A_1, A_2, \dots A_p) = \phi(l, m, l', m') F(a_0, a_1, a_2, \dots a_p),$$

where the form of  $\phi$  is at present unknown. We have then three expressions for  $F(A_0, A_1, A_2, \dots A_p)$ , which we can identify, and obtain

$$\begin{aligned} (lm' - l'm)^{ip-w} m'^{2w-ip} e^{\frac{m}{m'}\Omega} e^{\frac{l'm'}{lm'-l'm}O} F(\alpha_0, \alpha_1, \alpha_2, \dots \alpha_p) \\ = \phi(l, m, l', m') F(\alpha_0, \alpha_1, \alpha_2, \dots \alpha_p) \\ = (lm' - l'm)^w l^{ip-2w} e^{\frac{l'}{l}O} e^{\frac{lm}{lm'-l'm}\Omega} F(\alpha_0, \alpha_1, \alpha_2, \dots \alpha_p). \end{aligned}$$

In the first equality put  $l' = 0$ . It becomes

$$\begin{aligned} l^{ip-w} m'^w e^{\frac{m}{m'}\Omega} F(\alpha_0, \alpha_1, \alpha_2, \dots \alpha_p) \\ = \phi(l, m, 0, m') F(\alpha_0, \alpha_1, \alpha_2, \dots \alpha_p). \end{aligned}$$

Now  $\Omega F$  is of lower weight than  $F$ , and  $\Omega^2 F$ ,  $\Omega^3 F$ , &c. of lower weights still. The terms of different weights on the two sides must be separately equal. Hence

$$\Omega F = 0, \Omega^2 F = 0, \&c.$$

Again, put  $m = 0$ . We obtain in like manner

$$\begin{aligned} l^{ip-w} m'^w e^{\frac{l'}{l}O} F(\alpha_0, \alpha_1, \alpha_2, \dots \alpha_p) \\ = \phi(l, 0, l', m') F(\alpha_0, \alpha_1, \alpha_2, \dots \alpha_p); \end{aligned}$$

whence, by considering terms of different weights, since operation with  $O$  increases weight,

$$OF = 0, O^2 F = 0, \&c.$$

Thus the facts that an invariant is annihilated by  $\Omega$  and by  $O$  are obtained. This being so, the general equalities become

$$(lm' - l'm)^{ip-w} m'^{2w-ip} F = \phi(l, m, l', m') F = (lm' - l'm)^w l^{ip-2w} F,$$

in which  $l$ ,  $m'$  and  $lm' - l'm$  are independent. The equality of the first and third expressions requires then that the indices of  $m'^{2w-ip}$  and  $l^{ip-2w}$  vanish, and that the indices of  $(lm' - l'm)^{ip-w}$  and  $(lm' - l'm)^w$  be equal. These are all satisfied if and only if

$$ip - 2w = 0.$$

Lastly, using this fact, the value of  $\phi(l, m, l', m')$  is  $(lm' - l'm)^w$ .

100.] A similar analysis will lead us to a theorem of great importance which we shall use hereafter.

By the early part of § 98 we have, putting  $-\tau^{-1}$  for  $t$ ,

$$\begin{aligned} e^{-\tau^{-1}\Omega} e^{\tau O} (a_0, a_1, a_2, \dots a_p) (x, y)^p \\ &= (a_0, a_1, a_2, \dots a_p) (-\tau^{-1}y, \tau x + y)^p \\ &= e^{-\tau O} (a_0, a_1, a_2, \dots a_p) (-\tau^{-1}y, \tau x)^p \\ &= e^{-\tau O} (a_p, a_{p-1}, a_{p-2}, \dots a_0) (\tau x, -\tau^{-1}y)^p; \end{aligned}$$

whence, equating the coefficients of  $x^{p-r}y^r$  on the two sides

$$e^{-\tau^{-1}\Omega} e^{\tau O} a_r = (-1)^r \tau^{p-2r} e^{-\tau O} a_{p-r},$$

for every value of  $r$  from 0 to  $p$  inclusive.

We may hence pass to products, and linear functions of products, of coefficients  $a_0, a_1, a_2, \dots a_p$  as in § 98, and obtain that, if  $F(a_0, a_1, a_2, \dots a_p)$  be any rational integral homogeneous isobaric function of degree  $i$  and weight  $w$ ,

$$\begin{aligned} e^{-\tau^{-1}\Omega} e^{\tau O} F(a_0, a_1, a_2, \dots a_p) \\ &= (-1)^w \tau^{ip-2w} e^{-\tau O} F(a_p, a_{p-1}, a_{p-2}, \dots a_0), \end{aligned}$$

i. e.

$$\begin{aligned} (1 - \tau^{-1}\Omega + \tau^{-2} \frac{\Omega^2}{1 \cdot 2} - \dots) (1 + \tau O + \tau^2 \frac{O^2}{1 \cdot 2} + \dots) F(a_0, a_1, a_2, \dots a_p) \\ &= (-1)^w \tau^{ip-2w} (1 - \tau O + \tau^2 \frac{O^2}{1 \cdot 2} - \dots) F(a_p, a_{p-1}, a_{p-2}, \dots a_0). \end{aligned}$$

Now equate terms free from  $\tau$  on the two sides, as we may do since the equality is identical, holding for all values of  $\tau$ . The conclusion is that

$$\begin{aligned} (1 - \frac{\Omega O}{1^2} + \frac{\Omega^2 O^2}{1^2 \cdot 2^2} - \frac{\Omega^3 O^3}{1^2 \cdot 2^2 \cdot 3^2} + \dots) F(a_0, a_1, a_2, \dots a_p) \\ &= 0, \text{ if } ip - 2w > 0, \end{aligned}$$

but  $= (-1)^w F(a_p, a_{p-1}, a_{p-2}, \dots a_0)$ , if  $ip - 2w = 0$ ,

and  $= (-1)^{ip-w} \frac{O^{2w-ip}}{(2w-ip)!} F(a_p, a_{p-1}, a_{p-2}, \dots a_0)$ , if  $ip - 2w < 0$ .

The first part of the conclusion is the one to which we wish to draw particular attention. It tells us that, if  $ip - 2w > 0$ ,

$$\begin{aligned} F(a_0, a_1, a_2, \dots a_p) \\ &= \Omega \left\{ \frac{O}{1^2} - \frac{\Omega O^2}{1^2 \cdot 2^2} + \frac{\Omega^2 O^3}{1^2 \cdot 2^2 \cdot 3^2} - \dots \right\} F(a_0, a_1, a_2, \dots a_p), \end{aligned}$$

7 not in int.

i.e. that any rational integral homogeneous isobaric function, for which  $ip > 2w$ , can be obtained by operation with  $\Omega$  on another rational integral homogeneous isobaric function.

101.] **Formation of invariants by aid of  $\Omega$ .** We return to invariants. A rational integral invariant of the binary  $p$ -ic is, we have seen, a rational integral homogeneous isobaric function, whose degree and weight are connected with  $p$  by the relation  $ip = 2w$ , and which is annihilated by  $\Omega$ . These two requirements are necessary, and we shall see in § 112 that they suffice. They lead to the following method of formation of all invariants of given degree  $i$ .

Write down all possible products of  $i$  factors chosen from among  $a_0, a_1, a_2, \dots, a_p$ , repeated factors being allowed, which are such that the sum of the suffixes in every one is  $\frac{1}{2} ip$ , and take the sum of arbitrary multiples of those products. Operate on the sum with  $\Omega$ , and express that the result vanishes. This will give a number of equations in the arbitrary multipliers, since the multiplier of every distinct product of  $a_0, a_1, a_2, \dots, a_p$  in the result of operating must vanish separately. If the number of the arbitrary multipliers be not greater than the number of independent equations to be satisfied, values of them different from zero cannot be found to accord with the requirements, and there is no invariant of degree  $i$ . If the number of independent equations be one less than the number of arbitrary multipliers, the ratios of these can be chosen in one way to satisfy them, and there is one invariant. If the number of independent equations be more than one less than the number of multipliers the equations can be satisfied in more than one way. In fact, if the excess of the one number over the other be  $r$ ,  $r$  of the multipliers may be left arbitrary, and the equations can still be satisfied by proper choice of the rest. We thus get an invariant

$$\lambda_1 I_1 + \lambda_2 I_2 + \dots + \lambda_r I_r,$$

where  $\lambda_1, \lambda_2, \dots, \lambda_r$  are arbitrary. This is expressed by saying that there are  $r$  linearly independent invariants  $I_1, I_2, \dots, I_r$  of degree  $i$ .

It will be proved later, by means of the last article, that all the equations for determining the multipliers are independent,

so that the number of linearly independent invariants of degree  $i$  is the excess of the number of products of  $a_0, a_1, a_2, \dots, a_p$ , of degree  $i$  and weight  $\frac{1}{2}ip$ , over the number of products which occur in the results of operating with  $\Omega$  on these products, i.e. over the number of products of degree  $i$  and weight  $\frac{1}{2}ip-1$ . By deferring the proof we shall avoid repetition, as the theorem is a case of a more general one which will be required later.

Notice that the number of products of degree  $i$  and weight  $w$  is the number of ways in which the number  $w$  can be formed by the addition of  $i$  or fewer numbers, none exceeding  $p$ . This number of partitions of  $w$  is usually denoted by  $(w; i, p)$ . Thus the number of linearly independent invariants of degree  $i$  is

$$\left(\frac{ip}{2}; i, p\right) - \left(\frac{ip}{2} - 1; i, p\right).$$

102.] As an example of this method let us prove the statement made in § 80, that  $I$  and  $J$  are the only invariants of degrees 2 and 3 respectively of the binary quartic.

Here  $p = 4$ . Take first  $i = 2$ . Then  $w = \frac{1}{2}2 \cdot 4 = 4$ . Now the only partitions of 4 into two or fewer parts, none exceeding 4, are  $0+4, 1+3, 2+2$ . The only possible terms in an invariant of degree 2 are, then,  $a_0a_4, a_1a_3, a_2^2$ ; or  $ae, bd, c^2$  say. Now suppose that  $ae + \lambda bd + \mu c^2$  is an invariant. The result of operating on it with  $\Omega$ , i.e. with  $a\delta_b + 2b\delta_c + 3c\delta_d + 4d\delta_e$ , where, for instance,  $\delta_b$  means  $\frac{d}{db}$ , is

$$ad(4 + \lambda) + bc(3\lambda + 4\mu);$$

for which to vanish we must have  $4 + \lambda = 0$  and  $3\lambda + 4\mu = 0$ , i.e.

$$\lambda = -4, \mu = 3.$$

Thus

$$I \equiv ae - 4bd + 3c^2$$

is the only invariant of degree 2, any other being merely a numerical multiple of it.

Again, take  $i = 3$ , so that  $w = \frac{1}{2}3 \cdot 4 = 6$ . The only partitions of 6 to be dealt with are

$$0+2+4, 0+3+3, 1+1+4, 1+2+3, 2+2+2.$$

The necessary form is then

$$ace + \lambda ad^2 + \mu b^2e + \nu bcd + \rho c^3,$$

and  $\Omega$  operating on this produces

$$abe \begin{vmatrix} 2 \\ + 2\mu \end{vmatrix} + acd \begin{vmatrix} 4 \\ + 6\lambda \\ + \nu \end{vmatrix} + b^2d \begin{vmatrix} 4\mu \\ + 2\nu \end{vmatrix} + bc^2 \begin{vmatrix} 3\nu \\ + 6\rho \end{vmatrix},$$

for which to vanish

$$2 + 2\mu = 0, 4 + 6\lambda + \nu = 0, 4\mu + 2\nu = 0, 3\nu + 6\rho = 0,$$

i. e.  $\mu = -1, \nu = 2, \lambda = -1, \rho = -1.$

Thus  $J \equiv ace + 2bcd - ad^2 - b^2e - c^3$

is the only invariant of degree 3 which the quartic possesses.

Ex. 1. Show that a binary  $p$ -ic has no invariant of the second degree if  $p$  is odd, and only one, that of § 48, if  $p$  is even.

Ex. 2. Show that no binary quantic has an invariant of the first degree.

Ex. 3. Show that a binary cubic has one invariant, its discriminant, of the third degree.

Ex. 4. Show that  $a_0^2a_3 - 3a_0a_1a_2 + 2a_1^3$  is annihilated by  $\Omega$ , for any value of  $p$  not less than 3. Is it an invariant for any value of  $p$ ?

Ans. No.

103.] Invariants of several binary quantics. The methods of the earlier part of this chapter apply, *mutatis mutandis*, to a system of binary quantics. The facts developed by them are as follows.

$$\begin{aligned} \text{Let} \quad & (a_0, a_1, a_2, \dots, a_{p_1})(x, y)^{p_1}, \\ & (b_0, b_1, b_2, \dots, b_{p_2})(x, y)^{p_2}, \\ & (c_0, c_1, c_2, \dots, c_{p_3})(x, y)^{p_3}, \\ & \dots \end{aligned}$$

be a system of binary quantics in  $x, y$ . Take the operators

$$\begin{aligned} & \left( a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + \dots + p_1 a_{p_1-1} \frac{d}{da_{p_1}} \right) \\ & + \left( b_0 \frac{d}{db_1} + 2b_1 \frac{d}{db_2} + \dots + p_2 b_{p_2-1} \frac{d}{db_{p_2}} \right) \\ & + \left( c_0 \frac{d}{dc_1} + 2c_1 \frac{d}{dc_2} + \dots + p_3 c_{p_3-1} \frac{d}{dc_{p_3}} \right) + \dots \equiv \Sigma\Omega, \text{ say,} \end{aligned}$$



$$\begin{aligned}
 \text{and } & \left( p_1 a_1 \frac{d}{da_0} + \overline{p_1 - 1} a_2 \frac{d}{da_1} + \dots + a_{p_1} \frac{d}{da_{p_1-1}} \right) \\
 & + \left( p_2 b_1 \frac{d}{db_0} + \overline{p_2 - 1} b_2 \frac{d}{db_1} + \dots + b_{p_2} \frac{d}{db_{p_2-1}} \right) \\
 & + \left( p_3 c_1 \frac{d}{dc_0} + \overline{p_3 - 1} c_2 \frac{d}{dc_1} + \dots + c_{p_3} \frac{d}{dc_{p_3-1}} \right) + \dots \\
 & \equiv \Sigma O, \text{ say.}
 \end{aligned}$$

Then any invariant of the system is annihilated by  $\Sigma\Omega$  and by  $\Sigma O$ . Also, conversely, any rational integral function of the different sets of coefficients which is homogeneous in each set, of partial degrees  $i_1, i_2, i_3, \dots$  say, and of the same total weight  $w$  in the sets jointly throughout, and which has both  $\Sigma\Omega$  and  $\Sigma O$  for annihilators, is an invariant of the system. Moreover, the possession of these two annihilators necessitates that the several partial degrees and the total weight must be connected with one another and the orders of the quantics by the relation

$$i_1 p_1 + i_2 p_2 + i_3 p_3 + \dots = 2w.$$

Also, as in § 95, an invariant is unaltered if its weight be even, or altered only in sign if its weight be odd, when we interchange  $a_0$  and  $a_{p_1}$ ,  $a_1$  and  $a_{p_1-1}, \dots, b_0$  and  $b_{p_2}$ ,  $b_1$  and  $b_{p_2-1}, \dots, c_0$  and  $c_{p_3}$ ,  $c_1$  and  $c_{p_3-1}, \dots, \dots$ . Thus, since these interchanges make  $\Sigma\Omega$  into  $\Sigma O$ , and vice versa, we need not, when we have found a function which  $\Sigma\Omega$  annihilates, test by direct operation whether it is also annihilated by  $\Sigma O$  before being sure whether it is an invariant. If it have the correct symmetry of form as above there is no doubt of the fact.

Ex. 5. Find the invariant of partial degrees 1, 1, and consequently of weight 2, of the two quadratics

$$(a_0, a_1, a_2) (x, y)^2, (b_0, b_1, b_2) (x, y)^2.$$

Ans. Its form must be  $\lambda a_0 b_2 + \mu a_1 b_1 + \nu a_2 b_0$ . Now  $\Sigma\Omega$  on this produces  $(2\lambda + \mu) a_0 b_1 + (\mu + 2\nu) a_1 b_0$ . Thus  $\mu = -2\lambda = -2\nu$ . The one invariant is then that of § 7, Ex. 4,  $\lambda (a_0 b_2 - 2a_1 b_1 + a_2 b_0)$ .

Ex. 6. Show more generally that the invariant of § 49 is the only lineo-linear invariant of two binary  $p$ -ics.

Ex. 7. Show that the quartic and quadratic

$$(a_0, a_1, \dots, a_4) (x, y)^4, (b_0, b_1, b_2) (x, y)^2$$

have no lineo-linear invariant.

*Ans.* The weight would have to be  $\frac{1}{2}(4+2) = 3$ . The only possible form is then  $\lambda a_1 b_2 + \mu a_2 b_1 + \nu a_3 b_0$ ; and  $\Sigma\Omega$  on this produces

$$\lambda a_0 b_2 + (2\lambda + 2\mu) a_1 b_1 + (\mu + 3\nu) a_2 b_0;$$

for which to vanish would require  $\lambda = 0, \mu = 0, \nu = 0$ .

Ex. 8. No two binary quantics of different orders can have a lineo-linear invariant.

Ex. 9. Find the only invariant of partial degrees 2, 1 of a linear form and a quadratic.

*Ans.* The invariant of § 7, Ex. 3.

Ex. 10. Find an invariant of partial degrees 1, 2 of a quadratic and cubic.

*Ans.*  $a_0(b_1 b_3 - b_2^2) - a_1(b_0 b_3 - b_1 b_2) + a_2(b_0 b_2 - b_1^2)$ .

Ex. 11. Find an invariant of partial degrees 1, 1, 1 of three quadratics.

*Ans.* That of § 17, Ex. 25.

104.] **Annihilators of covariants.** In § 69 it has been seen that the covariants of a binary quantic  $u$  are identical with the results of replacing  $x'$  and  $y'$  by  $x$  and  $y$  in the invariants of  $u$  and the linear form  $xy' - x'y$ .

Now invariants of these two quantics have, by the preceding article, the two annihilators

$$\Omega - y' \frac{d}{dx'}, \quad O - x' \frac{d}{dy'}.$$

It follows that covariants of  $u$  have the annihilators

$$\Omega - y \frac{d}{dx}, \quad O - x \frac{d}{dy}.$$

It seems best, however, to prove this fact and develop its consequences *ab initio*, as was done in the matter of invariants.

105.] Let  $F(a_0, a_1, a_2, \dots, a_p; x, y)$  be a covariant of  $(a_0, a_1, a_2, \dots, a_p)(x, y)^p$ ; and let the quantic be transformed into

$$(A_0, A_1, A_2, \dots, A_p)(X, Y)^p$$

by the substitution

$$x = X + mY, \quad y = Y,$$

whose modulus is unity. We seek first the necessary and sufficient condition that we may have

$$F(A_0, A_1, A_2, \dots, A_p; X, Y) = F(a_0, a_1, a_2, \dots, a_p; x, y),$$

which will be the case when  $F$  is a covariant, though not then only.

If  $A_0, A_1, A_2, \dots, A_p$  are expressed in terms of  $m$  and  $a_0, a_1, a_2, \dots, a_p$ , we have, as in § 92,

$$\frac{dA_0}{dm} = 0, \quad \frac{dA_1}{dm} = A_0, \quad \frac{dA_2}{dm} = 2A_1, \dots, \quad \frac{dA_p}{dm} = pA_{p-1}.$$

Also the expressions for  $X$  and  $Y$  in terms of  $m$  and  $x$  and  $y$  are

$$X = x - my, \quad Y = y,$$

so that

$$\frac{dX}{dm} = -y = -Y, \quad \frac{dY}{dm} = 0.$$

Consequently

$$\begin{aligned} & \frac{d}{dm} F(A_0, A_1, A_2, \dots, A_p; X, Y) \\ &= \frac{dF}{dA_0} \cdot \frac{dA_0}{dm} + \frac{dF}{dA_1} \cdot \frac{dA_1}{dm} + \dots + \frac{dF}{dA_p} \cdot \frac{dA_p}{dm} + \frac{dF}{dX} \cdot \frac{dX}{dm} + \frac{dF}{dY} \cdot \frac{dY}{dm} \\ &= A_0 \frac{dF}{dA_1} + 2A_1 \frac{dF}{dA_2} + \dots + pA_{p-1} \frac{dF}{dA_p} - Y \frac{dF}{dX}. \end{aligned}$$

Now for the left-hand member here to vanish is the necessary and sufficient condition that  $F(A_0, A_1, A_2, \dots, A_p; X, Y)$  is a function of  $a_0, a_1, a_2, \dots, a_p; x, y$  which is free from  $m$ , and consequently equal to  $F(a_0, a_1, a_2, \dots, a_p; x, y)$  its value when  $m = 0$ . The vanishing of the right-hand member must then express the same thing. Thus the necessary and sufficient condition required is that

$$\begin{aligned} & (A_0 \frac{d}{dA_1} + 2A_1 \frac{d}{dA_2} + 3A_2 \frac{d}{dA_3} + \dots + pA_{p-1} \frac{d}{dA_p} - Y \frac{d}{dX}) \\ & F(A_0, A_1, A_2, \dots, A_p; X, Y) = 0, \end{aligned}$$

or, replacing capital by small letters, that

$$(\Omega - y \frac{d}{dx}) F(a_0, a_1, a_2, \dots, a_p; x, y) = 0.$$

We have also, as in § 93, that even when this condition is not satisfied

$$F(A_0, A_1, A_2, \dots, A_p; X, Y) = e^{m(\Omega - y \frac{d}{dx})} F(a_0, a_1, a_2, \dots, a_p; x, y).$$

*F(---) any function, etc.*

Thus we have the means of writing down the result of applying the transformation of this article to any function of the coefficients and variables.

We might also have adopted the method of § 91.

106.] In precisely the same way, the necessary and sufficient condition for the persistence in form of  $F$  after transformation by the substitution

$$x = X, \quad y = l'X + Y,$$

is  $(O - x \frac{d}{dy}) F(a_0, a_1, a_2, \dots, a_p; x, y) = 0,$

where  $O$  is the second operator of § 90.

Any covariant has, then, the two annihilators

$$\Omega - y \frac{d}{dx}, \quad O - x \frac{d}{dy}.$$

*2 2 the theorem*  
*| a1 a2 |*

107.] **Symmetry of a covariant.** Again, as in § 95, we see that there is a symmetry in any covariant. The simultaneous interchange of  $x$  and  $y$ , of  $a_0$  and  $a_p$ ,  $a_1$  and  $a_{p-1}$ , &c. in its expression must, since the interchange means a substitution of modulus  $-1$ , have the effect only of multiplying it by  $(-1)^{\frac{1}{2}(ip - \varpi)}$ , where  $i$  is its degree and  $\varpi$  its order. Now  $\frac{1}{2}(ip + \varpi)$  is the weight of the covariant (chap. iii), and  $\frac{1}{2}(ip - \varpi)$  consequently the weight of the coefficient of  $x^\varpi$  in it. Thus a covariant is unaltered, or altered only in sign, by these interchanges, according as the weight of its leading coefficient is even or odd.

Hence for covariants one of the two conditions of § 106 is necessitated by the other and symmetry. The reasoning is as in § 96.

108.] **Sufficiency of the two conditions of annihilation.** We can also prove the converse of § 106 for homogeneous isobaric functions. Stated at length, the fact is that any rational integral function  $F$  of  $a_0, a_1, a_2, \dots, a_p$  and  $x, y$ , which is homogeneous, of degree  $i$ , in the coefficients, homogeneous, of order

$\varpi$ , in  $x, y$ , and isobaric on the whole, reckoning  $a_0, a_1, a_2, \dots a_p$ ,  $x, y$  as of weights 0, 1, 2, ...  $p, 1, 0$ , and which has both  $\Omega - y \frac{d}{dx}$  and  $O - x \frac{d}{dy}$  for annihilators, is a covariant.

The proof, which proceeds exactly as in § 97, need not be repeated at length. The only variation is that in passing from

$$(a_0, a_1, a_2, \dots a_p) (x, y)^p$$

to

$$(a'_0, a'_1, a'_2, \dots a'_p) (x', y')^p,$$

we have, as well as  $a'_r = \lambda^{p-r} \mu^r a_r$ ,

also  $x' = \lambda^{-1} x, y' = \mu^{-1} y$ ,

so that  $x'^s y'^{\varpi-s} = \lambda^{-s} \mu^{s-\varpi} x^s y^{\varpi-s}$ ,

where the index of the power of  $\lambda$  is minus the weight of the product  $x^s y^{\varpi-s}$ , and that of the power of  $\mu$  is the weight of the product lessened by  $\varpi$ . Also the weight of the function of  $a_0, a_1, a_2, \dots a_p$  which multiplies  $x^s y^{\varpi-s}$  in the function  $F$  which we are considering is  $w-s$ , where  $w$  is the weight of  $F$ . Thus, in place of the

$$F(a'_0, a'_1, a'_2, \dots a'_p) = \lambda^{ip-w} \mu^w F(a_0, a_1, a_2, \dots a_p)$$

of § 97, what we now have is, since

$$\lambda^{ip-(w-s)} \mu^{w-s} \cdot \lambda^{-s} \mu^{s-\varpi} = \lambda^{ip-w} \mu^{w-\varpi}$$

is the same for every  $s$ ,

$$F(a'_0, a'_1, a'_2, \dots a'_p; x', y') = \lambda^{ip-w} \mu^{w-\varpi} F(a_0, a_1, a_2, \dots a_p; x, y).$$

This difference of the factor will not affect the argument. The supplementary conclusion, from the fact (§ 37) that when we have proved  $F$  to be a covariant we know that the factor must be a power of the modulus, is in this case

$$ip-w = w-\varpi = \frac{1}{2} (ip-\varpi),$$

i.e. is

$$ip+\varpi = 2w,$$

which accords with chapter iii.

109.] A covariant completely given by an end term. We are now in a position to find the covariants of a given degree and order by a method like that of § 101. A further theorem of great importance will, however, much facilitate the process. It is due to M. Roberts.

Let a covariant of order  $\varpi$  of a binary  $p$ -ic be arranged as a quantic in  $x$  and  $y$ . We may write it

$$C_0 x^\varpi + \varpi C_1 x^{\varpi-1} y + \frac{\varpi(\varpi-1)}{1 \cdot 2} C_2 x^{\varpi-2} y^2 + \dots \\ + \varpi C_{\varpi-1} x y^{\varpi-1} + C_\varpi y^\varpi,$$

where  $C_0, C_1, C_2, \dots, C_\varpi$  are all of degree  $i$  in  $a_0, a_1, a_2, \dots, a_p$ , and of weights respectively

$$\frac{1}{2}(ip - \varpi), \frac{1}{2}(ip - \varpi) + 1, \frac{1}{2}(ip - \varpi) + 2, \dots, \frac{1}{2}(ip + \varpi).$$

This is annihilated by  $\Omega - y \frac{d}{dx}$ . We must have then

$$\Omega C_0 \cdot x^\varpi + \varpi(\Omega C_1 - C_0) x^{\varpi-1} y + \frac{\varpi(\varpi-1)}{1 \cdot 2} (\Omega C_2 - 2C_1) x^{\varpi-2} y^2 + \dots \\ + \varpi(\Omega C_{\varpi-1} - \overline{\varpi-1} C_{\varpi-2}) x y^{\varpi-1} + (\Omega C_\varpi - \varpi C_{\varpi-1}) y^\varpi = 0,$$

for all values of  $x$  and  $y$ . The various coefficients of  $x^\varpi, x^{\varpi-1} y, \dots$  must therefore vanish separately. In other words,

$$\Omega C_0 = 0,$$

$$\Omega C_1 = C_0,$$

$$\Omega C_2 = 2C_1,$$

$$\dots$$

$$\Omega C_{\varpi-1} = (\varpi-1) C_{\varpi-2},$$

$$\Omega C_\varpi = \varpi C_{\varpi-1}.$$

We have then the two most interesting conclusions which follow:

(1)  $C_0$ , the leading coefficient in the covariant, is annihilated by  $\Omega$ , the first of the two annihilators of invariants. For this reason it is called a semi-invariant or *seminvariant*.

(2) When  $C_\varpi$ , the last coefficient in a covariant, is known, all the other coefficients are determined from it by mere operations with  $\Omega$ , i.e. by differentiations only. In fact, we see that

$$C_r = \frac{1}{\varpi(\varpi-1)(\varpi-2)\dots\varpi(r+1)} \Omega^{\varpi-r} C_\varpi;$$

and that the whole covariant is

$$\frac{\Omega^\varpi C_\varpi}{\varpi!} x^\varpi + \frac{\Omega^{\varpi-1} C_\varpi}{(\varpi-1)!} x^{\varpi-1} y + \frac{\Omega^{\varpi-2} C_\varpi}{(\varpi-2)!} x^{\varpi-2} y^2 + \dots \\ + \Omega C_\varpi x y^{\varpi-1} + C_\varpi y^\varpi,$$

which, since  $\Omega^{\varpi+1}C_{\varpi}$ ,  $\Omega^{\varpi+2}C_{\varpi}$ , &c., vanish, the first of them being only  $\varpi! \Omega C_0$  which is zero by the first equality above, may briefly be written

$$y^{\varpi} e^{\frac{x}{y} \Omega} C_{\varpi}.$$

For this reason  $C_{\varpi}$ , which is (§ 107) merely the result of interchanging  $a_0$  and  $a_p$ ,  $a_1$  and  $a_{p-1}$ , &c. in  $(-1)^{\frac{1}{2}(i p - \varpi)} C_0$ , has been called the *source* of the covariant. In the next article we shall see that the same name might equally and for a like reason be given to  $C_0$  itself.

110.] Express, in fact, precisely as in the preceding article, that the covariant is annihilated by  $O - x \frac{d}{dy}$ . The conclusions are that

$$\begin{aligned} OC_0 &= \varpi C_1 \\ OC_1 &= (\varpi - 1) C_2 \\ OC_2 &= (\varpi - 2) C_3 \\ &\dots \dots \dots \\ OC_{\varpi-1} &= C_{\varpi}, \\ OC_{\varpi} &= 0. \end{aligned}$$

Thus (1)  $C_{\varpi}$  is annihilated by  $O$  the second annihilator of invariants, and may be called an *anti-seminvariant*; and (2) for every value of the number  $r$  from 0 to  $\varpi$  inclusive

$$C_r = \frac{1}{\varpi(\varpi-1)(\varpi-2)\dots(\varpi-r+1)} O^r C_0,$$

so that the covariant is

$$C_0 x^{\varpi} + OC_0 x^{\varpi-1} y + \frac{O^2 C_0}{1 \cdot 2} x^{\varpi-2} y^2 + \dots + \frac{O^{\varpi} C_0}{\varpi!} y^{\varpi},$$

or, as it may be written,

$$x^{\varpi} e^{\frac{y}{x} O} C_0,$$

for  $O^{\varpi+1}C_0$ ,  $O^{\varpi+2}C_0$ , ... vanish since  $OC_{\varpi} = 0$ .

Thus, when we have the seminvariant  $C_0$  which is the leading coefficient of a covariant, all the coefficients in the covariant can be obtained from it by mere operations with  $O$ , i.e. by differentiations.

In fact, given any coefficient in a covariant, all the coefficients can be found. Successive operations with  $\Omega$  give the

coefficients on the one side of it, and successive operations with  $O$  give those on the other.

111.] **Seminvariants.** We may define a *seminvariant* as any homogeneous isobaric function of the coefficients  $a_0, a_1, a_2, \dots a_p$  which is annihilated by  $\Omega$ . We now confine attention, however, to seminvariants which are rational and integral.

Looking back at §§ 91, 92 we see that the half invariant property which seminvariants possess is that of being absolutely invariantic for such linear substitutions as

$$x = X + mY, \quad y = Y.$$

(From § 97 we gather that they are really invariantic for the somewhat more general substitution

$$x = lX + mY, \quad y = m'Y,$$

though the factor for any one is in this case not as a rule a power of the modulus.)

Consequently, when expressed in terms of  $a_0$  and the roots of the quantic, a seminvariant can involve only differences of these latter. If of degree  $i$ , and not divisible by  $a_0$ , it is a product of  $a_0^i$  and a function of the differences, which involves each particular root to the  $i$ -th degree, since the ratios of  $a_1, a_2, a_3, \&c.$ , to  $a_0$  are all of the first degree in every root. Conversely, any rational integral symmetric function of the roots, which can be expressed in terms of their differences only, becomes a rational integral seminvariant when multiplied by such a power of  $a_0$  that it can be expressed integrally in terms of the coefficients. The least power of  $a_0$  which suffices is the  $i$ -th, where  $i$  is the degree of the symmetric function in any particular root.

We may now see that in the two preceding articles  $C_0$  may be any rational integral seminvariant; that is to say, that any rational integral seminvariant whatever may be taken as the leading coefficient of a covariant, and determines that covariant uniquely.

Take any rational integral seminvariant  $S$  of degree  $i$ , without  $a_0$  for a factor, and write it as  $a_0^i$  multiplied into a symmetric function of the roots  $a_1, a_2, \dots a_p$ . This symmetric function



will be a function of the differences  $a_r - a_s$ , &c. Now for every difference  $a_r - a_s$  write  $\frac{a_r - a_s}{(x - a_r y)(x - a_s y)}$ . In this fraction  $a_r$  occurs once in the numerator and once in the denominator, and so does  $a_s$ .

Clear the function obtained of fractions by multiplying by the lowest necessary power, the  $i$ -th, of

$$(x - a_1 y)(x - a_2 y) \dots (x - a_p y).$$

In this multiplier  $a_1, a_2, \dots a_p$  occur in equal numbers  $i$  of factors.

The result, when expressed in terms of  $x, y$  and coefficients only, will be a covariant whose leading coefficient  $C_0$  is the seminvariant  $S$ . That the leading coefficient is  $S$  is clear from the method of construction. That the whole expression is a covariant follows from the fact (§ 83) that it is a power of  $a_0$  multiplied into a function of differences  $x - ay$  and differences  $a_r - a_s$ , which is symmetrical in the roots, homogeneous in both kinds of differences, and such that all roots  $a$  occur in equal numbers of factors in any product, and in the same number in all products.

The covariant is unique of its degree and order; for § 110 shows that a leading coefficient  $C_0$  determines a covariant uniquely, giving  $\varpi$  as the least number for which  $O^{\varpi+1}C_0 = 0$ , and giving  $C_1, C_2, \dots C_\varpi$  by a succession of operations with  $O$  on  $C_0$ . The uniqueness also follows from § 39. Note, however, that the order  $\varpi$  depends on the value of  $p$ . For quantities of different orders  $p$ , the same <sup>covariant</sup> covariant will lead covariants of different orders  $\varpi$ . We had above, in forming the covariant from its seminvariant leader by means of the roots, to divide terms by products of order  $2w'$ , where  $w'$  is the weight of the seminvariant, and to multiply through by a product of order  $ip$ . Altogether the covariant obtained is of order  $ip - 2w'$ , which accords with the known fact that  $w' = \frac{1}{2}(ip - \varpi)$ .

A seminvariant with a power of  $a_0, a_0^j$  say, for a factor is of the form  $a_0^j S$ , where  $S$  is a seminvariant to which the reasoning above applies. The unique covariant which it leads is the product of the covariant led by  $S$  and the  $j$ -th power of the  $p$ -ic.

Another proof, making no explicit use of the roots, of the

important theorem of the present article will be given in the next chapter. (Cf. § 126, Ex. 6.)

112.] We now see that the problem of finding covariants by aid of annihilators is reduced to the much simpler one of finding seminvariants. We have found, suppose, a seminvariant of degree  $i$  and weight  $w$ . Take it for  $C_0$  in § 110. The leading term in the corresponding covariant of the binary  $p$ -ic is  $C_0 x^{ip-2w}$  as above, when  $a_0$  is a factor of  $C_0$  as well as in other cases, where  $w$  is the weight of  $C_0$  and not that of the covariant; and the full expression of the covariant is, by § 110,

$$x^{ip-2w} e^{\frac{y}{x}} O C_0.$$

That the order  $ip-2w$  of the covariant cannot be negative,  $C_0$  being a rational integral seminvariant, is clear. For the procedure of the last article determined a covariant of essentially non-negative order in  $x$  and  $y$  from the seminvariant with which it started. Thus there are no rational integral seminvariants for which  $ip-2w$  is negative. Should  $ip-2w$  be zero the seminvariant is an invariant. For the covariant derived from it is of zero order as above and consequently an invariant, and is in fact the seminvariant itself. It must clearly be borne in mind, however, that we are dealing only with rational integral seminvariants. The argument does not apply, for instance, to the fractional seminvariant  $\frac{1}{a}(a^2d-3abc+2b^3)$ . Here, for the cubic,  $ip-2w=0$ , but nevertheless the function is not an invariant of the cubic.  $O$  does not annihilate it.

We may, in fact, state succinctly the conclusions arrived at as to invariants of a binary  $p$ -ic. If  $\Omega$  and  $O$  annihilate a function, it is an invariant, and  $ip-2w=0$  (§ 97). If  $\Omega$  annihilate a function for which  $ip-2w=0$ , it is also annihilated by  $O$  and is consequently an invariant *provided it is rational and integral*, but not necessarily if it is fractional.

113.] **Determination of seminvariants.** The determination of the linearly independent seminvariants of degree  $i$  and weight  $w$  of a binary  $p$ -ic proceeds as in § 101. If  $ip < 2w$  there are none, as above. If  $ip \leq 2w$ , write down all the products of weight  $w$  of  $i$  constituents chosen from among

$a_0, a_1, a_2, \dots, a_p$ , repetitions of factors allowed, and add together arbitrary multiples of these. The number of such products is  $(w; i, p)$ , this symbol denoting, as in § 101, the number of different partitions of the number  $w$  into  $i$  or fewer numbers, none exceeding  $p$ . Operate on this sum with  $\Omega$ , thus obtaining a sum of multiples of the  $(w-1; i, p)$  products of degree  $i$  and weight  $w-1$ . It will be proved in the next chapter that the  $(w-1; i, p)$  coefficients of these products are linearly independent linear functions of the  $(w; i, p)$  arbitrary multipliers of the  $(w; i, p)$  products. They have to vanish. Their vanishing gives  $(w-1; i, p)$  relations which have to be satisfied by the  $(w; i, p)$  multipliers. If then  $(w; i, p)$  exceed  $(w-1; i, p)$  we can satisfy them and leave

$$(w; i, p) - (w-1; i, p)$$

of the multipliers still arbitrary. Suppose that this excess is  $r$ . We have as the most general seminvariant of the type under consideration a sum of the form

$$\lambda_1 S_1 + \lambda_2 S_2 + \dots + \lambda_r S_r$$

where  $\lambda_1, \lambda_2, \dots, \lambda_r$  are arbitrary, and  $S_1, S_2, \dots, S_r$  are known linear functions of the products of type  $w, i$ . We express this by saying that there are  $r$  linearly independent seminvariants

$$S_1, S_2, \dots, S_r$$

of this type belonging to the  $p$ -ic.

114.] **Seminvariants of the second degree.** As an example let us discover all the seminvariants of degree 2 of a binary  $p$ -ic. The condition  $ip \leq 2w$  is for this case  $w \geq p$ . Two cases will arise.

(1) For an even weight  $w$ , not exceeding  $p$ , the general form to be assumed is

$$a_0 a_w + \lambda_1 a_1 a_{w-1} + \lambda_2 a_2 a_{w-2} + \dots + \lambda_{\frac{1}{2}w} a_{\frac{1}{2}w}^2.$$

Expressing that  $\Omega$  annihilates this, we obtain the conditions

$$\begin{aligned} w + \lambda_1 &= 0, & (w-1)\lambda_1 + 2\lambda_2 &= 0, \\ (w-2)\lambda_2 + 3\lambda_3 &= 0, \dots & \left(\frac{w}{2} + 1\right)\lambda_{\frac{1}{2}w-1} + w\lambda_{\frac{1}{2}w} &= 0, \end{aligned}$$

the coefficient  $w$  in the last of these being double what it

would be according to the law of all the other second terms. Solving these for the  $\lambda$ 's we have a unique seminvariant of weight  $w$ , viz.

$$a_0 a_w - w a_1 a_{w-1} + \frac{w(w-1)}{1 \cdot 2} a_2 a_{w-2} - \dots \\ + (-1)^{\frac{1}{2}w} \frac{w(w-1) \dots (\frac{1}{2}w+1)}{2(\frac{1}{2}w)!} a_{\frac{1}{2}w}^2,$$

where the law of coefficients is that of the expansion of  $(1+z)^w$  up to its middle coefficient, which one alone is halved.

For every even weight not exceeding  $p$  there is then a single seminvariant of degree 2. In particular, of course, it is an invariant for a weight equal to  $p$  if  $p$  is even.

(2) For an odd weight  $w$ , not exceeding  $p$ , the form to be assumed is

$$a_0 a_w + \lambda_1 a_1 a_{w-1} + \lambda_2 a_2 a_{w-2} + \dots + \lambda_{\frac{1}{2}(w-1)} a_{\frac{1}{2}(w-1)} a_{\frac{1}{2}(w+1)},$$

and the conditions obtained from the annihilation by  $\Omega$  are

$$w + \lambda_1 = 0, \quad (w-1)\lambda_1 + 2\lambda_2 = 0, \quad (w-2)\lambda_2 + 3\lambda_3 = 0, \dots \\ \frac{1}{2}(w+3)\lambda_{\frac{1}{2}(w-3)} + \frac{1}{2}(w-1)\lambda_{\frac{1}{2}(w-1)} = 0, \quad \lambda_{\frac{1}{2}(w-1)} = 0,$$

of which the last tells us that  $\lambda_{\frac{1}{2}(w-1)}$  vanishes, and the rest, taken in order backwards, tells us that all the other  $\lambda$ 's vanish.

For no odd weight, then, is there a seminvariant of degree 2.

Accordingly the complete series of seminvariants of degree 2 is

$$a_0^2, \\ a_0 a_2 - a_1^2, \\ a_0 a_4 - 4 a_1 a_3 + 3 a_2^2, \\ a_0 a_6 - 6 a_1 a_5 + 15 a_2 a_4 - 10 a_3^2, \\ a_0 a_8 - 8 a_1 a_7 + 28 a_2 a_6 - 56 a_3 a_5 + 35 a_4^2, \\ \&c., \&c.,$$

the series terminating with the weight  $p$  or  $p-1$  according as  $p$  is even or odd. In the former case the last of the series is an invariant.

The orders of the corresponding covariants are  $2p, 2p-4, 2p-8, \dots$ . They are the covariants of § 57, Ex. 12, together

with the invariant, if  $p$  be even, of § 48. In other words, they are the transvectants (§ 61) of the  $p$ -ic and itself. Thus we have the theorem that a binary quantic  $u$  has no covariants of the second degree in the coefficients besides the 0th, second, fourth, sixth, &c., transvectants of  $u$  and itself. The 0th, led by  $a_0^2 x^{2p}$ , is  $u^2$ .

Ex. 12. Use § 110 to write down in full the covariants of degree 2.

$$a_0^2 x^{10} + \dots, (a_0 a_2 - a_1^2) x^8 + \dots, (a_0 a_4 - 4 a_1 a_3 + 3 a_2^2) x^2 + \dots$$

of the binary quintic.

*Ans.* For the last see § 61, Ex. 29.

Ex. 13. A binary quantic of order not less than 4 has one and only one seminvariant of degree 3 and weight 6.

$$\text{Ans.} \quad a_0 a_2 a_4 + 2 a_1 a_2 a_3 - a_0 a_3^2 - a_1^2 a_4 - a_2^3.$$

Ex. 14. Hence obtain the single covariant of degree 3 and order 3 of the quintic.

$$\begin{aligned} \text{Ans.} \quad & (ace + 2bcd - ad^2 - b^2e - c^3)x^3 \\ & + (acf - ade - b^2f + bce + bd^2 - c^2d)x^2y \\ & + (adf - bcf - ae^2 + bde + c^2e - cd^2)xy^2 \\ & + (bdf + 2cde - c^2f - be^2 - d^3)y^3. \end{aligned}$$

Ex. 15. Show that a binary quantic of order not less than 3 has one and only one seminvariant of degree 3 and weight 3.

$$\text{Ans.} \quad a_0^2 a_3 - 3 a_0 a_1 a_2 + 2 a_1^3.$$

Ex. 16. Hence, and from what has been proved above, prove the statement of § 86 that  $H$  and  $G$  are the only covariants of a binary cubic whose degrees are 2, 3 and orders 2, 3 respectively.

Ex. 17. Show that a binary quartic, or binary quantic of higher order, has two and only two seminvariants of degree 4 and weight 4.

$$\text{Ans.} \quad \lambda a^2 (ae - 4bd + 3c^2) + \mu (ac - b^2)^2,$$

which may also be written

$$\lambda (a^3e - 4a^2bd + 6ab^2c - 3b^4) + \mu' (ac - b^2)^2.$$

Ex. 18. Find the sum  $\sigma_4$  of the six fourth powers of differences between the roots of the quartic  $(a, b, c, d, e) (x, 1)^4$ .

$$\text{Ans.} \quad a^4 \sigma_4 = 720 (ac - b^2)^2 - 16 (a^3e - 4a^2bd + 6ab^2c - 3b^4).$$

Determine  $\lambda$  and  $\mu'$  in the second form above by taking two particular quartics, e.g.  $x^2(x^2 - 1)$  and  $x^4 - 1$ , in both of which  $b = 0$ , and in one of which  $e = 0$  and in the other  $c = 0$ .

Ex. 19. Show that a binary quintic has two and only two independent seminvariants of degree 5 and weight 5.

$$\text{Ans. } \lambda (a^4f - 5a^3be + 10a^2b^2d - 10ab^3c + 4b^5) + \mu (ac - b^2) (a^2d - 3abc + 2b^3).$$

Ex. 20. The sum of the numerical coefficients in any seminvariant which is not a mere power of  $a_0$  vanishes.

Ex. 21. If  $(C_0, C_1, C_2, \dots, C_m) (x, y)^m$  is a covariant of

$$(a_0, a_1, a_2, \dots, a_p) (x, y)^p$$

prove from § 109 that  $a_0C_1 - a_1C_0$  is a seminvariant. (*M. Roberts.*)

Ex. 22. By application of this result to the first linear covariant (§ 57, Ex. 17) of a binary quintic, prove that the quintic has a covariant of degree 6 and order 4, and an invariant of degree 18, the catalecticant of this covariant.

115.] **Seminvariants and covariants of several binary quantics.** Referring to § 103 for the notation, we define a seminvariant of a system of quantics in the same variables  $x, y$  as a function of the several sets of coefficients—in general a rational integral function—which is homogeneous in each set separately, and isobaric on the whole, though not necessarily in the sets separately, and which has  $\Sigma\Omega$  for an annihilator.

The methods which have preceded are applicable to covariants and seminvariants of systems of quantics. It is left to the student to convince himself, as in § 104 or §§ 105, 106, that every covariant of the system is annihilated by  $\Sigma\Omega - y \frac{d}{dx}$  and  $\Sigma O - x \frac{d}{dy}$ ; and, as in § 108, that conversely a function which is homogeneous in the variables and in every set of coefficients separately, and isobaric on the whole, is a covariant if these operators annihilate it. He will also see, as in § 107, that if  $x$  and  $y$ ,  $a_0$  and  $a_{p_1}$ ,  $a_1$  and  $a_{p_1-1}$ , ...,  $b_0$  and  $b_{p_2}$ ,  $b_1$  and  $b_{p_2-1}$ , ...,  $c_0$  and  $c_{p_3}$ ,  $c_1$  and  $c_{p_3-1}$ , ..., ... are interchanged in a covariant the effect is only to multiply the covariant by  $(-1)^{\frac{1}{2}(\Sigma \cdot ip - \omega)}$ , where the index is the total weight of the leading coefficient, that of  $x^m$ . That this leading coefficient  $C_0$  is a seminvariant, i. e. is annihilated by  $\Sigma\Omega$ , he will see as in § 109, and that all other coefficients can be derived from it

by operations with  $\Sigma O$  he will see as in § 110. In fact, the covariant may be written either as

$$x^{\varpi} e^{x^{\frac{y}{x} \Sigma O}} C_0,$$

or as

$$y^{\varpi} e^{y^{\frac{x}{y} \Sigma \Omega}} C_{\varpi}.$$

Once more, as in § 111, he will see that any rational integral seminvariant whatever may be taken as the leading coefficient  $C_0$ , and determines  $\varpi$ , the order, and the full expression for the covariant uniquely. The order  $\varpi$ , the partial degrees  $i_1, i_2, i_3, \dots$  in the coefficients of the  $p_1$ -ic, the  $p_2$ -ic, the  $p_3$ -ic, &c., and the weight  $w$  of the seminvariant are seen to be connected by the relation

$$i_1 p_1 + i_2 p_2 + i_3 p_3 + \dots - \varpi = 2w.$$

Thus there is no seminvariant for which  $\Sigma . ip - 2w$  is negative. If  $\Sigma . ip - 2w = 0$  the derived covariant and the seminvariant are identical. The seminvariant is in fact an invariant, and is annihilated by  $\Sigma O$ .

The method of §§ 98–100 also applies; and the results of § 100 hold, when we put  $\Sigma \Omega, \Sigma O, \Sigma . ip - 2w$  in place of  $\Omega, O, ip - 2w$  respectively, for operations on functions of any or all of the sets of coefficients.

All the linearly independent seminvariants of given weight  $w$  and partial degrees  $i_1, i_2, i_3, \dots$  are found, as in § 113, by writing down the most general rational integral function of the type in question and determining the multipliers in it so that it may be annihilated by  $\Sigma \Omega$ .

Ex. 23. Find a seminvariant of weight 2 and partial degrees, 1, 1 of the quadratic and cubic  $(a_0, a_1, a_2) (x, y)^2, (b_0, b_1, b_2, b_3) (x, y)^3$ , and show that the covariant to which it leads is linear.

Ans. The covariant is

$$(a_0 b_2 - 2a_1 b_1 + a_2 b_0) x + (a_0 b_3 - 2a_1 b_2 + a_2 b_1) y.$$

Ex. 24. Find a linear covariant of partial degrees 2, 1 of the quadratic and cubic.

$$\text{Ans. } (a_0^2 b_3 - 3a_0 a_1 b_2 + a_0 a_2 b_1 + 2a_1^2 b_1 - a_1 a_2 b_0) x \\ - (a_2^2 b_0 - 3a_1 a_2 b_1 + a_0 a_2 b_2 + 2a_1^2 b_2 - a_0 a_1 b_3) y.$$

Ex. 25. Remembering that a cubic has a cubicovariant (§ 45, Ex. 13) deduce two other linear covariants of a quadratic and cubic.

Ex. 26. Two different binary quantics of orders  $p$  and  $p'$  have a single lineo-linear seminvariant of every weight not exceeding the smaller of  $p, p'$ , and none of higher weight than this.

$$\text{Ans. } a_0 b_w - w a_1 b_{w-1} + \frac{w(w-1)}{1 \cdot 2} a_2 b_{w-2} + \dots + (-1)^w a_w b_0.$$

Ex. 27. The covariants led by these seminvariants are the mutual transvectants (§§ 59, 61) of the two quantics. Hence the mutual transvectants of  $u$  and  $v$  are the only covariants of  $u$  and  $v$  which are lineo-linear in the coefficients.

### ADDITIONAL EXAMPLES.

Ex. 28. Prove that  $I = 0, J = 0$ , where  $I$  and  $J$  are the invariants of the quartic  $u \equiv (a, b, c, d, e) (x, y)^4$  are two results which can be obtained by elimination of  $x$  between

$$\frac{d^2 u}{dx^2} = 0, \quad \frac{d^2 u}{dx dy} = 0, \quad \frac{d^2 u}{dy^2} = 0. \quad (\text{Cayley.})$$

Ex. 29. If  $a, \beta, \gamma, \delta$  the roots of  $(a, b, c, d, e) (x, y)^4$  be taken in pairs  $a, \beta; \gamma, \delta$  in any way, the substitution

$$\begin{aligned} x &= (\gamma\delta - a\beta)X - \{a\beta(\gamma + \delta) - \gamma\delta(a + \beta)\} Y, \\ y &= (\gamma + \delta - a - \beta)X - (\gamma\delta - a\beta) Y, \end{aligned}$$

transforms the quartic into the same quartic  $(a, b, c, d, e) (X, Y)^4$  multiplied by a function of the roots.

Ans. Consider the quartic in its factorized form.

Ex. 30. If in any covariant of  $(a_0, a_1, a_2, \dots, a_p) (x, y)^p$  we put

$$a_0 x + a_1 y, a_1 x + a_2 y, a_2 x + a_3 y, \dots, a_{p-1} x + a_p y$$

for  $a_0, a_1, a_2, \dots, a_p$  respectively, we deduce the covariant with the same leading coefficient of

$$(a_0, a_1, a_2, \dots, a_p, a_{p+1}) (x, y)^{p+1}. \quad (\text{Cayley.})$$

Ans. Symmetrical and annihilated by  $\Omega_{p+1} - y \frac{d}{dx}$ .

Ex. 31. To substitute  $(a_0, a_1, a_2) (x, y)^2, (a_1, a_2, a_3) (x, y)^2, \dots$  for  $a_0, a_1, a_2, \dots$  is to repeat the same process twice, and to deduce a covariant of  $(a_0, a_1, a_2, \dots, a_p, a_{p+1}, a_{p+2}) (x, y)^{p+2}$ . (Cayley.)

Ex. 32. In any seminvariant or invariant of  $(a_0, a_1, a_2, \dots, a_p) (x, y)^p$  put  $k, k-1, k-2, \dots, k-p$  for  $a_0, a_1, a_2, \dots, a_p$ , and equate to the result of putting  $k, -1$  for  $a_0, a_1$  in the one term, if there be any,



which involves  $a_0, a_1$  only, or to zero if no such term occurs (i. e. if  $w > i$ ). The result is an identity for all values of  $k$ .

*Ans.* Follows from the equality of seminvariants of

$$k(x+y)^p - p(x+y)^{p-1}y \quad \text{and} \quad kX^p - pX^{p-1}Y.$$

Ex. 33. Hence by giving  $k$  the values 0, 1, 2, ... in succession obtain facts with regard to the numerical coefficients of terms free from  $a_0$ , terms free from  $a_1$ , terms free from  $a_2$ , ... in any seminvariant or invariant.

Ex. 34. In this way determine the seminvariants of degree 3 and weight 3, and of degree 2 and weight 4.

Ex. 35. In the same way determine the terms free from  $b$  in the discriminant of the cubic  $(a, b, c, d)(x, y)^3$ , and deduce the full expression for the discriminant, by considering the transformation of the cubic to a form without a second term.

Ex. 36. By consideration of the special binary quantic

$$k(x+y)^p - \frac{p(p-1)}{1 \cdot 2}(x+y)^{p-2}y^2,$$

prove that if in any seminvariant or invariant  $a_0, a_1, a_2, a_3, a_4, \dots$  are replaced by  $k, k, k-1, k-3, k-6, \dots$ , where 1, 3, 6, 10, 15, ... are the figurate numbers of the third order, the result is equal to that of replacing  $a_0$  and  $a_2$  by  $k$  and  $-1$  in the one term which involves  $a_0$  and  $a_2$  only, or to zero if there be no such term.

Ex. 37. Generally, if in any seminvariant or invariant of a binary  $p$ -ic we replace  $a_0, a_1, \dots, a_{r-1}$  all by  $k$ , and  $a_r, a_{r+1}, \dots, a_p$  by  $k$  diminished respectively by the first, second, ...  $(p-r+1)$ th figurate numbers of the  $(r+1)$ th order, the result is equal to that of replacing  $a_0$  and  $a_r$  by  $k$  and  $-1$  in the one term which involves  $a_0$  and  $a_r$  only, or to zero if there be no such term.

Ex. 38. If  $F(a_0, a_1, a_2, \dots, a_p)$  is a rational integral homogeneous isobaric function, of the coefficients in  $(a_0, a_1, a_2, \dots, a_p)(x, y)^p$ , for which  $ip - 2w = 0$ , prove that

$$\left(2 - \frac{\Omega O}{1^2} + \frac{\Omega^2 O^2}{1^2 \cdot 2^2} - \frac{\Omega^3 O^3}{1^2 \cdot 2^2 \cdot 3^2} + \dots\right) \{F(a_0, a_1, a_2, \dots, a_p) - (-1)^w F(a_p, a_{p-1}, a_{p-2}, \dots, a_0)\} = 0.$$

*Ans.* Use § 100.

Ex. 39. In the same case prove that

$$F(a_0, a_1, a_2, \dots, a_p) - (-1)^w F(a_p, a_{p-1}, a_{p-2}, \dots, a_0)$$

is of the form  $\Omega G(a_0, a_1, a_2, \dots, a_p)$ , where  $G$  is a rational integral homogeneous isobaric function.

Ex. 40. Prove that the method of § 19 applies to seminvariants, irrespective of the orders of quantics to which they belong, so that

from covariants with like seminvariant leaders of two quantics of different orders intermediate covariants follow.

Ex. 41. If  $J$  is the Jacobian  $(ab' - a'b)x^{p+p'-2} + \dots$  of two binary quantics  $u, u'$ , and if  $H$  and  $H'$  are their Hessians

$$(ac - b^2)x^{2p-4} + \dots, (a'c' - b'^2)x^{2p'-4} + \dots,$$

and  $H''$  the intermediate covariant

$$(ac' + a'c - 2bb')x^{p+p'-4} + \dots$$

between  $H$  and  $H'$ , prove that

$$J^2 = -u^2H' + uu'H'' - u'^2H. \quad (\text{Faa de Bruno.})$$

*Ans.* It suffices to prove the relation among the seminvariant leaders.

Ex. 42. Any factor of a seminvariant is a seminvariant. (*Sylvester.*)

*Ans.* If  $\Omega P^n = 0$  then  $P^{n-1}\Omega P = 0$ , i. e.  $\Omega P = 0$ . If  $\Omega \cdot PQ = 0$ , then  $\frac{\Omega P}{\Omega Q} = -\frac{P}{Q}$ , whence, if  $Q$  and  $P$  have no common factor,  $\Omega P = 0, \Omega Q = 0$ .

Ex. 43. If  $A = PB + QC$ , where  $A, B, C$  are seminvariants but  $P$  and  $Q$  are not, then there is a relation  $A = P'B + Q'C$  in which  $P'$  and  $Q'$  are seminvariants. (*Sylvester.*)

*Ans.*  $B\Omega P + C\Omega Q = 0$ .  $\therefore \Omega P = -CK, \Omega Q = BK$ . Therefore, if  $B$  and  $C$  have no common factor, (if they have, use first Ex. 42.)

$$P = -C\Omega^{-1}K + P', \quad Q = B\Omega^{-1}K + Q'. \quad (\text{Cf. } \S 100.)$$

Ex. 44. If  $\phi(a, b, c, d, e, f, g, \dots)$  be a seminvariant, then

$$\phi(0, a, 2b, 3c, 4d, 5e, 6f, \dots), \quad \phi(0, 0, a, 3b, 6c, 10d, 15e, \dots), \\ \phi(0, 0, 0, a, 4b, 10c, 20d, \dots), \dots$$

are other seminvariants, the series of numbers being figurate.

## CHAPTER VII.

FURTHER THEORY OF THE OPERATORS  $\Omega$  AND  $O$ . RECIPROCITY.

116.] FOR brevity of statement we henceforth use, with Sylvester, a single word to denote a function of the  $p$  quantities  $a_0, a_1, a_2, \dots a_p$  which is rational and integral and of the same degree and the same weight throughout. The name adopted for such a function is a *gradient* in  $a_0, a_1, a_2, \dots a_p$ .

So, too, by a *gradient* in more sets of quantities  $a_0, a_1, \dots a_{p_1}$ ;  $b_0, b_1, \dots b_{p_2}$ ;  $c_0, c_1, \dots c_{p_3}$ ; ... than one we mean a rational integral function of some or all of the quantities which is of constant degrees throughout in the sets of quantities separately, and of constant weight throughout in the sets collectively.

For the present we deal with gradients in one set  $a_0, a_1, a_2, \dots a_p$  only. It is in accordance with what has preceded to denote in general the *degree* of a gradient by  $i$  and its *weight* by  $w$ .

We need not always have in view that  $a_0, a_1, a_2, \dots a_p$  are the coefficients in a binary  $p$ -ic, but may specify that a gradient involves  $a_p$ , but no element with a greater suffix than  $p$ , by describing it as of *extent*  $p$ .

For instance,

$$a_0 a_5 + \lambda a_1 a_4 + \mu a_2 a_3,$$

and 
$$a_0 a_2 a_4 + \lambda a_1 a_2 a_3 + \mu a_0 a_3^2 + \nu a_1^2 a_4 + \varpi a_2^3,$$

where the coefficients  $\lambda, \mu, \nu, \varpi$  are arbitrary but independent of  $a_0, a_1, a_2, a_3, a_4, a_5$ , are gradients of degrees 2, 3, weights 5, 6, and extents 5, 4 respectively.

117.] **Expressions of homogeneity and isobarism.** Any gradient whatever, of given degree and weight, satisfies two linear differential equations.

Take  $G_{w, i, p}$  a gradient of weight  $w$ , degree  $i$ , and extent  $p$ . One of the two differential equations is Euler's equation which expresses its homogeneity, of degree  $i$ , viz.

$$\left( a_0 \frac{d}{da_0} + a_1 \frac{d}{da_1} + a_2 \frac{d}{da_2} + \dots + a_p \frac{d}{da_p} \right) G_{w, i, p} = i G_{w, i, p}.$$

The other expresses that it is isobaric, of constant weight  $w$  throughout. It is

$$\left( a_1 \frac{d}{da_1} + 2a_2 \frac{d}{da_2} + \dots + pa_p \frac{d}{da_p} \right) G_{w, i, p} = w G_{w, i, p}.$$

This also follows from Euler's theorem of homogeneous functions, for constancy of weight has been seen to be the same thing as homogeneity in magnitudes which are roots of  $(a_0, a_1, a_2, \dots, a_p)(x, y)^p$ , in which  $a_0, a_1, a_2, \dots, a_p$  are homogeneous and of dimensions 0, 1, 2, ...  $p$  respectively. It is, however, at once clear when we notice that, if  $a_r^\rho a_s^\sigma a_t^\tau \dots$  be any term in  $G_{w, i, p}$ ,

$$\begin{aligned} \left( a_1 \frac{d}{da_1} + 2a_2 \frac{d}{da_2} + \dots + pa_p \frac{d}{da_p} \right) \cdot a_r^\rho a_s^\sigma a_t^\tau \dots \\ = (\rho r + \sigma s + \tau t + \dots) a_r^\rho a_s^\sigma a_t^\tau \dots \\ = w a_r^\rho a_s^\sigma a_t^\tau \dots \end{aligned}$$

118.] **Seminvariants as particular Gradients.** Referring to § 111 we see that, according to the definition there given, those gradients  $G_{w, i, p}$ , of type  $w, i, p$ , whose arbitrary coefficients are so chosen that they satisfy the third linear differential equation

$$\Omega G_{w, i, p} = 0,$$

where  $\Omega \equiv a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + 3a_2 \frac{d}{da_3} + \dots + pa_{p-1} \frac{d}{da_p}$ , are the seminvariants of the  $p$ -ic

$$(a_0, a_1, a_2, \dots, a_p)(x, y)^p,$$

and are equally seminvariants, though not all the seminvariants, of the  $(p+q)$ -ic

$$(a_0, a_1, a_2, \dots, a_p, a_{p+1}, \dots, a_{p+q})(x, y)^{p+q}$$

where  $q$  is any positive integer.

It has already been shown, and will be otherwise exhibited presently, that gradients which are seminvariants exist only

when their weights, degrees, and extents are such as to make  $ip - 2w < 0$ . It has also been seen that a gradient which is a seminvariant of type  $w, i, p$  is the coefficient of the highest power of  $x$ , i. e.  $x^{ip-2w}$ , in a covariant of

$$(a_0, a_1, a_2, \dots a_p)(x, y)^p,$$

and, more generally, the coefficient of the highest power  $x^{i(p+q)-2w}$  of  $x$  in a covariant of the quantic of higher order

$$(a_0, a_1, a_2, \dots a_p, a_{p+1}, \dots a_{p+q})(x, y)^{p+q}.$$

Thus, for instance,  $a_0^2 a_3 - 3a_0 a_1 a_2 + 2a_1^3$ , a gradient annihilated by  $\Omega$ , for which  $w = 3, i = 3, p = 3, ip - 2w = 3$  is a seminvariant of  $(a_0, a_1, a_2, \dots a_{3+q})(x, y)^{3+q}$ , where  $q$  is zero or any positive integer, and is the leading coefficient of covariants

$$(a_0^2 a_3 - 3a_0 a_1 a_2 + 2a_1^3)x^3 + \dots$$

$$(a_0^2 a_3 - 3a_0 a_1 a_2 + 2a_1^3)x^6 + \dots$$

$$(a_0^2 a_3 - 3a_0 a_1 a_2 + 2a_1^3)x^9 + \dots$$

&c., &c.,

of the cubic  $(a_0, a_1, a_2, a_3)(x, y)^3,$

the quartic  $(a_0, a_1, a_2, a_3, a_4)(x, y)^4,$

the quintic  $(a_0, a_1, a_2, a_3, a_4, a_5)(x, y)^5,$

&c., &c., respectively.

It has also been seen (§ 112) that a gradient  $G_{w, i, p}$  of type  $w, i, p$ , which satisfies

$$\Omega G_{w, i, p} = 0,$$

and is such that  $ip - 2w = 0$ , has the further property of satisfying the fourth linear differential equation

$$O G_{w, i, p} = 0,$$

where  $O \equiv pa_1 \frac{d}{da_0} + (p-1)a_2 \frac{d}{da_1} + \dots + a_p \frac{d}{da_{p-1}}$ ,

so that it is an *invariant* of the  $p$ -ic

$$(a_0, a_1, a_2, \dots a_p)(x, y)^p,$$

while still only a seminvariant of the higher binary quantics

$$(a_0, a_1, a_2, \dots a_p, a_{p+1}, \dots a_{p+q})(x, y)^{p+q}.$$

Thus, for instance,

$$a_0 a_4 - 4 a_1 a_3 + 3 a_2^2,$$

for which  $w = 4$ ,  $i = 2$ ,  $p = 4$ ,  $ip - 2w = 0$ , is an invariant of the quartic

$$(a_0, a_1, a_2, a_3, a_4)(x, y)^4,$$

and a seminvariant of the quintic, sextic, &c.,

$$(a_0, a_1, \dots, a_4, a_5)(x, y)^5,$$

$$(a_0, a_1, \dots, a_4, a_5, a_6)(x, y)^6,$$

&c.,

being the leading coefficient in covariants

$$(a_0 a_4 - 4 a_1 a_3 + 3 a_2^2) x^2 + \dots,$$

$$(a_0 a_4 - 4 a_1 a_3 + 3 a_2^2) x^4 + \dots,$$

&c.,

of the quintic, sextic, &c., respectively.

We need not then dissociate invariants from seminvariants in searching for them by means of the annihilator  $\Omega$ . A seminvariant found will be in particular an invariant for the binary  $p$ -ic, in case the *excess* (to use another word of Sylvester's)  $ip - 2w$  vanishes. It has already been established, and will again appear, that there is no rational integral invariant of a binary  $p$ -ic which is not thus given.

119.] A seminvariant of extent  $p$  involves all of  $a_0, a_1, a_2, \dots, a_p$ . Suppose if possible that  $a_r$ , where  $r < p$ , is absent from a seminvariant  $S$  of extent  $p$ . We may write the fact

$$\Omega S = 0$$

in the form

$$(r+1) a_r \frac{d}{da_{r+1}} S + \left\{ a_0 \frac{d}{da_1} + 2 a_1 \frac{d}{da_2} + \dots + r a_{r-1} \frac{d}{da_r} \right. \\ \left. + (r+2) a_{r+1} \frac{d}{da_{r+2}} + \dots + p a_{p-1} \frac{d}{da_p} \right\} S = 0.$$

Now, on our supposition, all of the left-hand side but

$$(r+1) a_r \frac{d}{da_{r+1}} S$$

is free from  $a_r$ . But the sum vanishes. Therefore

$$a_r \frac{d}{da_{r+1}} S$$

is free from  $a_r$ . Therefore  $\frac{d}{da_{r+1}} S = 0$ , i.e.  $S$  is free from  $a_{r+1}$ .

A seminvariant free from  $a_r$  is thus free from  $a_{r+1}$ , and therefore from  $a_{r+2}$ , from  $a_{r+3}$ , &c. Finally it is free from  $a_p$ .

Our supposition then that a seminvariant of extent  $p$  exists which does not contain all of  $a_0, a_1, a_2, \dots a_p$  is untenable.

120.] An invariant of a binary  $p$ -ic involves all the coefficients. Being a seminvariant, it must by the preceding article involve all of  $a_0, a_1, a_2, \dots a_r$  if it extend as far as  $a_r$ . Also being an anti-seminvariant, annihilated by  $O$ , it must by the same reasoning involve all of  $a_p, a_{p-1}, a_{p-2}, \dots a_0$ , since, reckoning extent from  $a_p$  back to  $a_0$ , it extends to  $a_0$ .

121.] Repeated operations with  $\Omega$  and  $O$ . We proceed to pay attention to the results of operating with  $\Omega$  or with  $O$ , once or any number of times in succession, on any gradient whatever.

$$\text{The operator } \Omega \equiv a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + \dots + pa_{p-1} \frac{d}{da_p}$$

acting on any gradient  $G_{w, i, p}$  produces another gradient. The degree of the produced gradient is  $i$ , that of  $G_{w, i, p}$ . The weight of the produced gradient is  $w-1$ , where  $w$  is the weight of  $G_{w, i, p}$ . The extent of the produced gradient is either  $p$ , the extent of  $G_{w, i, p}$ , or less. These facts are clear

when we remark that any term of  $\Omega$ ,  $ra_{r-1} \frac{d}{da_r}$  for instance,

operating on a term involving  $a_r$  in  $G_{w, i, p}$ , on  $a_r^\rho a_s^\sigma a_t^\tau \dots$  for instance, has the effect of replacing that term by another,  $\rho ra_{r-1} a_r^{\rho-1} a_s^\sigma a_t^\tau \dots$ , of the same degree and weight one less, one suffix being diminished by unity and none increased.

Thus if  $G$  be a gradient of weight  $w$ , degree  $i$ , and extent not exceeding  $p$ ,  $\Omega G$  is a gradient of weight  $w-1$ , degree  $i$ , and extent not exceeding  $p$ .

Consequently  $\Omega^2 G = \Omega . \Omega G$  is a gradient of weight  $w-2$ , degree  $i$ , and extent not exceeding  $p$ .

And generally, by  $r$  repetitions of the  $\Omega$  process,  $\Omega^r G$  is a gradient of weight  $w-r$ , degree  $i$ , and extent not exceeding  $p$ .

This  $\Omega$  process cannot be repeated indefinitely without leading to a vanishing result. For a gradient of negative weight is an impossibility, the weight of a gradient being a sum of numbers chosen from among  $0, 1, 2, \dots p$ . Thus when we take  $w+1$  for  $r$ , if not sooner, we must have

$$\Omega^{w+1} G = 0,$$

and consequently also  $\Omega^{w+2} G = 0, \Omega^{w+3} G = 0, \&c.$

122.] We may reason in a similar way with regard to the operator

$$O \equiv p a_1 \frac{d}{da_0} + (p-1) a_2 \frac{d}{da_1} + \dots + a_p \frac{d}{da_{p-1}}.$$

Operation with this produces from a gradient of extent  $p$  or less another gradient of the same degree, of weight greater by one, and of extent not greater than  $p$ . The term *extent* is here used in the sense of the definition (§ 116). That the extent cannot be raised beyond  $p$  by operation with  $O$  results from the fact that  $O$  itself is taken as involving no letter  $a_r$  with a suffix (extent) greater than  $p$ . Thus,  $G$  being of weight  $w$ , degree  $i$ , and extent not exceeding  $p$ ,

$$O^r G$$

is a gradient of weight  $w+r$ , degree  $i$ , and extent not exceeding  $p$ .

Here, again, the succession of gradients produced is not indefinitely continued. For the greatest possible weight of a product of  $i$  constituents of weights chosen from among  $0, 1, 2, 3, \dots p$  is  $ip$ , that of  $a^i_p$ . Consequently

$$O^{ip-w} G$$

can be nothing more than a (non-vanishing or vanishing) multiple of  $a^i_p$ , and therefore

$$O^{ip-w+1} G = 0, O^{ip-w+2} G = 0, \&c.$$

We may also consider the results of successive operations with both of  $\Omega$  and  $O$  in any order on a gradient. The



conclusion, drawn readily from the above, is that,  $G$  being any gradient of weight  $w$ , degree  $i$ , and extent  $p$  or less,

$$\Omega^{m_1} O^{n_1} \Omega^{m_2} O^{n_2} \Omega^{m_3} O^{n_3} \dots G,$$

where  $m_1, m_2, m_3, \dots, n_1, n_2, n_3, \dots$  are positive numbers, or one or more of them zero, is, unless it vanish, a gradient of weight

$$w - m_1 - m_2 - m_3 - \dots + n_1 + n_2 + n_3 + \dots,$$

of degree  $i$ , and of extent not exceeding  $p$ .

123.] The alternant of  $\Omega$  and  $O$ . The operators  $\Omega, O$  are linear: but this is not the case with the operators  $\Omega^2, O^2, \Omega O, O \Omega$ , &c. Thus, for instance,

$$\begin{aligned} \Omega^2 &\equiv \left( a_0 \frac{d}{da_1} + 2 a_1 \frac{d}{da_2} + \dots + p a_{p-1} \frac{d}{da_p} \right)^2 \\ &\equiv 1 \cdot 2 a_0 \frac{d}{da_2} + 2 \cdot 3 a_1 \frac{d}{da_3} + \dots + (p-1) p a_{p-2} \frac{d}{da_p} \\ &\quad + \left\{ a_0^2 \frac{d^2}{da_1^2} + 4 a_0 a_1 \frac{d^2}{da_1 da_2} + 4 a_1^2 \frac{d^2}{da_2^2} + \dots \right\}. \end{aligned}$$

Moreover the operators  $\Omega, O$  are not commutative; i. e. the compound operators  $\Omega O$  and  $O \Omega$  are not identical in meaning. Thus while

$$\begin{aligned} \Omega O &\equiv \left( a_0 \frac{d}{da_1} + 2 a_1 \frac{d}{da_2} + \dots + p a_{p-1} \frac{d}{da_p} \right) \\ &\quad \left( p a_1 \frac{d}{da_0} + (p-1) a_2 \frac{d}{da_1} + \dots + a_p \frac{d}{da_{p-1}} \right) \\ &\equiv 1 \cdot p a_0 \frac{d}{da_0} + 2 (p-1) a_1 \frac{d}{da_1} + 3 (p-2) a_2 \frac{d}{da_2} + \dots \\ &\quad + p \cdot 1 a_{p-1} \frac{d}{da_{p-1}} \\ &\quad + \text{terms involving } \frac{d^2}{da_0 da_1}, \frac{d^2}{da_1^2}, \dots, \frac{d^2}{da_{p-1} da_p}, \\ O \Omega &\equiv p \cdot 1 a_1 \frac{d}{da_1} + (p-1) 2 a_2 \frac{d}{da_2} + (p-2) 3 a_3 \frac{d}{da_3} + \dots \\ &\quad + 1 \cdot p a_p \frac{d}{da_p} \\ &\quad + \text{the same terms in } \frac{d^2}{da_0 da_1}, \dots \text{ as in } \Omega O. \end{aligned}$$

We thus see, however, that the two compound operators differ only in their linear parts. The non-linear parts of both are just the algebraical product of  $\Omega$  and  $O$ . This leads us to consider the difference  $\Omega O - O\Omega$  of the two compound operators, the *alternant*, as it is called, of  $\Omega$  and  $O$ . It is always a fact that the alternant  $\theta\phi - \phi\theta$  of two linear operators  $\theta$  and  $\phi$  is a linear operator. In the present case

$$\begin{aligned} \Omega O - O\Omega &\equiv p a_0 \frac{d}{da_0} + \{2(p-1) - 1 \cdot p\} a_1 \frac{d}{da_1} \\ &\quad + \{3(p-2) - 2(p-1)\} a_2 \frac{d}{da_2} + \dots \\ &\quad + \{p \cdot 1 - (p-1)2\} a_{p-1} \frac{d}{da_{p-1}} - p a_p \frac{d}{da_p} \\ &\equiv p a_0 \frac{d}{da_0} + (p-2) a_1 \frac{d}{da_1} + (p-4) a_2 \frac{d}{da_2} + \dots \\ &\quad - (p-2) a_{p-1} \frac{d}{da_{p-1}} - p a_p \frac{d}{da_p} \\ &\equiv p \left\{ a_0 \frac{d}{da_0} + a_1 \frac{d}{da_1} + a_2 \frac{d}{da_2} + \dots + a_p \frac{d}{da_p} \right\} \\ &\quad - 2 \left\{ a_1 \frac{d}{da_1} + 2 a_2 \frac{d}{da_2} + \dots + p a_p \frac{d}{da_p} \right\}. \end{aligned}$$

Now let the operation be on  $G$  a gradient of degree  $i$  and weight  $w$ .  $G$  satisfies the two linear differential equations of § 117, which express its homogeneity and isobarism. Using the two equations, we see that what we are led to is

$$\begin{aligned} (\Omega O - O\Omega)G &= p \cdot iG - 2 \cdot wG \\ &= (ip - 2w)G. \end{aligned}$$

124.] An important application of this result has already been mentioned in § 97. Let the gradient  $G$  be an invariant  $I$  of  $(a_0, a_1, a_2, \dots, a_p)(x, y)^p$ , so that  $\Omega I = 0$  and  $O I = 0$ , and therefore  $O\Omega I = 0$  and  $\Omega O I = 0$ . We have the consequence that  $(ip - 2w)I = 0$ , i.e. that the degree and weight of an invariant of a binary  $p$ -ic are connected with  $p$  by the relation

$$ip - 2w = 0.$$

As another application let the gradient  $G$  be a semi-invariant  $S$ . Then  $\Omega S = 0$ , and therefore  $O\Omega S = 0$ , so that

$$\Omega O S = (ip - 2w)S,$$

which tells us that the result of operating first with  $O$  and then with  $\Omega$  on a seminvariant of extent not exceeding  $p$  is to reproduce that seminvariant multiplied by a numerical factor.

This is in accord with the conclusions of §§ 109, 110.

125.] **Alternant of  $\Omega$  and  $O'$ .** Important information is to be gathered from the alternants of  $\Omega$  and  $O^2, O^3, \dots$  which, though not linear, have simple equivalents when the functions on which they operate are gradients.

It is assumed throughout this article and in what follows, except where otherwise stated, that the operation is on a gradient  $G$  of weight  $w$ , degree  $i$ , and extent not exceeding  $p$ . For brevity the  $G$  is not as a rule written.

The 'excess'  $ip - 2w$ , in which  $p$  is always the suffix of the highest element which occurs in  $\Omega$  and  $O$ , and may, it must be remembered, be greater than the extent of  $G$ , is, also for brevity, denoted by  $\eta$ .

Thus instead of writing

$$(\Omega O - O \Omega)G = (ip - 2w)G,$$

we write merely  $\Omega O - O \Omega = \eta$ . ...(1)

Now notice that

$$\Omega O^2 - O^2 \Omega = (\Omega O - O \Omega)O + O(\Omega O - O \Omega),$$

and also observe that,  $G$  which is operated on being of weight  $w$  and degree  $i$ ,  $OG$  is by § 122 of weight  $w + 1$  and degree  $i$ , so that the excess for  $OG$ , corresponding to  $\eta$  for  $G$ , is  $ip - 2(w + 1) = \eta - 2$ . Thus

$$\begin{aligned} \Omega O^2 - O^2 \Omega &= (\eta - 2)O + O \eta \\ &= 2(\eta - 1)O, \end{aligned} \quad \dots(2)$$

since  $\eta$ , being numerical, is commutative with  $O$ .

Again

$$\begin{aligned} \Omega O^3 - O^3 \Omega &= (\Omega O - O \Omega)O^2 + O(\Omega O - O \Omega)O + O^2(\Omega O - O \Omega) \\ &= (\eta - 4)O^2 + O(\eta - 2)O + O^2 \eta, \end{aligned}$$

the excess for  $O^2G$  being  $ip - 2(w + 2)$ , i.e.  $\eta - 4$ ,

$$= 3(\eta - 2)O^2. \quad \dots(3)$$

In like manner we notice generally that the excess for  $O^r G$  is  $ip - 2(w + r)$ , i. e.  $\eta - 2r$ , and that generally

$$\begin{aligned} \Omega O^r - O^r \Omega &= (\Omega O - O \Omega) O^{r-1} + O(\Omega O - O \Omega) O^{r-2} \\ &\quad + O^2(\Omega O - O \Omega) O^{r-3} + \dots + O^{r-1}(\Omega O - O \Omega) \\ &= (\eta - 2 \cdot \overline{r-1}) O^{r-1} + O(\eta - 2 \cdot \overline{r-2}) O^{r-2} \\ &\quad + O^2(\eta - 2 \cdot \overline{r-3}) O^{r-3} + \dots + O^{r-1} \eta \\ &= \{r\eta - 2(1 + 2 + 3 + \dots + \overline{r-1})\} O^{r-1} \\ &= r(\eta - r + 1) O^{r-1}. \end{aligned} \quad \dots(R)$$

Ex. 1. Deduce that

$$O \Omega^r - \Omega^r O = r(-\eta - r + 1) \Omega^{r-1}.$$

Ex. 2. Prove that

$$\begin{aligned} \Omega^r O^r &= \Omega^{r-1} O^{r-1} \{O \Omega + r(\eta - r + 1)\} \\ &= (O \Omega + 1 \cdot \eta) (O \Omega + 2 \cdot \overline{\eta-1}) (O \Omega + 3 \cdot \overline{\eta-2}) \\ &\quad \dots (O \Omega + r \cdot \overline{\eta-r+1}) \\ &= \Omega O (\Omega O + \eta - 2) (\Omega O + 2 \cdot \overline{\eta-3}) \dots (\Omega O + \overline{r-1} \cdot \overline{\eta-r}). \end{aligned}$$

Ex. 3. In like manner

$$\begin{aligned} O^r \Omega^r &= (\Omega O - 1 \cdot \eta) (\Omega O - 2 \cdot \overline{\eta+1}) (\Omega O - 3 \cdot \overline{\eta+2}) \\ &\quad \dots (\Omega O - r \cdot \overline{\eta+r-1}) \\ &= O \Omega (O \Omega - \eta - 2) (O \Omega - 2 \cdot \overline{\eta+3}) \dots (O \Omega - \overline{r-1} \cdot \overline{\eta+r}). \end{aligned}$$

Ex. 4. Prove that

$$\begin{aligned} \Omega^r O^r \cdot \Omega^s O^s &= \Omega^s O^s \cdot \Omega^r O^r, \\ O^r \Omega^r \cdot O^s \Omega^s &= O^s \Omega^s \cdot O^r \Omega^r, \\ \Omega^r O^r \cdot O^s \Omega^s &= O^s \Omega^s \cdot \Omega^r O^r. \end{aligned}$$

Ex. 5. Prove by mathematical induction that

$$\begin{aligned} O^r \Omega^s &= \Omega^s O^r - (\eta - r + s) r s \Omega^{s-1} O^{r-1} \\ &\quad + \frac{(\eta - r + s)(\eta - r + s + 1)}{1 \cdot 2} r(r-1) s(s-1) \Omega^{s-2} O^{r-2} - \dots \end{aligned}$$

(Hilbert.)

126.] **The excess non-negative for a seminvariant.** Use of the results of the preceding article gives a proof (Sylvester's) of the fact (§ 112) that for no seminvariant can the 'excess'  $ip - 2w$  be negative. Since, if  $S$  be a seminvariant of extent

or less,  $\Omega S = 0$ , the results give

$$\begin{aligned}\Omega O . S &= \eta S, \\ \Omega O^2 . S &= 2(\eta - 1)OS, \\ \Omega O^3 . S &= 3(\eta - 2)O^2S, \\ &\dots \dots \dots \\ \Omega O^r . S &= r(\eta - r + 1)O^{r-1}S, \\ &\text{\&c., \&c.}\end{aligned}$$

If  $\eta$ , or  $ip - 2w$ , be negative, the coefficients on the right in these equalities form a numerically increasing series of negative numbers. None of them can vanish. Now (§ 122) there must be a number  $r$ , equal to or less than  $ip - w + 1$ , for which and all greater numbers  $O^r S = 0$ , and consequently  $\Omega O^r S = 0$ . The  $r$ th of the above equalities gives then

$$0 = r(\eta - r + 1)O^{r-1}S,$$

and therefore  $O^{r-1}S = 0$ . This necessitates  $\Omega O^{r-1}S = 0$ , and this again, by the  $(r-1)$ th equality, that

$$0 = (r-1)(\eta - r + 2)O^{r-2}S,$$

i.e.  $O^{r-2}S = 0$ . Proceeding thus backwards step by step, we eventually find from the first equality that

$$0 = \eta S,$$

i.e. that

$$S = 0,$$

since  $\eta$  is negative and not zero. In other words, the supposition that there is a seminvariant  $S$  for which  $\eta$  is negative is untenable.

We repeat that the  $\eta$  which it is here proved cannot be negative is  $ip - 2w$  where  $p$  is the greatest suffix occurring in  $\Omega$  and  $O$ . The extent  $p'$  of a seminvariant of

$$(a_0, a_1, a_2, \dots, a_p)(x, y)^p,$$

if not  $p$  itself, is less than  $p$ . In this latter case the seminvariant is also one of  $(a_0, a_1, a_2, \dots, a_{p'}) (x, y)^{p'}$ , and we might have taken  $p'$  as our  $p$  in the above reasoning. Thus, if  $p'$  be the extent of a seminvariant of weight  $w$  and degree  $i$ ,  $ip' - 2w$  cannot be negative.

EX. 6. Use the results of this and the preceding article to prove the theorem of § 111, that any seminvariant  $S$ , of extent not exceeding  $p$ , leads a covariant of order  $\eta$  of  $(a_0, a_1, a_2, \dots, a_p)$   $(x, y, y^p)$ , i.e. that  $x^\eta e^{\frac{y}{x} O} S$  is not fractional in  $x$ , and is annihilated by  $\Omega - y \frac{d}{dx}$ .

*Ans.*  $\Omega O^{\eta+1} S = 0$ . Therefore  $O^{\eta+1} S$ , whose excess is negative,  $= 0$ . The coefficients in the result of operating with  $\Omega - y \frac{d}{dx}$  are of the form  $\frac{1}{r!} \{ \Omega O^r - r(\eta - r + 1) O^{r-1} \} S$ , i.e. of the form  $\frac{1}{r!} O^r \Omega S$ , which vanishes.

127.] It is clear that the above reasoning, which shows there to be no seminvariant with a negative  $\eta$ , has no application to the cases of  $\eta$  zero and  $\eta$  positive. In these cases one of the series of multipliers  $\eta, \eta - 1, \eta - 2, \eta - 3, \dots$  on the right of the equalities of the last article vanishes. Thus from the fact that  $O^{\eta+1} S = 0$  it does not follow that  $O^\eta S = 0$ . It is the factor  $\eta - \eta$  on the right of the critical equality which vanishes, and not the other factor  $O^\eta S$ .

For a positive or vanishing  $\eta$ , a number

$$(w; i, p) - (w - 1; i, p)$$

has been found in §§ 113, 101 which cannot exceed the number of linearly independent seminvariants (or invariants) of weight  $w$ , degree  $i$ , and extent  $p$  or less. It is now to be proved that we have a means of assuring ourselves that the number is precise.

This famous theorem, stated by Cayley and much used, remained long without proof, and was even doubted. The first demonstration of it was given by Sylvester, by means of the results of § 125. The method to be here given is different from his, but is based upon the same results, though, as we shall see, an alternative basis is the theorem of § 100.

128.] **Exactness of Cayley's number of linearly independent seminvariants of given type.** Let  $G$  be any gradient whatever of degree  $i$ , weight  $w$ , and extent not exceeding  $p$ . Let  $\eta$  be  $ip - 2w$  the excess for  $G$ . For  $\Omega G, \Omega^2 G, \Omega^3 G, \&c.$  the excesses are  $\eta + 2, \eta + 4, \eta + 6, \&c.$

Take the operative equalities (1) to (R) of § 125, and operate, not always on  $G$ , but on  $G$  in the first case, on  $\Omega G$  in

the second, on  $\Omega^2 G$  in the third, and so on, so that  $\eta$  has to be replaced by  $\eta, \eta + 2, \eta + 4, \eta + 6, \dots$  in the successive cases. We obtain

$$\begin{aligned} \Omega O G - O \Omega G &= \eta G, \\ \Omega O^2 \Omega G - O^2 \Omega^2 G &= 2(\eta + 1) O \Omega G, \\ \Omega O^3 \Omega^2 G - O^3 \Omega^3 G &= 3(\eta + 2) O^2 \Omega^2 G, \\ &\dots \\ \Omega O^r \Omega^{r-1} G - O^r \Omega^r G &= r(\eta + r - 1) O^{r-1} \Omega^{r-1} G, \\ &\text{\&c., \&c.} \end{aligned}$$

Multiply the first of these by  $\frac{1}{\eta}$ , the second by  $-\frac{1}{2 \cdot \eta(\eta + 1)}$ , the third by  $\frac{1}{2 \cdot 3 \cdot \eta(\eta + 1)(\eta + 2)}$ , ..., the  $r$ th by

$$(-1)^{r-1} \frac{1}{2 \cdot 3 \dots r \cdot \eta(\eta + 1) \dots (\eta + r - 1)}$$

&c., and add. We thus obtain

$$\begin{aligned} \Omega O \left\{ \frac{1}{1 \cdot \eta} - \frac{1}{1 \cdot 2 \cdot \eta(\eta + 1)} O \Omega \right. \\ \left. + \frac{1}{1 \cdot 2 \cdot 3 \cdot \eta(\eta + 1)(\eta + 2)} O^2 \Omega^2 - \dots \right\} G = G, \end{aligned}$$

since for a great value of  $r$  the residual multiple of  $O^r \Omega^r G$  does not exist, for  $\Omega^{w+1} G$  vanishes, and therefore  $O^r \Omega^r G$  vanishes if  $r$  be  $w + 1$  or more.

Consequently, if  $\eta$  be positive, the result of operating on the gradient

$$\begin{aligned} O \left\{ \frac{1}{1 \cdot \eta} - \frac{1}{1 \cdot 2 \cdot \eta(\eta + 1)} O \Omega \right. \\ \left. + \frac{1}{1 \cdot 2 \cdot 3 \cdot \eta(\eta + 1)(\eta + 2)} O^2 \Omega^2 - \dots \right\} G \end{aligned}$$

with  $\Omega$  is to produce  $G$ .

The gradient is a finite one, for though the operative series is regarded as continuing to infinity it produces really only a finite number of terms, since  $O^r \Omega^r G$  vanishes when  $r$  exceeds  $w$  if not earlier.

We have thus proved that any gradient whatever, of weight  $w$ , degree  $i$ , and extent  $p$  or less, for which  $\eta = ip - 2w$  is positive, can be obtained by operating with  $\Omega$  on some

gradient or other of weight  $w+1$ , degree  $i$ , and extent  $p$  or less. The same was otherwise proved in § 100.

Now for  $w$  write  $w-1$ . It follows that every gradient of weight  $w-1$ , degree  $i$ , and extent  $p$  or less, can be obtained by operation with  $\Omega$  on some gradient of weight  $w$ , degree  $i$ , and extent  $p$  or less, provided that  $ip-2(w-1)$  is positive, i. e. that  $ip-2w \lessdot -1$ .

This tells us that if we write down the most general gradient of weight and degree  $w, i$ , and of extent  $p$  or less, where  $ip-2w \lessdot -1$  and operate on it with  $\Omega$ , the result must be the most general gradient of weight and degree  $w-1, i$ , and of extent  $p$  or less. For the arbitrary coefficients in the first gradient may be so chosen that the derived gradient may be any one of its type we choose, and so in particular may be any single product of its type we choose.

In other words, if the general gradient  $G'$  be such that  $ip-2w \lessdot -1$  the coefficients in the derived gradient  $\Omega G'$  are all linearly independent.

Now, in the notation of §§ 101, 113,  $G'$  contains

$$(w; i, p)$$

terms, and  $\Omega G'$  contains

$$(w-1; i, p)$$

terms. If  $G'$  be a seminvariant  $\Omega G' = 0$ , and conversely, i. e. the coefficients of these  $(w-1; i, p)$  terms have separately to vanish. These are all independent, by the above, if  $ip-2w \lessdot -1$ , and are linear functions of the  $(w; i, p)$  arbitrary coefficients in  $G'$ . Their vanishing determines then  $(w-1; i, p)$  of the coefficients in  $G'$  in terms of the rest. In other words, exactly

$$(w; i, p) - (w-1; i, p)$$

are left arbitrary. This, then, is the exact number of linearly independent seminvariants of weight  $w$ , degree  $i$ , and extent not greater than  $p$ .

Ex. 7. It may be proved by aid of § 125, Ex. 5 that, when the operation is on a gradient for which  $\eta = ip-2w > 0$ , the operator of § 100,

$$1 - \frac{\Omega O}{1^2} + \frac{\Omega^2 O^2}{1^2 \cdot 2^2} - \frac{\Omega^3 O^3}{1^2 \cdot 2^2 \cdot 3^2} + \dots,$$



and that of the present article,

$$1 - \frac{\Omega O}{1 \cdot \eta} + \frac{\Omega O^2 \Omega}{1 \cdot 2 \cdot \eta (\eta + 1)} - \frac{\Omega O^3 \Omega^2}{1 \cdot 2 \cdot 3 \cdot \eta (\eta + 1) (\eta + 2)} + \dots,$$

are identical. (*Proc. Lond. Math. Soc.* Vol. XXIV. p. 23.)

Ex. 8. If  $G$  be a gradient for which  $\eta$  is *negative* prove that

$$\left(1 - \frac{O \Omega}{1^2} + \frac{O^2 \Omega^2}{1^2 \cdot 2^2} - \frac{O^3 \Omega^3}{1^2 \cdot 2^2 \cdot 3^2} + \dots\right) G = 0,$$

and

$$\left(1 + \frac{O \Omega}{1 \cdot \eta} + \frac{O \Omega^2 O}{1 \cdot 2 \cdot \eta (\eta - 1)} + \frac{O \Omega^3 O^2}{1 \cdot 2 \cdot 3 \cdot \eta (\eta - 1) (\eta - 2)} + \dots\right) G = 0.$$

129.] **Arithmetical conclusions.** Some arithmetical conclusions of interest with regard to numbers of partitions may be drawn from results at which we have arrived.

Since the most general gradient of type  $w-1, i, p$  can when  $ip - 2w \ll -1$  be derived by operation with  $\Omega$  from the most general gradient of type  $w, i, p$ , the former cannot contain more arbitrariness than the latter, i. e. more terms than the latter. Hence if  $ip - 2w \ll -1$

$$(w; i, p) \ll (w-1; i, p).$$

Again, we have shown in § 126 and elsewhere that if  $ip - 2w < 0$  there is no seminvariant, and in § 128 that if  $ip - 2w \ll -1$  there are exactly  $(w; i, p) - (w-1; i, p)$  seminvariants. The case  $ip - 2w = -1$  is included in both categories. The conclusion from this case of  $ip - 2w = -1$ , i. e. of  $w = \frac{1}{2}(ip + 1)$ , where  $i$  and  $p$  must clearly both be odd, is that if  $i$  and  $p$  be any odd numbers

$$\left(\frac{ip+1}{2}; i, p\right) = \left(\frac{ip-1}{2}; i, p\right).$$

This is only a particular case of the fact that, for any  $w$  not exceeding  $ip$ , whatever numbers  $i$  and  $p$  be,

$$(w; i, p) = (ip - w; i, p),$$

which is immediately seen by noticing that the products of weight  $w$  and those of weight  $ip - w$  are conjugate in pairs. If, in fact,  $a_0^{a_0} a_1^{a_1} \dots a_p^{a_p}$  is one of the first type, the conjugate one of the second type is  $a_p^{a_0} a_{p-1}^{a_1} \dots a_0^{a_p}$ .

130.] **Reciprocal partitions.** The number  $(w; i, p)$  is, it will be remembered, the number of ways in which the number  $w$  may be written as a sum of  $i$  or fewer numbers, none exceeding  $p$ . It is an important fact that this number is also the number of ways in which  $w$  may be written as a sum of  $p$  or fewer numbers, none exceeding  $i$ ; in other words, that

$$(w; i, p) = (w; p, i).$$

The following proof is due to Ferrers. Another will present itself in the next chapter.

Let any partition of  $w$  into  $i$  or fewer parts, none exceeding  $p$ , be the partition into  $n_1 + n_2 + n_3 + \dots + n_i$ , where no part is greater than  $p$  nor than the preceding part, and where one or more at the end may be zero. Write down  $n_1$  dots in a row. Next write  $n_2$  dots under the first  $n_2$  of these dots in a second row. Then write in a third row  $n_3$  dots under the first  $n_3$  of the  $n_2$  dots: and so on, till in all  $n_1 + n_2 + n_3 + \dots + n_i = w$  dots have been written. We have thus visibly arranged a partition of  $w$  into  $i$  or fewer parts, none exceeding  $p$ .

Now read the arrangement by columns instead of rows. We have in the first column a number,  $m_1$ , of dots not greater than  $i$ . In the second column we have, say,  $m_2$  dots where  $m_2 \succ m_1$  and so  $\succ i$ . In the third we have  $m_3$  dots where  $m_3 \succ m_2$  and therefore  $\succ i$ : and so on. Finally, in the  $p$ th column we have either no dot or a number  $m_p$  of dots not greater than any previous  $m$ , and so not greater than  $i$ . We have thus visibly arranged a partition of  $w$  into

$$m_1 + m_2 + m_3 + \dots + m_p$$

a sum of  $p$  or fewer numbers, none greater than  $i$ .

Thus to every one of the  $(w; i, p)$  partitions we have a conjugate one of the  $(w; p, i)$  partitions. Similarly, considering columns first and then rows, to every one of the  $(w; p, i)$  partitions there is a conjugate one of the  $(w; i, p)$  partitions. And no two of the one set of partitions have the same conjugate in the other set, for a definite arrangement in the one way is also definite in the other. Consequently the numbers  $(w; i, p)$  and  $(w; p, i)$  are equal.

131.] **Hermite's law of reciprocity.** Hence we obtain a famous and most prolific theorem due to Hermite.

Since  $(w; i, p) = (w; p, i)$

for all numbers  $w$  it follows that

$$(w; i, p) - (w-1; i, p) = (w; p, i) - (w-1; p, i).$$

Accordingly:—*The number of rational integral seminvariants of weight  $w$ , degree  $i$ , and extent not exceeding  $p$ , is equal to the number of rational integral seminvariants of weight  $w$ , degree  $p$ , and extent not exceeding  $i$ .*

In particular take  $ip = 2w$ . We are told that:—*The number of invariants (i.e. linearly independent rational integral invariants) of degree  $i$  of a binary  $p$ -ic is equal to the number of invariants of degree  $p$  of a binary  $i$ -ic.*

Again take  $ip - 2w > 0$ , and denote  $ip - 2w$  by  $\varpi$ . Then, since when  $i, p, w$  are known  $\varpi$  is known, we may enunciate:—*The number of covariants of degree  $i$  (in the coefficients) and order  $\varpi$  (in the variables) of a binary  $p$ -ic is equal to the number of covariants of degree  $p$  and order  $\varpi$  of a binary  $i$ -ic.*

There are, of course, other ways of arriving at this law of reciprocity. For instance, we may take it in connexion with the one to one correspondence which clearly exists between the hyperdeterminant symbols (§ 60)

$$\frac{n_{12}}{12} \frac{n_{13}}{13} \frac{n_{23}}{23} \dots,$$

where the number of figures 1, 2, 3, ... is  $i$ , and where  $n_{12} + n_{13} + n_{23} + \dots$  is  $\frac{1}{2}(ip - \varpi)$ , for covariants of a binary  $p$ -ic, and the root expressions (§ 111)

$$\Sigma (a_1 - a_2)^{n_{12}} (a_1 - a_3)^{n_{13}} (a_2 - a_3)^{n_{23}} \dots,$$

where  $i$  is the number of roots, and where

$$n_{12} + n_{13} + n_{23} + \dots = w' = \frac{1}{2}(ip - \varpi),$$

for the seminvariant leaders, divided by powers of  $\alpha_0$ , of covariants of a binary  $i$ -ic.

This is, however, probably best regarded as a proof from the law of reciprocity that hyperdeterminants form a complete system of covariants, rather than as a proof of the law of reciprocity from this fact. The idea has been developed by Sylvester.

Ex. 9. A binary  $4n$ -ic has one invariant of degree 3, and a binary quantic whose order is not a multiple of 4 has none. (Cayley.)

Ans. Since all invariants of a cubic are powers of the discriminant whose degree is 4.

Ex. 10. The symbol of this invariant is  $\frac{-2n}{12} \frac{-2n}{23} \frac{-2n}{31}$ . Prove that if  $u, v, w$  all denote the quantic the invariant may be written

$$\frac{-2n-1}{12} \frac{-2n-1}{23} \frac{-2n-1}{31} \left| \begin{array}{ccc} \frac{d^2u}{dx^2} & \frac{d^2u}{dxdy} & \frac{d^2u}{dy^2} \\ \frac{d^2v}{dx^2} & \frac{d^2v}{dxdy} & \frac{d^2v}{dy^2} \\ \frac{d^2w}{dx^2} & \frac{d^2w}{dxdy} & \frac{d^2w}{dy^2} \end{array} \right|,$$

where  $u, v, w$  are not to be made identical till all the operations are performed; and hence that the invariant is a linear function of the determinants

$$\left\| \begin{array}{c} a_0, a_1, a_2, a_3, \dots, a_{4n-2} \\ a_1, a_2, a_3, a_4, \dots, a_{4n-1} \\ a_2, a_3, a_4, a_5, \dots, a_{4n} \end{array} \right\|. \quad (\text{Cayley.})$$

Ex. 11. A binary  $p$ -ic has as many invariants of degree 4 as there are ways of choosing positive integral or zero values of  $m$  and  $n$  to satisfy  $2m + 3n = p$ . (Cayley.)

Ans. Assume, as will be shown later, that  $I$  and  $J$  are the only irreducible invariants of the quartic.

Ex. 12. A binary  $p$ -ic has a single or no  $p$ -ic covariant of the second degree in the coefficients according as  $p$  is or is not a multiple of 4. (Cayley.)

Ans. Since covariants of equal order and degree of a quadratic must have the form  $(ac - b^2)^n (ax^2 + 2bxy + cy^2)^{2n}$ .

Ex. 13. The one invariant (Ex. 9) of degree 3 of a  $4n$ -ic is the lineo-linear invariant of the  $4n$ -ic and the covariant of Ex. 12. (Cayley.)

Ex. 14. A binary  $p$ -ic has as many covariants of degree 2 in the coefficients as there are solutions of  $2m + n = p$  in positive integers (and zeros); and  $2n$  is the order of any such covariant in the variables. (Hermite.)

Ex. 15. A binary quantic of odd order has a covariant of the second order and the second degree. (Hermite.)

Ex. 16. A binary quantic of order  $4n + 2$ , where  $n$  is any number, has a covariant of the second order and third degree. (Hermite.)

Ans. Use Ex. 15 for the case of the cubic.

Ex. 17. By the two preceding examples binary quantics whose orders are of the forms  $4n+1$ ,  $4n+2$ ,  $4n+3$  have quadratic covariants. Use the facts that a quintic has a quadratic covariant of degree 8, the Jacobian of the cubic covariant of § 17, Ex. 20 and the linear covariant of § 57, Exx. 16, 17, and an invariant of degree 4 (§ 61, Ex. 30) to complete the proof that every binary quantic except the quartic has a quadratic covariant whose degree in the coefficients does not exceed 5. (*Hermite.*)

Ex. 18. No covariant or invariant of the second degree in the coefficients can have an odd weight. In particular, no invariant of the second degree can be skew.

Ex. 19. No invariant of the third degree can be skew.

Ex. 20. A binary quantic of any odd order greater than 3 has a linear covariant of degree 5. (*Hermite.*)

*Ans.* Use the fact that a quintic has linear covariants of degrees 5 and 7 (§ 57, Ex. 17) and an invariant of degree 4 (§ 61, Ex. 30).

132.] **Gradients in more sets than one.** Just as we have dealt with gradients in one set of quantities  $a_0, a_1, a_2, \dots, a_p$  in the present chapter, we may deal with gradients in more sets than one  $a_0, a_1, a_2, \dots, a_{p_1}; b_0, b_1, b_2, \dots, b_{p_2}; c_0, c_1, c_2, \dots, c_{p_3}; \dots$ . We have merely throughout to insert  $\Sigma\Omega$ ,  $\Sigma O$ , and

$$i_1 p_1 + i_2 p_2 + i_3 p_3 + \dots - 2w$$

for  $\Omega$ ,  $O$ , and  $ip - 2w$ . As in § 119 a seminvariant which involves one letter of any set involves all the previous letters of that set. As in § 120 an invariant of the quantics whose coefficients are the sets involves all the coefficients of any one if it involve one of them. As in § 125

$$\Sigma\Omega (\Sigma O)^r - (\Sigma O)^r \Sigma\Omega = r (\eta - r + 1) (\Sigma O)^{r-1},$$

where

$$\eta = \Sigma(ip) - 2w.$$

As in § 126 there can be no seminvariant for which  $\Sigma(ip) - 2w$  is negative. As in § 128 any gradient for which  $\Sigma(ip) - 2w$  is positive can be written as the result of operating with  $\Sigma\Omega$  on another gradient, and hence if  $\Sigma(ip) - 2w \leq -1$  the exact number of linearly independent seminvariants of weight  $w$  and partial degrees  $i_1, i_2, i_3, \dots$  is

$$(w; i_1, p_1; i_2, p_2; i_3, p_3, \dots) - (w-1; i_1, p_1; i_2, p_2; i_3, p_3; \dots),$$

where  $(w; i_1, p_1; i_2, p_2; i_3, p_3; \dots)$  denotes the number of

ways in which  $w$  may be written as the sum of  $i_1$  or fewer numbers not greater than  $p_1$ , of  $i_2$  or fewer not greater than  $p_2$ , of  $i_3$  or fewer not greater than  $p_3$ , &c. Finally we have arithmetical conclusions corresponding to those of § 129.

The generalization of § 130 is also immediate. It will readily be seen, by considering as many Ferrers' diagrams as there are sets of  $a$ 's,  $b$ 's,  $c$ 's, &c., containing altogether  $w$  dots, that in

$$(w; i_1, p_1; i_2, p_2; i_3, p_3; \dots)$$

$i_1$  and  $p_1$ , or  $i_2$  and  $p_2$ , or  $i_3$  and  $p_3$ , ..., or more than one or all of these pairs, may be interchanged without altering the number of partitions. Hence a generalization of the law of reciprocity is easy.

Ex. 21. The number of covariants of any degree  $i$  and of order  $p$  of a binary  $p$ -ic, i.e. of invariants of partial degrees  $i, p$  of a  $p$ -ic and a linear form, is equal to the number of invariants of partial degrees  $i, 1$  of two binary  $p$ -ics.

Ex. 22. Prove that

$$(w; i_1, p_1; i_2, p_2; i_3, p_3; \dots) = \Sigma \Sigma \Sigma \dots (v_1; i_1, p_1) (v_2; i_2, p_2) (v_3; i_3, p_3) \dots,$$

where the summations indicate that to  $v_1, v_2, v_3, \dots$  are to be given all positive integral and zero values which make

$$v_1 + v_2 + v_3 + \dots = w. \quad (\text{Franklin.})$$

## CHAPTER VIII.

### GENERATING FUNCTIONS.

133.] UNFORTUNATELY no practically convenient algebraical formula<sup>1</sup> is known which gives in all cases the number of partitions denoted by  $(w ; i, p)$ , i.e. the number of ways in which the number  $w$  may be formed by adding together  $i$  or fewer numbers, every one of which is one of  $1, 2, 3, \dots, p$ , or, which is the same thing, of  $i$  parts, every one of which is one of  $0, 1, 2, 3, \dots, p$ . For tabulation of such numbers of partitions recourse must be had to a method known as that of *Generating Functions*.

The origin of the theory of numbers of partitions is due to Euler. The theory in its application to invariants, &c., was first studied by Cayley with a view to and in his second memoir on quantics (*Collected Works*, Vol. II). The subsequent writings on the subject are very numerous. Cayley himself, Sylvester, Franklin, MacMahon and Hammond as well as others have devoted themselves to it with remarkable success.

The investigation of the number  $(w ; i, p) - (w - 1 ; i, p)$  of linearly independent or 'asyzygetic' seminvariants of given weight, degree, and extent by means of generating functions is only a preliminary object of the researches. The ulterior aims are the discovery of the number and types of the *irreducible* concomitants of a binary quantic, and of the relations or syzygies which connect those irreducible concomitants.

The subject being a vast one only an introduction to it can be given here. We consider only quantics of the first few orders. In passing from order to order the complexity of the investigations necessary enormously increases.

---

<sup>1</sup> For a formula due to Brioschi see Faa de Bruno's *Formes Binaires*, § 89.

134.] **Generating function for  $(w ; i, p)$ .** By a *Generating Function* we mean a function of one or more variables which, when it is expanded in powers of that variable, or powers and products of powers of those variables, has for the general coefficient of a power or product of powers the number of an assigned class which is determined by the index of that power, or the indices of those powers. It may be that only a limited range of the coefficients is relevant. For instance, the expansion may be an infinite one, but the class of numbers a finite one given by the coefficients of a limited range of terms, the indices of other terms being parameters irrelevant to the matter we have in hand.

We proceed to see that a generating function can be formed whose expansion is

$$(0 ; i, p) + (1 ; i, p)z + (2 ; i, p)z^2 + \dots \\ + (w ; i, p)z^w + \dots + (ip ; i, p)z^{ip},$$

and which accordingly, when  $i$  and  $p$  are known, gives the number of partitions  $(w ; i, p)$  as the coefficient of  $z^w$  in its development.

It is at once clear that, by definition of  $(w ; i, p)$ , this number of partitions is the number of ways in which positive integral, or vanishing, values of  $r_0, r_1, r_2, \dots, r_p$  can be found which satisfy the two equations

$$r_0 + r_1 + r_2 + \dots + r_p = i, \\ r_1 + 2r_2 + \dots + pr_p = w.$$

Term  
( $a_0^{r_0} a_1^{r_1} a_2^{r_2} \dots a_p^{r_p}$ )

Now this number is the coefficient of  $z^w x^i$  in the product

$$(1 + x + x^2 + \dots + x^{r_0} + \dots)(1 + zx + z^2x^2 + \dots + z^{r_1}x^{r_1} + \dots) \\ (1 + z^2x + z^4x^2 + \dots + z^{2r_2}x^{r_2} + \dots) \dots \\ (1 + z^p x + z^{2p}x^2 + \dots + z^{pr_p}x^{r_p} + \dots),$$

where the series forming any factor may if we please be extended to infinity. This product may be written

$$\{(1-x)(1-zx)(1-z^2x) \dots (1-z^p x)\}^{-1}.$$

It can also be multiplied out and arranged according to ascending powers of  $x$ . Suppose that, thus arranged, it is

$$u_0 + u_1x + u_2x^2 + \dots + u_ix^i + \dots ;$$



then  $u_0 = 1$ , and  $u_1, u_2, \dots u_i, \dots$  are functions of  $z$ . In fact,

$$u_i = (0; i, p) + (1; i, p)z + (2; i, p)z^2 + \dots + (w; i, p)z^w + \dots + (ip; i, p)z^{ip}.$$

Notice that  $(0; i, p)$  denotes 1. This is reasonable, for there is one partition of zero into  $i$  parts not exceeding  $p$ , namely, into  $i$  zeros.  $a_0^i$  is the one corresponding term when we are thinking of gradients. In particular, by convention, we may think of  $(0; 0, p)$  as denoting 1.

Now in

$$1 + u_1x + u_2x^2 + \dots + u_ix^i + \dots = \{(1-x)(1-zx)(1-z^2x) \dots (1-z^px)\}^{-1},$$

put  $zx$  for  $x$ , getting

$$\begin{aligned} 1 + u_1zx + u_2z^2x^2 + \dots + u_iz^ix^i + \dots \\ = \{(1-zx)(1-z^2x)(1-z^3x) \dots (1-z^{p+1}x)\}^{-1} \\ = \frac{1-x}{1-z^{p+1}x} \{1 + u_1x + u_2x^2 + \dots + u_ix^i + \dots\}. \end{aligned}$$

Here multiply through by  $1 - z^{p+1}x$ , and equate the coefficients of  $x^i$ , the equality being identical. We obtain

$$u_iz^i - u_{i-1}z^{p+i} = u_i - u_{i-1},$$

so that

$$\begin{aligned} u_i &= u_{i-1} \frac{1 - z^{p+i}}{1 - z^i} \\ &= u_{i-2} \frac{1 - z^{p+i-1}}{1 - z^{i-1}} \cdot \frac{1 - z^{p+i}}{1 - z^i} \\ &= \dots \dots \dots \\ &= u_0 \frac{1 - z^{p+1}}{1 - z} \cdot \frac{1 - z^{p+2}}{1 - z^2} \cdot \dots \cdot \frac{1 - z^{p+i-1}}{1 - z^{i-1}} \cdot \frac{1 - z^{p+i}}{1 - z^i} \\ &= \frac{(1 - z^{p+1})(1 - z^{p+2}) \dots (1 - z^{p+i})}{(1 - z)(1 - z^2) \dots (1 - z^i)}. \end{aligned}$$

Consequently  $(w; i, p)$  is the coefficient of  $z^w$  in the expansion of this function in ascending powers of  $z$ . Notice that it is incidentally proved that this expansion is a terminating one of degree  $ip$ , i.e. that the numerator of the generating function  $u_i$  is divisible by the denominator whatever numbers  $i$  and  $p$  be.

Notice also that  $u_i$  is exactly  $z^{ip}$  times the result of replacing

in it  $z$  by  $\frac{1}{z}$ . Coefficients equidistant from the beginning and the end in the development are then equal. We have thus a proof of one of the facts of § 129, i.e. that

$$(ip - w ; i, p) = (w ; i, p).$$

135.] **Generating function for number of seminvariants of given type.** It is easy to deduce a generating function in which the coefficient of  $z^w$  is the difference of numbers of partitions  $(w ; i, p) - (w - 1 ; i, p)$ . This difference is the number of linearly independent seminvariants of

$$(a_0, a_1, a_2, \dots, a_p)(x, y)^p$$

whose weight and degree are  $w$  and  $i$ , if  $ip > 2w$ , and the number of invariants of the type if  $ip = 2w$ . For values of  $w$ , such that  $ip - 2w < 0$ , or rather  $< -1$ , we are not really concerned with the difference in connexion with seminvariants.

Since

$$u_i = (0 ; i, p) + (1 ; i, p)z + (2 ; i, p)z^2 + \dots \\ + (w ; i, p)z^w + \dots + (ip ; i, p)z^{ip},$$

the value of  $(w ; i, p) - (w - 1 ; i, p)$ , for values of  $w$  from 1 to  $ip$  inclusive, is the coefficient of  $z^w$  in  $(1 - z)u_i$ , i.e. in the development of

$$\frac{(1 - z^{p+1})(1 - z^{p+2})(1 - z^{p+3}) \dots (1 - z^{p+i})}{(1 - z^2)(1 - z^3) \dots (1 - z^i)}.$$

This development is a terminating one of degree  $ip + 1$ .

Notice that from the last remark of the preceding article the middle coefficient in the development, if there be one, i.e. if  $ip$  is odd, vanishes; and that coefficients equidistant from the beginning and end are equal but of opposite signs. Now (§ 129) we know that when  $ip - 2w \leq -1$ , i.e. when  $w \geq \frac{1}{2}(ip + 1)$ , the difference  $(w ; i, p) - (w - 1 ; i, p)$  is never negative. The development of the generating function consists then of a series of terms with positive coefficients followed by a series with the same coefficients taken negatively in reversed order.

A word as to the first coefficient  $(0 ; i, p) = 1$ , which is not presented as a difference. It is correctly the number of semin-

variants of degree  $i$  and zero weight. The one is  $a_0^i$ . We may if we like regard it as a difference  $(0; i, p) - (-1; i, p)$  like the rest. For  $(-1; i, p)$  is of course zero.

136.] **Reciprocity.** The generating function of § 134 may, upon multiplication of numerator and denominator by

$$(1-z)(1-z^2) \dots (1-z^p),$$

be written

$$\frac{(1-z)(1-z^2)(1-z^3) \dots (1-z^{p+i})}{(1-z)(1-z^2) \dots (1-z^p) \cdot (1-z)(1-z^2) \dots (1-z^i)},$$

which is unaltered by interchange of  $i$  and  $p$ . It may, in fact, be written

$$\frac{(1-z^{i+1})(1-z^{i+2}) \dots (1-z^{i+p})}{(1-z)(1-z^2) \dots (1-z^p)}.$$

The coefficient of  $z^w$  in its expansion is then  $(w; p, i)$  for exactly the same reason that it is  $(w; i, p)$ . Thus we have another proof of the theorem of § 130 that

$$(w; i, p) = (w; p, i),$$

and of Hermite's law of reciprocity (§ 131).

The generating function of § 135 may also be written

$$\frac{(1-z^{i+1})(1-z^{i+2})(1-z^{i+3}) \dots (1-z^{i+p})}{(1-z^2)(1-z^3) \dots (1-z^p)}.$$

Ex. 1. Prove by aid of generating functions that

$$(w; i, p) - (w; i-1, p) = (w-i; i, p-1).$$

Ex. 2. Prove that

$$(w; i, p) = (w; 0, p-1) + (w-1; 1, p-1) + (w-2; 2, p-1) + \dots + (w-i; i, p-1).$$

137.] **The whole number of seminvariants and invariants of given degree.** The whole number of linearly independent seminvariants, including invariants if there be any, of degree  $i$  which a binary  $p$ -ic possesses may be found as follows.

We have seen (§§ 111, 126) that there are none for which  $ip - 2w$  is negative. Thus the greatest weight of any is  $\frac{1}{2} ip$  or  $\frac{1}{2}(ip-1)$ , according as  $i$  and  $p$  are not or are both odd. Call this maximum weight  $W$ .

The number of seminvariants (or invariants if  $W$  be  $\frac{1}{2}ip$ ) of weight  $W$  is

$$(W; i, p) - (W-1; i, p);$$

the number (all seminvariants necessarily) of weight  $W-1$  is

$$(W-1; i, p) - (W-2; i, p);$$

the number of weight  $W-2$  is

$$(W-2; i, p) - (W-3; i, p);$$

and so on. Finally the number of weight zero is unity or

$$\cdot (0; i, p).$$

Upon addition we have then for the whole number

$$(W; i, p).$$

We restore to  $W$  its value, and have the two following results.

(1) Unless  $i$  and  $p$  are both odd, the whole number of linearly independent seminvariants and invariants (i.e. of covariants and invariants) of degree  $i$  in the coefficients of the binary  $p$ -ic is

$$\left(\frac{ip}{2}; i, p\right),$$

and these consist of

$$\left(\frac{ip}{2}; i, p\right) - \left(\frac{ip}{2} - 1; i, p\right)$$

invariants, and  $\left(\frac{ip}{2} - 1; i, p\right)$

seminvariants (covariants).

(2) If  $i$  and  $p$  are both odd the whole number of linearly independent seminvariants, i.e. of covariants, of degree  $i$  in the coefficients is

$$\left(\frac{ip-1}{2}; i, p\right),$$

none of them being invariants.

By § 134 the whole number of degree  $i$  is thus seen to be the coefficient of  $z^{\frac{1}{2}ip}$  or  $z^{\frac{1}{2}(ip-1)}$  as the case may be in the development of

$$\frac{(1-z^{p+1})(1-z^{p+2}) \dots (1-z^{p+i})}{(1-z)(1-z^2) \dots (1-z^i)},$$

or its equivalent

$$\frac{(1-z^{i+1})(1-z^{i+2}) \dots (1-z^{i+p})}{(1-z)(1-z^2) \dots (1-z^p)}.$$

We can now illustrate by a few simple cases the way in which generating functions give information as to the number and nature of *irreducible* concomitants.

138.] Has a linear form invariants or covariants? For the linear form  $ax + by$ ,  $p = 1$ . The whole number of linearly independent seminvariants (including invariants) of degree  $i$  is by the preceding article

$$\text{co. } z^W \text{ in developement of } \frac{1 - z^{i+1}}{1 - z},$$

where  $W = \frac{1}{2}i$  or  $\frac{1}{2}(i-1)$  according as  $i$  is even or odd,

$$= \text{co. } z^W \text{ in } 1 + z + z^2 + \dots + z^i$$

$$= 1.$$

Thus of each degree there is a single seminvariant. What it is is clear. For degree 1 it is  $a$ , and for degree  $i$  it is  $a^i$ . It is a seminvariant and not an invariant, for  $ip - 2w = i > 0$ . The covariant which it leads is  $(ax + by)^i$ , the  $i$ -th power of the linear form itself.

Thus a linear form has no invariant, and its only covariants are powers of itself.

139.] Irreducible concomitants of a quadratic. For the quadratic  $ax^2 + 2bxy + cy^2$ ,  $p = 2$ . Here  $W = \frac{1}{2}2i = i$ . The whole number of linearly independent seminvariants and invariants of degree  $i$  is then

$$\begin{aligned} \text{co. } z^i \text{ in developement of } & \frac{(1 - z^{i+1})(1 - z^{i+2})}{(1 - z)(1 - z^2)} \\ = \text{ " " " } & (1 - z)^{-1}(1 - z^2)^{-1} \\ = \text{ " " " } & (1 + z + z^2 + z^3 + \dots) \\ & (1 + z^2 + z^4 + z^6 + \dots) \\ = \text{co. } z^i \text{ in } & 1 + z + 2z^2 + 2z^3 + 3z^4 + 3z^5 + \dots \end{aligned}$$

There is, then, one of degree 1, viz.  $a$ ; and there are two of degree 2, viz. the square  $a^2$  of the one of degree 1 and another distinct from it, which we know otherwise to be the discriminant  $ac - b^2$ . Of any higher degree  $i$  we see, by considering the product  $(1 + z + z^2 + z^3 + \dots)(1 + z^2 + z^4 + z^6 + \dots)$ , that there are just as many as there are ways of making up the number  $i$  as a sum of multiples, including zero multiples, of 1 and 2; and

these are of course the products of powers of the two independent ones  $a, ac - b^2$  of degrees 1, 2. For instance, if  $r + 2s$  is one of the partitions of  $i$  in question,  $a^r(ac - b^2)^s$  is a seminvariant of degree  $i$ . All seminvariants of the quadratic are then rationally and integrally expressible in terms of the two  $a$  and  $ac - b^2$ , without the necessity of introducing any other. The binary quadratic has therefore no irreducible seminvariant besides  $a$  and  $ac - b^2$ .

The second of these is an invariant. The first leads the quadratic  $ax^2 + 2bxy + cy^2$  itself. Consequently the complete system of irreducible concomitants of the binary quadratic consists of the quadratic itself and its discriminant. (Cf. § 85.)

Had we been looking for the irreducible *invariants* only we might have taken the generating function of § 135. For the quadratic the weight of an invariant of degree  $i$  is  $\frac{1}{2}2i = i$ . Thus the number of invariants of degree  $i$

$$\begin{aligned} &= \text{co. } z^i \text{ in developement of } \frac{(1 - z^{i+1})(1 - z^{i+2})}{1 - z^2} \\ &= \text{ " " " } (1 - z^2)^{-1} \\ &= \text{co. } z^i \text{ in } 1 + z^2 + z^4 + z^6 + \dots \end{aligned}$$

There is then no invariant of any odd degree, and a single one of every even degree. Thus there is one, the discriminant  $ac - b^2$ , of the second degree, and no other which is irreducible, all others being powers of this. (Cf. § 78.)

140.] **Invariants of a cubic.** Take  $p = 3$ , the case of the binary cubic  $(a, b, c, d)(x, y)^3$ ; and first consider the question of *invariants* only.

An invariant of degree  $i$  is of weight  $\frac{3i}{2}$ . For there to be one then  $i$  must be even.

The number of degree  $i$  is, by § 135,

$$\begin{aligned} &\text{co. } z^{\frac{3i}{2}} \text{ in developement of } \frac{(1 - z^{i+1})(1 - z^{i+2})(1 - z^{i+3})}{(1 - z^2)(1 - z^3)} \\ &= \text{ " " " } \frac{1 - z^i(z + z^2 + z^3)}{(1 - z^2)(1 - z^3)} \\ &= \text{ " " " } \frac{1}{(1 - z^2)(1 - z^3)} - \frac{z^{i+1}}{(1 - z)(1 - z^2)} \end{aligned}$$

$$\begin{aligned}
 &= \text{co. } z^{\frac{3i}{2}} \text{ in developement of } \frac{1}{(1-z^2)(1-z^3)} \\
 &\quad - \text{co. } z^{\frac{i}{2}} \text{ in developement of } \frac{z}{(1-z)(1-z^2)} \\
 &= \text{co. } z^{\frac{3i}{2}} \text{ in developement of } \frac{1}{(1-z^2)(1-z^3)} - \frac{z^3}{(1-z^3)(1-z^6)} \\
 &= \text{co. } z^{3i} \quad \quad \quad \text{''} \quad \frac{1}{(1-z^4)(1-z^6)} - \frac{z^6}{(1-z^6)(1-z^{12})} \\
 &= \quad \quad \quad \text{''} \quad \quad \quad \frac{1+z^4+z^8-z^6}{(1-z^6)(1-z^{12})} \\
 &= \quad \quad \quad \text{''} \quad \quad \quad \frac{1-z^6}{(1-z^6)(1-z^{12})},
 \end{aligned}$$

since the terms  $z^4$  and  $z^8$  in the numerator cannot when multiplied by powers of  $z^6$  and  $z^{12}$  produce terms of form  $z^{3i}$ ,

$$\begin{aligned}
 &= \text{co. } z^{3i} \text{ in developement of } \frac{1}{1-z^{12}} \\
 &= \text{co. } z^i \quad \quad \quad \text{''} \quad \frac{1}{1-z^4} \\
 &= \text{co. } z^i \text{ in } 1+z^4+z^8+z^{12}+\dots
 \end{aligned}$$

Thus for a degree not divisible by 4 there is no invariant; and for a degree divisible by 4 there is a single one, which must accordingly be a power of the one of degree 4, i.e. the discriminant  $(ad-bc)^2 - 4(ac-b^2)(bd-c^2)$ . This then is the only irreducible invariant of the cubic. (Cf. § 78.)

141.] Irreducible concomitants and syzygy for a cubic. We seek now the complete system of *seminvariants* and *invariants* of the cubic.

Here, by § 137, the number that are linearly independent of degree  $i$  is the coefficient of  $z^{\frac{3i}{2}}$  or  $z^{\frac{3i-1}{2}}$ , according as  $i$  is even or odd, in the developement of

$$\frac{(1-z^{i+1})(1-z^{i+2})(1-z^{i+3})}{(1-z)(1-z^2)(1-z^3)}.$$

The two cases may be combined by saying that it is the

coefficient of  $z^{\frac{3i}{2}}$  in the development of

$$(1+z^{\frac{1}{2}}) \frac{(1-z^{i+1})(1-z^{i+2})(1-z^{i+3})}{(1-z)(1-z^2)(1-z^3)},$$

i.e. in that of

$$\frac{1-z^i(z+z^2+z^3)}{(1-z^{\frac{1}{2}})(1-z^2)(1-z^3)}$$

$$= \text{co. } z^{\frac{3i}{2}} \text{ in development of } \frac{1}{(1-z^{\frac{1}{2}})(1-z^2)(1-z^3)}$$

$$- \text{co. } z^{\frac{i}{2}} \text{ in development of } \frac{z}{(1-z^{\frac{1}{2}})(1-z)(1-z^2)}$$

$$= \text{co. } z^{\frac{3i}{2}} \text{ in development of } \frac{1}{(1-z^{\frac{1}{2}})(1-z^2)(1-z^3)} - \frac{z^3}{(1-z^{\frac{3}{2}})(1-z^3)(1-z^6)}$$

$$= \text{co. } z^{3i} \quad " \quad \frac{1}{(1-z)(1-z^4)(1-z^6)} - \frac{z^6}{(1-z^3)(1-z^6)(1-z^{12})}$$

$$= \quad " \quad " \quad \frac{(1+z+z^2)(1+z^4+z^8)-z^6}{(1-z^3)(1-z^6)(1-z^{12})}$$

$$= \quad " \quad " \quad \frac{1+z^9}{(1-z^3)(1-z^6)(1-z^{12})},$$

in the numerator of which all powers of  $z$  with indices not divisible by 3 have been omitted as incapable of producing  $z^{3i}$  when multiplied into powers of  $z$  arising from the denominator, where all indices are multiples of 3,

$$= \text{co. } z^i \text{ in development of } \frac{1+z^3}{(1-z)(1-z^2)(1-z^4)}$$

$$= \quad " \quad " \quad \frac{1-z^6}{(1-z)(1-z^2)(1-z^3)(1-z^4)}$$

$$= \quad " \quad " \quad \frac{(1-z^6)(1+z+z^2+\dots)}{(1+z^2+z^4+\dots)(1+z^3+z^6+\dots)(1+z^4+z^8+\dots)},$$

of which the first few terms are

$$1+z+2z^2+3z^3+5z^4+6z^5+8z^6+\dots$$



We have then the following conclusions, gathered from the form before multiplying out.

(1) There is one seminvariant of degree 1, arising from the factor  $(1-z)^{-1}$  in the reduced generating function. This is the seminvariant  $a$ .

(2) Besides the square of this there is another seminvariant of degree 2, arising from the factor  $(1-z^2)^{-1}$ . This is  $ac-b^2$ , the seminvariant which leads the covariant which in § 86 has been called  $H$ . Denote it by  $H'$ .

(3) Besides  $a^3$  and  $a(ac-b^2)$  there is another of degree 3. This is  $a^2d-3abc+2b^3$ , the leader of the covariant which in § 86 we have called  $G$ . Denote it by  $G'$ . This arises from the factor  $(1-z^3)^{-1}$ .

(4) Besides the four seminvariants of degree 4 which can be formed by compounding  $a$ ,  $H'$  and  $G'$  rationally, there is an additional one arising from the factor  $(1-x^4)^{-1}$ . This is the discriminant  $(ad-bc)^2-4(ac-b^2)(bd-c^2)$  which we have called  $\Delta$ . It is an invariant.

(5) There is no other irreducible seminvariant. For all the factors of the denominator of the prepared generating function are now exhausted, and there are no positive terms in the numerator except 1; and this tells us that there is nothing which in the developement can increase the coefficient of  $x^i$ , whatever  $i$  be, beyond the number of ways in which  $i$  can be made up of sums of multiples of 1, 2, 3, 4 the indices of the  $z$ ,  $z^2$ ,  $z^3$ ,  $z^4$  in the denominator.

There are then four, and only four, irreducible seminvariants, including the one invariant  $\Delta$ . All of degree higher than 4 can be expressed rationally and integrally in terms of these four  $a$ ,  $H'$ ,  $G'$ ,  $\Delta$ .

(6) But there is one fact more, given by the existence of the negative term  $-z^6$  in the numerator of the reduced generating function. The four  $a$ ,  $H'$ ,  $G'$ ,  $\Delta$ , though irreducible are not independent. A relation, or 'syzygy' as it is called, connects them. And this syzygy is of the sixth degree. The presence of the  $-z^6$  reduces the coefficient of  $z^6$  in the developement from 9, which would be its value were the numerator 1 only, to 8. The number of linearly independent seminvariants of degree 6 is then one less than the number of products of degree 6 of powers of  $a$ ,  $H'$ ,  $G'$  and  $\Delta$ . These

products are consequently connected by a linear relation which reduces the most general linear function of them to one with 8 arbitrary coefficients instead of 9. The products are

$$a^6, a^4H', a^3G', a^2H'^2, a^2\Delta, aH'G', H'^3, H'\Delta, G'^2.$$

The syzygy which connects them must of course connect a number of them which are of the same weight. Now their weights are

$$0, 2, 3, 4, 6, 5, 6, 8, 6,$$

the only three of which are the same are those of  $a^2\Delta$ ,  $H'^3$  and  $G'^2$ . The syzygy then connects these. It is found to be

$$a^2\Delta = G'^2 + 4H'^3,$$

which is of course the same relation as the

$$u^2\Delta = G^2 + 4H^3$$

of § 87. For  $a$ ,  $H'$ ,  $G'$  are the seminvariants which lead the covariants  $u$ ,  $H$ ,  $G$ ; and a syzygy connecting the seminvariant leaders of covariants connects also the covariants led, as otherwise by means of the syzygy we could form a covariant whose seminvariant leader vanishes, i.e. a covariant with  $y$  for a factor, which is impossible.

The complete system of irreducible concomitants of a binary cubic consists then of itself, its quadratic and cubic covariants  $H$  and  $G$ , and its discriminant  $\Delta$ . The four are connected by a syzygy of the sixth degree in the coefficients, and, it may be noticed, of the sixth order in the variables.

142.] Irreducible invariants of the quartic. For the case  $p = 4$ , that of the quartic, we confine attention to the investigation of the number of irreducible *invariants*.

Since here  $\frac{1}{2}ip = 2i$ , the number of linearly independent invariants of degree  $i$  is, by § 135,

co.  $z^{2i}$  in developement of

$$\frac{(1-z^{i+1})(1-z^{i+2})(1-z^{i+3})(1-z^{i+4})}{(1-z^2)(1-z^3)(1-z^4)}$$

$$= \text{co. } z^{2i} \text{ in developement of } \frac{1}{(1-z^2)(1-z^3)(1-z^4)} \\ - \frac{z^{i+1}}{(1-z^2)(1-z^3)(1-z)}$$

$$\begin{aligned}
&= \text{co. } z^{2i} \text{ in developement of } \frac{1}{(1-z^2)(1-z^3)(1-z^4)} \\
&\quad - \text{co. } z^i \text{ in developement of } \frac{z}{(1-z)(1-z^2)(1-z^3)} \\
&= \text{co. } z^{2i} \text{ in developement of } \frac{1}{(1-z^2)(1-z^3)(1-z^4)} \\
&\quad - \frac{z^2}{(1-z^2)(1-z^4)(1-z^6)} \\
&= \quad \quad \quad \quad \quad \quad \frac{1+z^3-z^2}{(1-z^2)(1-z^4)(1-z^6)} \\
&= \quad \quad \quad \quad \quad \quad \frac{1-z^2}{(1-z^2)(1-z^4)(1-z^6)},
\end{aligned}$$

for  $z^3$  in the numerator can be a factor only of odd powers in the developement,

$$\begin{aligned}
&= \text{co. } z^{2i} \text{ in developement of } \frac{1}{(1-z^4)(1-z^6)} \\
&= \text{co. } z^i \quad \quad \quad \quad \quad \quad \frac{1}{(1-z^2)(1-z^3)}.
\end{aligned}$$

Hence there are two, and only two, irreducible invariants, one of degree 2 and one of degree 3, since there are just as many linearly independent invariants of any higher degree as there are combinations of that degree of these two. The two are (§ 80)

$$I \equiv ae - 4bd + 3c^2,$$

$$J \equiv ace + 2bcd - ad^2 - b^2e - c^3.$$

There is no invariant of higher degree which cannot be expressed rationally and integrally in terms of them.

143.] **Invariants of the quintic and sextic.** The application of these methods has been continued a good many stages further. The labour and ingenuity required increase considerably as we advance.

For the case of the quintic,  $p = 5$ , the result is that the number of linearly independent invariants of degree  $i$  is the coefficient of  $z^i$  in the developement of

$$\frac{1-z^{36}}{(1-z^4)(1-z^8)(1-z^{12})(1-z^{18})}.$$

N

There are then *four* irreducible invariants of the quintic, of degrees 4, 8, 12, 18. They are not, however, independent, as the presence of  $-z^{36}$  in the numerator implies. This presence diminishes the number of linearly independent invariants of degree 36 to one below the number of ways of making up 36 by means of repetitions of 4, 8, 12, 18. In other words, there is a 'syzygy,' of degree 36 in  $a, b, c, d, e, f$ , the coefficients of the quintic, which connects the irreducible invariants of degrees 4, 8, 12, 18. This syzygy will be exhibited in a later chapter. It expresses the square of  $I_{18}$  in terms of  $I_4, I_8,$  and  $I_{12}$ .

For the sextic the number of linearly independent invariants of degree  $i$  is the coefficient of  $z^i$  in the development of

$$\frac{1 - z^{30}}{(1 - z^2)(1 - z^4)(1 - z^6)(1 - z^{10})(1 - z^{15})}$$

There are *five* irreducible invariants, of degrees 2, 4, 6, 10, 15; and these are connected by a syzygy of degree 30, which expresses the square of  $I_{15}$  rationally and integrally in terms of  $I_2, I_4, I_6,$  and  $I_{10}$ .

For the full investigation of these facts reference should be made to Cayley's second memoir on quantics.

144.] **Generating function for concomitants of given degree and order.** A new departure in the use of generating functions dates from Cayley's ninth memoir on quantics (*Collected Works*, Vol. VII). The earlier use of them had not succeeded in exhibiting complete systems of irreducible covariants for higher quantics than the quartic, and indeed mistaken inferences from it had indicated the erroneous conclusion that there were not complete systems of finite number. That there were had meanwhile been conclusively established by Gordan's method of transvectants. The two theories have now been completely reconciled, and verify one another's conclusions. The error arose from considering all syzygies independent, whereas there are syzygies of the second order connecting syzygies, for values of  $p$  exceeding 4.

Let us return to § 134, where it was shown that the number of linearly independent partitions of  $w$  into  $i$  or fewer parts,

none exceeding  $p$ , is the coefficient of  $z^w \xi^i$  (notice that we have changed the notation) in the developement of

$$\frac{1}{(1-\xi)(1-z\xi)(1-z^2\xi)\dots(1-z^p\xi)}$$

in ascending powers of  $\xi$ , and therefore of  $z$ ; and consequently that the number of linearly independent seminvariants of weight  $w$  and degree  $i$  of a binary  $p$ -ic,

i. e.  $(w; i, p) - (w-1; i, p),$

is the coefficient of  $z^w \xi^i$  in the developement of

$$\frac{1-z}{(1-\xi)(1-z\xi)(1-z^2\xi)\dots(1-z^p\xi)}$$

Here put  $x^{-2}$  for  $z$  and  $ax^p$  for  $\xi$ . The number of the seminvariants is therefore the coefficient of  $a^i x^{ip-2w}$  in the developement of

$$\frac{1-x^{-2}}{(1-ax^p)(1-ax^{p-2})(1-ax^{p-4})\dots(1-ax^{-p+4})(1-ax^{-p+2})(1-ax^{-p})}$$

in ascending powers of  $a$ , and powers of  $x$  ascending and descending as they present themselves.

Now  $ip-2w$  is the order  $\varpi$  of that covariant of the  $p$ -ic which any seminvariant of weight  $w$  and degree  $i$  leads. Consequently the number of linearly independent covariants of degree  $i$  and order  $\varpi$ , where  $\varpi$  is essentially non-negative, is the coefficient of  $a^i x^\varpi$  in the developement of this last written generating function. In particular, the number of linearly independent *invariants* of degree  $i$  is the coefficient of  $a^i$  in the part of the developement which is free from  $x$ .

For the quadratic, the cubic, and the quartic the generating functions are

$$\frac{1-x^{-2}}{(1-ax^2)(1-a)(1-ax^{-2})},$$

$$\frac{1-x^{-2}}{(1-ax^3)(1-ax)(1-ax^{-1})(1-ax^{-3})},$$

$$\frac{1-x^{-2}}{(1-ax^4)(1-ax^2)(1-a)(1-ax^{-2})(1-ax^{-4})}.$$

145.] **Reduced generating functions.** We need only those terms in the developement which involve positive powers of

$x$  as well as of  $a$ . Now it proves to be possible to separate those parts of the generating function, for a given value of  $p$ , which give rise to positive powers of  $x$  from those which give negative powers. It will be verified without difficulty that the three last written generating functions, for the cases of the quadratic, the cubic, and the quartic, may respectively be written

$$A(x) - x^{-2} A(x^{-1}),$$

$$B(x) - x^{-2} B(x^{-1}),$$

$$C(x) - x^{-2} C(x^{-1}),$$

where

$$A(x) = \frac{1}{(1-ax^2)(1-a^2)},$$

$$B(x) = \frac{1-a^6x^6}{(1-ax^3)(1-a^2x^2)(1-a^3x^3)(1-a^4)},$$

$$C(x) = \frac{1-a^6x^{12}}{(1-ax^4)(1-a^2)(1-a^2x^4)(1-a^3)(1-a^3x^6)}.$$

Hence the numbers of linearly independent covariants of degree  $i$  and order  $\varpi$  for the quadratic, cubic, and quartic are respectively the coefficients of  $a^i x^\varpi$  in the developements in ascending powers of  $a$  of  $A(x)$ ,  $B(x)$ , and  $C(x)$ .

From  $A(x)$  and  $B(x)$  we at once gather the information of §§ 139, 141 with regard to the irreducible concomitants of the quadratic and cubic respectively.

From  $C(x)$  we gather in like manner the full information as to the quartic. It has *five* irreducible concomitants whose degrees and orders are given by  $ax^4$ ,  $a^2$ ,  $a^2x^4$ ,  $a^3$ ,  $\bar{a}^3x^6$ . The first is the quartic itself

$$u \equiv (a, b, c, d, e)(x, y)^4;$$

the second is its invariant of the second degree

$$I \equiv ae - 4bd + 3c^2;$$

the third is its Hessian

$$H \equiv (ac - b^2)x^4 + \dots \equiv x^4 e^{\frac{y}{x} 0} (ac - b^2);$$

the fourth is its invariant of the third degree

$$J \equiv ace + 2bcd - ad^2 - b^2e - c^3;$$

and the fifth is its covariant of degree 3 and order 6

$$G \equiv (a^2d - 3abc + 2b^3)x^6 + \dots \equiv x^6 e^{\frac{y}{x}} (a^2d - 3abc + 2b^3).$$

The second term  $-a^6x^{12}$  in the numerator tells us that the five are connected by a syzygy of degree 6 and order 12, which reduces the number of linearly independent covariants of this degree and order to one below that of the number of products of the degree and order in question of  $u, I, H, J, G$ . This is readily found to be

$$Ju^3 - IHu^2 + G^2 + 4H^3 = 0.$$

As to invariants alone, the terms in  $A(x), B(x),$  and  $C(x)$  which are free from  $x$  are the developements of

$$\frac{1}{1-a^2},$$

$$\frac{1}{1-a^4},$$

$$\frac{1}{(1-a^2)(1-a^3)};$$

whence the information that the discriminant is the only invariant of a quadratic, and that of §§ 140, 142 for the cubic and quartic, is at once gathered.

146.] **Reduced and representative generating functions.**  
**The Quintic.** To methods and results for quantics above the fourth order we have only space to allude. Most of the investigations are due to Sylvester who, with the collaboration of Franklin, has obtained for quantics of the first ten and the twelfth orders the numbers and types of complete systems of concomitants, or rather, as he himself points out, the types and numbers of systems which must, if they err from completeness in the higher cases, err by defect and not by excess; the possibility of there being more arising from the fact that the labour of discovery has been reduced to tractable dimensions by the adoption as a fundamental postulate for all cases of a fact observed for the first six orders, viz. that new syzygies and irreducible concomitants do not exist for the same degree and order. To this postulate Hammond has shown that there is an exception in the case of the septic, for

which the investigations had previously indicated a speciality not occurring in other cases so far as examined.

The first step in the process is general for all cases, and consists in showing that, when, as in § 145, the 'crude' generating function of § 144 is written as the difference of two parts, one of which gives all the terms in the development which proceed by positive and zero powers of  $x$ , and the other those which proceed by negative powers, the former part may be written in the form

$$\frac{C_0 + C_1x + C_2x^2 + \dots}{(1-ax^p)(1-ax^{p-2})(1-ax^{p-4})\dots(1-a^k)(1-a^k)\dots},$$

where the order of the numerator in  $x$  is less than that of the denominator, and where  $C_0, C_1, C_2, \dots$  are finite rational and integral functions of  $a$ . This is called the *reduced* generating function.

The second step is one which has to be performed for the cases of quantics of successive orders separately. The numerator and denominator have to be multiplied by such factors as to reduce the latter to a product of  $1-ax^p$  and such factors as

$$1-a^ix^\varpi, 1-a^j,$$

where  $i$  and  $\varpi$  are recognized as the degree and order of an irreducible covariant, and  $j$  as the degree of a known irreducible invariant. In all cases examined but that of the septic this has proved to be so possible as to keep the numerator a finite expression. In the case of the septic, however, there is a factor  $1-a^{10}$  in the denominator, and no irreducible invariant exists whose degree is 10 or a multiple of 10. In this case then the denominator can only be given the required form by multiplying by the infinite series

$$1 + a^{10} + a^{20} + a^{30} + \dots,$$

so that for this case the multiplied numerator does not terminate.

The reduced generating function thus prepared is called the *representative* generating function. For orders of quantics which have been examined the denominators of the representative generating functions are products of  $1-ax^p$  and of factors of the simple forms

$$1-a^2x^\varpi, 1-a^j.$$



For the *quintic* ( $p = 5$ ) the numerator of the representative generating function proves to be

$$\begin{aligned}
 & 1 + a^3(x^3 + x^5 + x^9) + a^4(x^4 + x^6) + a^5(x + x^3 + x^7 - x^{11}) \\
 & + a^6(x^2 + x^4) + a^7(x + x^5 - x^9) + a^8(x^2 + x^4) + a^9(x^3 + x^5 - x^7) \\
 & + a^{10}(x^2 + x^4 - x^{10}) + a^{11}(x + x^3 - x^9) + a^{12}(x^2 - x^3 - x^{10}) \\
 & + a^{13}(x - x^7 - x^9) + a^{14}(x^4 - x^6 - x^8) + a^{15}(-x^7 - x^9) \\
 & + a^{16}(x^2 - x^6 - x^{10}) + a^{17}(-x^7 - x^9) + a^{18}(1 - x^4 - x^8 - x^{10}) \\
 & + a^{19}(-x^5 - x^7) + a^{20}(-x^2 - x^6 - x^8) + a^{23}(-x^{11}),
 \end{aligned}$$

and the denominator

$$(1 - ax^5)(1 - a^2x^2)(1 - a^2x^6)(1 - a^4)(1 - a^8)(1 - a^{12}).$$

The third step in the process is one of sifting, or 'tamisage' as it is called. We have certain irreducible concomitants, or say 'ground forms,' to use a common designation, represented in the denominator. The earlier terms, after the first 1, in the numerator are positive, and indicate the existence of other ground forms. Proceed onwards from term to term in the numerator. As long as the degree and order of a term  $a^i x^w$  in the numerator cannot be made up as a sum of the degrees and orders of previously occurring terms, we have revealed the existence of as many new ground forms of that degree and order as there are positive units in its coefficient. When we have reached a term whose degree and order can be made up as a sum of degrees and orders of ground forms whose existence has been previously revealed by the numerator, *not* also those represented in the denominator, the excess of the coefficient of that term above the number of ways in which this can be done is the number of ground forms of the degree and order of the term in question, diminished by the number of syzygies of the degree and order which connect ground forms that have previously occurred, in the denominator as well as in the numerator, but increased, as will presently happen, by the number of syzygies of the second order which connect previous syzygies and are of the degree and order in question.

For instance, regarding the representative generating function for the quintic above, the eight terms in the

numerator which immediately follow the first have all the coefficient + 1, and the degree-orders of these terms,  $a^3x^3$ ,  $a^3x^5$ ,  $a^3x^9$ ,  $a^4x^4$ ,  $a^4x^6$ ,  $a^5x$ ,  $a^5x^3$ ,  $a^5x^7$ , are such as to make it clear that none of the terms can be written as a product of powers of the preceding terms. They indicate, then, that besides the ground forms of degree-orders (1, 5), (2, 2), (2, 6), (4, 0), (8, 0), (12, 0) given by factors of the denominator, there are others of degree-orders (3, 3), (3, 5), (3, 9), (4, 4), (4, 6), (5, 1), (5, 3), (5, 7).

Again, the coefficient of the next term  $a^5x^{11}$  in the numerator is -1. This indicates that there must be one syzygy of degree 5 and order 11 connecting some of the fourteen ground forms of degree-orders less than (5, 11). It is found to connect the products  $C_{1,5} C_{4,6}$ ,  $C_{2,2} C_{3,9}$ ,  $C_{2,6} C_{3,5}$ , where  $C_{r,s}$  is the ground form of degree  $r$  and order  $s$ . In particular  $C_{1,5}$  is the quintic itself.

As soon as the degree-order ( $i, \varpi$ ) of each new ground form in succession is found, we may, if we please, alter the form of the representative generating function by multiplying its numerator and denominator by  $1 - a^i x^\varpi$ , and so put that ground form in the same position as those whose representatives were before in the denominator, thus narrowing the further search by means of the numerator. In particular, when we know all the ground forms, we may write the generating function with the product of all their representative factors  $1 - a^i x^\varpi$ ,  $1 - a^j$ , &c. in the denominator. The sifting of the numerator is then a process of search for syzygies only. This idea has been developed by Hammond. The  $A(x)$ ,  $B(x)$ ,  $C(x)$  of § 145 for the quadratic cubic and quartic are generating functions thus written.

Notice that the terms free from  $x$  in the developement of the representative generating function for the quintic above are the terms in the developement of the result of putting  $x = 0$  in it, i. e. of

$$\frac{1 + a^{18}}{(1 - a^4)(1 - a^8)(1 - a^{12})}$$

This then is the representative generating function for invariants of the quintic. It leads to the conclusions already stated in § 143.

There prove to be twenty-three ground forms of the quintic, of which four,  $I_4$ ,  $I_8$ ,  $I_{12}$ ,  $I_{18}$ , are invariants.

147.] Sylvester and Franklin have also exhibited generating functions for the whole number of *seminvariants* of any degree for the quantics they have studied.

Moreover, they have obtained representative generating functions of two or more quantics of low degrees, and studied their indications as to systems of ground forms.

For these researches, and for the full theories above illustrated, reference should be made to the first four volumes of the *American Journal of Mathematics*. A few exercises are here left to the student.

Ex. 3. By § 137 the whole number of seminvariants (including invariants) of degree  $i$  of a binary  $p$ -ic is  $\left(\frac{ip}{2}; i, p\right)$  or  $\left(\frac{ip-1}{2}; i, p\right)$ , according as  $ip$  is even or odd. Show by the method of § 144 that this number is the coefficient of  $a^i$  in the part of the development of

$$\frac{1+x^{-1}}{(1-ax^p)(1-ax^{p-2})\dots(1-ax^{-p+2})(1-ax^{-p})},$$

in ascending powers of  $a$ , which is free from  $x$ .

Ex. 4. Prove that the number of linearly independent seminvariants of weight  $w$  and partial degrees  $i, i'$  of a  $p$ -ic and a  $p'$ -ic is the coefficient of  $z^w \xi^i \xi'^{i'}$  in the expansion of

$$\frac{1-z}{(1-\xi)(1-z\xi)\dots(1-z^p\xi) \cdot (1-\xi')(1-z\xi')\dots(1-z^{p'}\xi')}.$$

Ex. 5. Show, as in § 144, that the number of covariants of order  $\omega$  and partial degrees  $i, i'$  of a  $p$ -ic and a  $p'$ -ic is the coefficient of  $a^i a'^{i'} x^\omega$  in the development of

$$\frac{1-x^{-2}}{ax^p(1-ax^{p-2})\dots(1-ax^{-p+2})(1-ax^{-p}) \cdot (1-a'x^{p'})(1-a'x^{p'-2})\dots(1-a'x^{-p'+2})(1-a'x^{-p'})}$$

Ex. 6. Show that for two linear forms the reduced generating function for numbers of concomitants is

$$\frac{1}{(1-ax)(1-a'x)(1-aa')},$$

where  $a$  refers to one form and  $a'$  to the other.

Ex. 7. Show that for a linear form and a quadratic it is

$$\frac{1+abx}{(1-ax)(1-bx^2)(1-b^2)(1-a^2b)},$$

where  $a$  refers to the linear form and  $b$  to the quadratic.

Ex. 8. Show that for two quadratics it is

$$\frac{1 + bb'x^2}{(1 - bx^2)(1 - b'x^2)(1 - b^2)(1 - b'^2)(1 - bb')}.$$

148.] **Real generating functions.** From representative generating functions Cayley has passed to what he calls *Real Generating Functions*.

Let us return to § 145. The generating function

$$A(x) \equiv \frac{1}{(1 - ax^2)(1 - a^2)}$$

for the quadratic has told us that there are two ground forms, the quadratic  $u$  and the discriminant  $\Delta \equiv ac - b^2$ , and that there are just as many concomitants of any type as there are products of that type of powers of  $u$  and  $\Delta$ . This tells us that all concomitants of the quadratic are terms which actually occur in the expansion of

$$\frac{1}{(1 - u)(1 - \Delta)}.$$

This is the real generating function for the quadratic.

Again, for the cubic, that  $B(x)$ , which write in the form

$$\frac{1 + a^3x^3}{(1 - ax^3)(1 - a^2x^2)(1 - a^4)},$$

is the generating function, has told us that there are as many concomitants of any type as there are such products of  $u$ ,  $H$ ,  $\Delta$  and  $G$  of that type as do not involve  $G$  to a higher power than the first. And this information is exactly expressed by saying that concomitants of the cubic are linear functions of those products of the four ground forms which occur in the development of

$$\frac{1 + G}{(1 - u)(1 - H)(1 - \Delta)},$$

which is the real generating function of the cubic.

Once more, for the quartic,  $C(x)$ , or

$$\frac{1 + a^3x^6}{(1 - ax^4)(1 - a^2)(1 - a^2x^4)(1 - a^5)},$$

tells us in like manner that there is a real generating function

$$\frac{1 + G}{(1-u)(1-I)(1-H)(1-J)},$$

such that all concomitants of the quartic are linear functions of terms which actually occur in its development, i.e. of the products into which  $G^2$  does not enter.

For the quintic, and beyond, the form of a real generating function derived from the representative generating function of § 146 is not unique, owing to the number of different ways in which we may replace the many terms in the numerator by products of ground forms. Cayley has shown in his tenth memoir that the most useful form into which a real generating function of the quintic can be thrown is

$$\frac{\Sigma P(1-Q)}{(1-u)(1-C_{2,2})(1-C_{2,6})(1-I_4)(1-I_8)(1-I_{12})},$$

where every  $P$  and  $Q$ , in the products whose sum is the numerator, are products of ground forms and powers of ground forms chosen from among the complete system of 23 whose forms will be exhibited in a later chapter. All the 23 occur in the numerator and denominator together.

For invariants alone, real generating functions are

(1) for the quadratic  $\frac{1}{1-\Delta}$ ;

(2) for the cubic  $\frac{1}{1-\Delta}$ , not the same  $\Delta$  of course as in (1);

(3) for the quartic  $\frac{1}{(1-I)(1-J)}$ ;

(4) for the quintic  $\frac{1 + I_{18}}{(1-I_4)(1-I_8)(1-I_{12})}$ .

## CHAPTER IX.

### HILBERT'S PROOF OF GORDAN'S THEOREM.

149.] AN irreducible invariant has, it will be remembered, been defined as one which cannot be expressed rationally and integrally in terms of invariants of lower degree than its own belonging to the same quantic or quantics.

Similarly, an irreducible covariant is one which cannot be expressed rationally and integrally in terms of covariants and invariants of degree in the coefficients lower than its own.

In the cases of binary quantics of low orders, it has been seen in the last chapter that the number of irreducible invariants and covariants is limited.

And in § 61 it has been stated that Gordan, using the symbolic method of the German investigators, has proved that a complete system of transvectants is coextensive with a complete system of covariants and invariants, and does not comprise an unending series of irreducible forms; thus showing that any binary quantic, or system of binary quantics, has only a finite number of irreducible covariants and invariants.

That the same is true for ternary and higher quantics has been arrived at by Hilbert (*Mathematische Annalen*, Vol. XXXVI) as a consequence of a far-reaching argument as to the finiteness of systems obeying exact laws.

To Hilbert is also due (*Math. Ann.* XXX) the simplest existing proof of Gordan's theorem of the finiteness of the concomitant system for the case of *binary* quantics. This proof will be here exhibited.

150.] **Diophantine Equations.** Some lemmas as to the solutions in positive integers of a system of linear indeterminate, or Diophantine, equations are necessary.

(i) An equation

$$a_1x_1 + a_2x_2 + \dots + a_nx_n = 0,$$

where  $a_1, a_2, \dots, a_n$  are given positive integers, is not satisfied by any set of positive values of  $x_1, x_2, \dots, x_n$ . In fact, the only values, none negative, which satisfy the equation are  $x_1 = x_2 = \dots = x_n = 0$ .

(ii) An equation

$$a_1x_1 + a_2x_2 + \dots + a_nx_n = k,$$

where  $k$ , as well as  $a_1, a_2, \dots, a_n$ , is a positive integer, has, if any, only a finite number of positive integral solutions, zero being counted for the purpose a positive integer. For the whole number of ways in which the number  $k$  can be expressed as a sum of  $n$  or fewer positive integral parts is finite, and the limitation that the parts be integral multiples of some or all of  $a_1, a_2, \dots, a_n$  imposes a further restriction.

(iii) An equation of the form in (i), except that some of the coefficients are positive and some negative integers, is satisfied by an infinite number of positive integral, or some vanishing, values of  $x_1, x_2, \dots, x_n$ . But of this infinite number of positive integral sets of solutions only a finite number are what may be called *simple* sets, i. e. sets which cannot be obtained by adding together other sets of positive integral solutions.

Let the terms with negative coefficients be transposed to the other side of the equation, so that this may be written

$$a_1x_1 + a_2x_2 + \dots + a_mx_m = b_{m+1}x_{m+1} + b_{m+2}x_{m+2} + \dots + b_nx_n,$$

where every  $a$  and every  $b$  is a positive integer.

That there are positive integral solutions is clear. For instance,

$$\begin{aligned} x_1 = b_{m+1}, x_{m+1} = a_1, x_2 = x_3 = \dots = x_m \\ = x_{m+2} = x_{m+3} = \dots = x_n = 0, \end{aligned}$$

$$\text{and } x_1 = b_{m+2}, x_{m+2} = a_1, x_2 = x_3 = \dots = x_{m+1} \\ = x_{m+3} = x_{m+4} = \dots = x_n = 0,$$

are sets of solutions. Moreover, we may take for  $x_1, x_2, \dots, x_n$  any positive integral multiples, or any sums of positive integral multiples of the values of  $x_1, x_2, \dots, x_n$ , in one or

a number, respectively, of these particular sets, and thus obtain another set of solutions. The number of sets is thus infinite. Not all sets are as a rule obtained in this manner, for there will as a rule be other sets in considerable number, for which some or all of  $x_1, x_2, \dots, x_n$  have smaller non-vanishing values, than in sets of solutions comprised in the above aggregate.

We have, however, to establish that the number of *simple* sets of solutions is in all cases finite.

No set of solutions in which  $x_1 > b_{m+1}$ , and  $x_{m+1} > a_1$ , where  $a_1 x_1$  and  $b_{m+1} x_{m+1}$  are any terms on the left and right respectively, can be simple. For any such set of solutions is the sum of the first set of solutions written above and another set.

Thus in a simple set of solutions, if an  $x$  on the left,  $x_1$  say, exceed the greatest coefficient on the right, none of the  $x$ 's on the right can exceed  $a_1$ ; and if an  $x$  on the right,  $x_{m+1}$  say, exceed the greatest coefficient on the left, none of the  $x$ 's on the left can exceed  $b_{m+1}$ .

Simple sets of solutions can then occur only in one or both of two overlapping classes, the class in which no  $x$  on the right exceeds the greatest coefficient on the left, and the class in which no  $x$  on the left exceeds the greatest coefficient on the right.

In the first class we have

$$a_1 x_1 + a_2 x_2 + \dots + a_m x_m \nless (b_{m+1} + b_{m+2} + \dots + b_n) a_1,$$

where  $a_1$  denotes the greatest of  $a_1, a_2, \dots, a_m$ . Now by (ii) the number of sets of positive integral, and vanishing, sets of values of  $x_1, x_2, \dots, x_m$  for which this is the case, is finite; and each set gives for the determination of  $x_{m+1}, x_{m+2}, \dots, x_n$  an equation like

$$k = b_{m+1} x_{m+1} + b_{m+2} x_{m+2} + \dots + b_n x_n,$$

of which the number of sets of solutions is, again by (ii), finite.

And quite similarly in the second class there is only a finite number of sets of solutions.

Of these sets some will be simple; but the vast majority, as a rule, not so.





$x_1, x_2, \dots, x_n$  in the remaining  $r-1$  equations. These become  $r-1$  equations in  $x_{n+1}, x_{n+2}, \dots, x_p$  and  $t_1, t_2, t_3, \dots$ , a finite number of variables whose generality as positive integers or zeros is only limited by the  $r-1$  equations.

The first of these  $r-1$  equations may now be treated exactly as the first of the  $r$  equations was, and substitution may then be made in the remaining  $r-2$  equations; and so on continually till we get to a single equation only. To this the results of (iii) and (iv) apply. Thus we find eventually, on successive substitution backwards, that all the  $p$  variables  $x_1, x_2, \dots, x_p$  can have no more general values than are included in

$$x_1 = A_1\tau_1 + B_1\tau_2 + C_1\tau_3 + \dots,$$

$$x_2 = A_2\tau_1 + B_2\tau_2 + C_2\tau_3 + \dots,$$

$$\dots \dots \dots$$

$$x_p = A_p\tau_1 + B_p\tau_2 + C_p\tau_3 + \dots,$$

where  $A_1, B_1, C_1, \dots, A_2, B_2, C_2, \dots, \dots, A_p, B_p, C_p, \dots$  are definite positive integers, or some of them zeros, and where  $\tau_1, \tau_2, \tau_3, \dots$  are a finite number of arbitraries to which any positive integral and zero values can be assigned at will.

In other words, a system of any number  $r$  of linear Diophantine equations can, if soluble at all in positive integers, have only a finite number of *simple* sets of solutions

$$A_1, A_2, A_3, \dots; B_1, B_2, B_3, \dots; C_1, C_2, C_3, \dots; \&c.,$$

all other sets of positive integral solutions being sums of multiples of some or all of this finite number of sets.

Ex. 1. The only simple set of solutions, excluding the all zero set, of the equations

$$i = y + z = z + x = x + y$$

is the set

$$x = 1, y = 1, z = 1, i = 2.$$

Ex. 2. The simple sets of solutions, besides the all zero set, of

$$i = x + y + w = y + z + u = z + x + v = u + v + w$$

are those of the table

$$\underline{x, y, z, u, v, w, i}$$

$$1, 0, 0, 1, 0, 0, 1$$

$$0, 1, 0, 0, 1, 0, 1$$

$$0, 0, 1, 0, 0, 1, 1$$

Ans. It is easy to reduce the given set to

$$x = u, y = v, z = w, i = u + v + w.$$

Ex. 3. The only not all vanishing simple sets of solutions of

$$4(x+z+u+v) = 2(2z+3u+4v),$$

from which  $z$  disappears, leaving the equation  $2x = u + 2v$ , are given by the table

$x, z, u, v$
0, 1, 0, 0
1, 0, 0, 1
1, 0, 2, 0

This proves that any product of powers of  $a_0, a_2, a_3, a_4$  for which  $4i = 2w$  is a product of powers of  $a_2, a_0a_4, a_0a_3^2$ .

151.] **Application to invariants of one binary quantic.**  
 Now it has been seen in chapter v that if  $a_1, a_2, \dots a_p$  be the roots of the equation in  $x : y$

$$(a_0, a_1, a_2, \dots a_p)(x, y)^p = 0,$$

and if

$$\epsilon = a_0^i (a_1 - a_2)^{n_{12}} (a_1 - a_3)^{n_{13}} (a_2 - a_3)^{n_{23}} \dots$$

be  $a_0^i$  times a product of differences between its roots, such that  $n_{12}, n_{13}, n_{23}, \dots$  are all positive integers or some of them zero, and that all roots occur in the same number  $i$  of factors, so that

$$\begin{aligned} i &= n_{12} + n_{13} + \dots + n_{1p}, \\ &= n_{12} + n_{23} + \dots + n_{2p}, \\ &= n_{13} + n_{23} + \dots + n_{3p}, \\ &= \dots \dots \dots \\ &= n_{1p} + n_{2p} + \dots + n_{p-1, p}, \end{aligned}$$

then  $\Sigma \epsilon$ ,

where the  $\Sigma$  means that the roots are permuted in all possible ways and the sum taken, is, if it do not vanish identically, an invariant of

$$(a_0, a_1, a_2, \dots a_p)(x, y)^p;$$

and, conversely, that any invariant can be expressed as such a sum  $\Sigma \epsilon$ , or at any rate as a sum of numerical multiples of such sums for which  $i$  is the same though the individual  $n$ 's may be different.

We have to prove that any such sum  $\Sigma \epsilon$  can be expressed

rationality and integrally in terms of a finite number of elementary sums of the kind. This will show that any invariant is a rational integral function of a finite number of elementary invariants. It will not show that these elementary invariants are irreducible, but it will that all irreducible invariants occur among them.

It will be observed that the system of equations above in  $i$  and  $n_{12}, n_{13}, n_{23}, \dots$  is a system of Diophantine equations such as that contemplated in § 150 (v). Bringing to bear then on the system the theorem which has been proved, we learn that every such product  $\epsilon$  as above is a product of powers of a finite number of elemental products  $\epsilon_1, \epsilon_2, \epsilon_3, \dots, \epsilon_\mu$  which obey the same laws, one elemental product being given by every simple set of solutions of the system of Diophantine equations, so that generally

$$\epsilon = \epsilon_1^{r_1} \epsilon_2^{r_2} \dots \epsilon_\mu^{r_\mu},$$

for some set of positive integral, including zero, values of the indices  $r_1, r_2, \dots, r_\mu$ .

Every invariant is then of the form

$$\Sigma \epsilon_1^{r_1} \epsilon_2^{r_2} \dots \epsilon_\mu^{r_\mu}$$

for some such set of indices, or a linear function of such sums; and no such sum which does not vanish identically can fail to be an invariant.

The student must bear very clearly in mind the exact meaning of the summation denoted by the  $\Sigma$ . The summation is that of the  $1.2.3 \dots p$  terms, obtained by putting for the roots as they occur in the fully written expression of  $\epsilon_1^{r_1} \epsilon_2^{r_2} \dots \epsilon_\mu^{r_\mu}$  the corresponding roots in every one of the  $1.2.3 \dots p$  permutations of the  $p$  roots  $a_1, a_2, a_3, \dots, a_p$  of the quantic under consideration. The number of terms in the summation is in the first place, and is to be regarded as, the full number  $p!$  of these permutations, though among them in any particular case there may of course, and will, be repetitions or cancellings or both. Other meanings might, but must not, be attached to the  $\Sigma$ . For instance, the meaning might be attached that to each of  $\epsilon_1, \epsilon_2, \dots, \epsilon_\mu$  separately be given every one of its  $p!$  permutational values. We should then get it is true an invariant or an identical zero, but we should have no security that every invariant is thus obtained.

An instance of the need for this caution will occur in an example on the quartic to be given presently.

152.] We now bring in an idea from the theory of equations which will enable us to complete Hilbert's proof of Gordan's theorem.

Consider  $\epsilon_1$ , one of the elemental products  $\epsilon_1, \epsilon_2, \dots, \epsilon_\mu$ . It is one of  $p!$  similar products of which the rest are obtained from it by permuting the  $p$  roots  $a_1, a_2, \dots, a_p$  in all possible ways. These  $p!$  products are the roots of an equation

$$\epsilon_1^{p!} + P_1 \epsilon_1^{p!-1} + P_2 \epsilon_1^{p!-2} + \dots + P_{p!} = 0, \quad \dots (1)$$

of whose coefficients some will frequently vanish identically. Those which do not will be rational integral invariants. This is clear from § 75, but another exhibition of the fact is on the whole preferable here.

By Newton's theorem on the sums of powers of roots (Burnside and Panton's *Theory of Equations*, § 126)  $P_1, P_2, \dots, P_{p!}$  can be expressed rationally and integrally in terms of  $s_1, s_2, \dots, s_{p!}$ , the sums of the first, second, ...  $p!$ th powers of the  $p!$  values of  $\epsilon_1$  which are the roots of the equation. Now  $s_1, s_2, \dots, s_{p!}$ , or rather such of them as do not vanish identically, are invariants exhibited in a form which is a case of the general form of § 151.

Thus the equation (1) expresses  $\epsilon_1^{p!}$  as a linear function of the first  $p!-1$  powers  $\epsilon_1^{p!-1}, \epsilon_1^{p!-2}, \dots, \epsilon_1$  of  $\epsilon_1$  with an absolute term, the coefficients and absolute term being invariants expressible as rational integral functions of  $s_1, s_2, \dots, s_{p!}$ . If we multiply through by  $\epsilon_1$  we obtain an expression for  $\epsilon_1^{p!+1}$ , which upon insertion in it of the already obtained expression for  $\epsilon_1^{p!}$ , becomes a linear function of  $\epsilon_1^{p!-1}, \epsilon_1^{p!-2}, \dots, \epsilon_1$ , whose coefficients and absolute term are rational integral functions of  $s_1, s_2, \dots, s_{p!}$ . Multiply again by  $\epsilon_1$ , and again replace  $\epsilon_1^{p!}$  by the expression for it; and repeat the same process any number of times that may be desired. We thus obtain the fact that, the index  $r_1$  being any number not less than  $p!$ ,

$$\epsilon_1^{r_1} = Q_1 \epsilon_1^{p!-1} + Q_2 \epsilon_1^{p!-2} + \dots + Q_{p!},$$

where  $Q_1, Q_2, \dots, Q_{p!}$  are invariants capable of expression as rational integral functions of the  $p!$  invariants and zeros  $s_1, s_2, \dots, s_{p!}$ , i. e.  $\Sigma \epsilon_1, \Sigma \epsilon_1^2, \dots, \Sigma \epsilon_1^{p!}$ .

Proceed now in like manner with  $\epsilon_2$ , a second of the elemental products  $\epsilon_1, \epsilon_2, \dots, \epsilon_\mu$ . This again is a root of an equation of degree  $p!$  whose first coefficient is unity and whose other non-vanishing coefficients are invariants expressible rationally and integrally in terms of  $p!$  sums of which those which do not vanish are invariants, viz.  $\Sigma \epsilon_2, \Sigma \epsilon_2^2, \dots, \Sigma \epsilon_2^{p!}$ . Consequently, if  $r_2$  be not less than  $p!$ , we have

$$\epsilon_2^{r_2} = R_1 \epsilon_2^{p!-1} + R_2 \epsilon_2^{p!-2} + \dots + R_{p!},$$

where  $R_1, R_2, \dots, R_{p!}$  are invariants expressible as rational integral functions of  $\Sigma \epsilon_2, \Sigma \epsilon_2^2, \dots, \Sigma \epsilon_2^{p!}$ .

In like manner we have like expressions for powers not less than the  $p!$ th of  $\epsilon_3, \epsilon_4, \dots, \epsilon_\mu$ , the remaining elemental products of powers of  $a_0$  and differences between roots.

153.] The number of irreducible invariants of a binary quantic is finite. We now return to the general expression of § 151

$$\Sigma \epsilon_1^{r_1} \epsilon_2^{r_2} \dots \epsilon_\mu^{r_\mu}.$$

There is, in the first place, only a finite number of these expressions for which none of the exponents  $r_1, r_2, \dots, r_\mu$  exceeds  $p! - 1$ ; viz.  $(p!)^\mu - 1$ .

Take, however, any one in which one or more of  $r_1, r_2, \dots, r_\mu$  exceeds  $p! - 1$ , and express such higher  $r$ th powers of elemental factors in terms under the  $\Sigma$  by the expressions obtained in the last article in terms of powers less than the  $p!$ th. Having done so, multiply out the expression obtained. The result is an identity like

$$\begin{aligned} \Sigma \epsilon_1^{r_1} \epsilon_2^{r_2} \dots \epsilon_\mu^{r_\mu} &= \Sigma \{ K_1 \epsilon_1^{\rho_1} \epsilon_2^{\rho_2} \dots \epsilon_\mu^{\rho_\mu} + K_2 \epsilon_1^{\sigma_1} \epsilon_2^{\sigma_2} \dots \epsilon_\mu^{\sigma_\mu} + \dots \} \\ &= K_1 \Sigma \epsilon_1^{\rho_1} \epsilon_2^{\rho_2} \dots \epsilon_\mu^{\rho_\mu} + K_2 \Sigma \epsilon_1^{\sigma_1} \epsilon_2^{\sigma_2} \dots \epsilon_\mu^{\sigma_\mu} + \dots, \end{aligned}$$

where none of the indices on the right exceeds  $p! - 1$ , and where  $K_1, K_2, \dots$  are rational integral functions of the  $\mu$  times  $p!$  sums

$$\Sigma \epsilon_1, \Sigma \epsilon_1^2, \dots, \Sigma \epsilon_1^{p!},$$

$$\Sigma \epsilon_2, \Sigma \epsilon_2^2, \dots, \Sigma \epsilon_2^{p!},$$

$$\dots \dots \dots$$

$$\Sigma \epsilon_\mu, \Sigma \epsilon_\mu^2, \dots, \Sigma \epsilon_\mu^{p!},$$

which are all rational integral invariants, and are themselves

all included in the form  $\Sigma \epsilon_1^{r_1} \epsilon_2^{r_2} \dots \epsilon_\mu^{r_\mu}$  for values of the indices not exceeding  $p!$ .

Thus every sum  $\Sigma \epsilon_1^{r_1} \epsilon_2^{r_2} \dots \epsilon_\mu^{r_\mu}$ , and therefore (§ 151) every rational integral invariant, is a rational integral function of a finite number of rational integral invariants; viz. of those included in the form

$$\Sigma \epsilon_1^{r_1} \epsilon_2^{r_2} \dots \epsilon_\mu^{r_\mu}$$

for values of the indices none of which exceeds  $p!$ , and which are indeed all less than  $p!$  except that one of them may be equal to  $p!$  when all the rest are zero.

It is well to repeat that the summation is to be taken as including  $p!$  terms, one corresponding to every permutation of the  $p$  roots.

As already pointed out in § 151, it must not be supposed that we have here the exact number of irreducible invariants, of which all other invariants are rational integral functions, or the forms of invariants which are irreducible. The number of the invariants in terms of which all invariants are here shown to be capable of rational integral expression is, for quantics of low orders to which Gordan's method of transvectants and the arithmetical method of the last chapter have been applied, vastly in excess of the necessities. Moreover, no precise number is really assigned at all by the above reasoning, for even when the elemental products  $\epsilon_1, \epsilon_2, \dots, \epsilon_\mu$  are known for any quantic we have still no information as to how many of the sums  $\Sigma \epsilon_1^{r_1} \epsilon_2^{r_2} \dots \epsilon_\mu^{r_\mu}$  vanish identically.

But a finite number of expressions has been definitely specified, all non-vanishing individuals among which are invariants, and in terms of which all other invariants can be rationally and integrally expressed. That some of these only are strictly irreducible invariants, while the rest are rational integral functions of them, does not affect the argument that all invariants are rational integral functions of a finite irreducible system. A selection from a finite system is itself finite.

A modification of Hilbert's method has been proposed by Kempe. He succeeds in using a more readily exhibited system of products of differences in place of the elemental products contemplated in this chapter. See his paper 'On Regular Difference Terms,' *Proc. Lond. Math. Soc.* Vol. XXV. p. 343.

154.] **The cubic and quartic.** Little is taught us as to the invariants of particular quantities by exhibiting the method in their particular cases, but for the light thrown on the method itself we exemplify it in the cases of the cubic and quartic.

For the cubic  $(a, b, c, d) (x, y)^3$ , whose roots are  $a, \beta, \gamma$ , any invariant is a numerical multiple of

$$\Sigma a^i (\beta - \gamma)^r (\gamma - a)^s (a - \beta)^t,$$

where  $i = s + t = t + r = r + s$ . Of these equations (§ 150, Ex. 1) the only simple set of solutions is  $r = s = t = 1, i = 2$ . Thus the only elemental product is

$$\epsilon = a^2 (\beta - \gamma) (\gamma - a) (a - \beta).$$

The  $3!$  permutations of  $a, \beta, \gamma$  in this give three products each equal to  $\epsilon$  and three equal to  $-\epsilon$ . Thus the equation with invariant coefficients satisfied by  $\epsilon$  is

$$\{\epsilon^2 - a^4 (\beta - \gamma)^2 (\gamma - a)^2 (a - \beta)^2\}^3 = 0.$$

All irreducible invariants are then included among

$$\Sigma \epsilon, \Sigma \epsilon^2, \Sigma \epsilon^3, \Sigma \epsilon^4, \Sigma \epsilon^5, \Sigma \epsilon^6,$$

of which the first, third and fifth vanish, while the second, fourth and sixth are

$$6\epsilon^2, 6\epsilon^4, 6\epsilon^6.$$

Of these the second and third are numerical multiples of powers of the first. Thus  $6\epsilon^2$ , or, say,

$$\epsilon^2 = a^4 (\beta - \gamma)^2 (\gamma - a)^2 (a - \beta)^2,$$

is the only irreducible invariant of the cubic.

For the quartic  $(a, b, c, d, e) (x, y)^4$ , whose roots are  $a, \beta, \gamma, \delta$ , invariants have the form

$$\Sigma a^i (\beta - \gamma)^r (\gamma - a)^s (a - \beta)^t (a - \delta)^{r'} (\beta - \delta)^{s'} (\gamma - \delta)^{t'},$$

where  $i = s + t + r' = t + r + s' = r + s + t' = r' + s' + t'$ ,

of which the simple sets of solutions are given in § 150, Ex. 2, and tell us that the elemental products are

$$\epsilon_1 = a (\beta - \gamma) (a - \delta),$$

$$\epsilon_2 = a (\gamma - a) (\beta - \delta),$$

$$\epsilon_3 = a (a - \beta) (\gamma - \delta),$$



These are, it will be noticed, as is *not* the case in general for the  $\epsilon$ 's corresponding to higher quantities, of one type, and are the  $au, av, aw$  of § 80.

The  $4!$  permutational values of  $\epsilon_1$  are

$$au, av, aw, -au, -av, -aw$$

each four times repeated; and the values of  $\epsilon_2$  and  $\epsilon_3$  are the same in different orders.

The equation with invariant coefficients whose roots are the 24 values of either  $\epsilon_1$  or  $\epsilon_2$  or  $\epsilon_3$  is (§ 81, Ex. 4)

$$\{(\epsilon^3 - 12I\epsilon + a^3uvw)(\epsilon^3 - 12I\epsilon - a^3uvw)\}^4 = 0. \quad \dots(1)$$

The irreducible invariants are included in the finite number like

$$\Sigma \epsilon_1^\lambda \epsilon_2^\mu \epsilon_3^\nu,$$

where neither of  $\lambda, \mu, \nu$  exceeds 24, and where if either one is 24 the other two vanish.

This example, then, affords an instance of the great excess over the number of irreducible invariants of the finite number of invariants among which they are included according to the present investigation. For we know from previous chapters that the only really irreducible invariants are

$$\Sigma \epsilon_1^2 = \Sigma \epsilon_2^2 = \Sigma \epsilon_3^2 = 8a^2(u^2 + v^2 + w^2)$$

and

$$\begin{aligned} \Sigma \epsilon_1^2 \epsilon_2 = \Sigma \epsilon_2^2 \epsilon_3 = \Sigma \epsilon_3^2 \epsilon_1 = -\Sigma \epsilon_1 \epsilon_2^2 = -\Sigma \epsilon_2 \epsilon_3^2 = -\Sigma \epsilon_3 \epsilon_1^2 \\ = 4a^3\{u^2(v-w) + v^2(w-u) + w^2(u-v)\}. \end{aligned}$$

It also affords an illustration of the care which must be taken to attach the right meaning (see § 151) to the summation  $\Sigma$ . If in  $\Sigma \epsilon_1^2 \epsilon_2$ , instead of taking this as meaning the sum of 24 terms obtained by permuting the roots as they occur in  $\epsilon_1^2 \epsilon_2$  in all possible ways, we had wrongly taken it as meaning that  $\epsilon_1$  is to be given its 24 (or its 6 essentially different) values, and  $\epsilon_2$  similarly, we should have had a sum of  $24^2$  (or  $6^2$ ) terms which vanishes identically, and should not have obtained the, in fact irreducible, invariant

$$a^3\{u^2(v-w) + v^2(w-u) + w^2(u-v)\}$$

at all. Or, if we had taken it as meaning the sum of the 6.5 terms  $\epsilon^2 \epsilon'$ , where  $\epsilon$  and  $\epsilon'$  are two different roots of the

equation (1) for  $\epsilon_1$  or for  $\epsilon_2$ , the same failure would have resulted.

155.] To secure clearness we have in the last four articles restricted our field of investigation as much as possible, and have confined attention to *invariants*, and to invariants of a single binary quantic only. Neither Gordan's theorem, however, nor Hilbert's line of argument is of such restricted application.

Equal fulness of treatment is unnecessary in the next two articles, which deal respectively with the case of covariants of a single binary quantic and the general case of covariants and invariants of more binary quantics than one.

156.] The number of irreducible covariants and invariants of a binary quantic is finite. Let us use the word covariant as including invariant as a particular case, and also as including the quantic itself.

Any covariant of the binary  $p$ -ic

$$(a_0, a_1, a_2, \dots a_p)(x, y)^p$$

whose roots are  $a_1, a_2, a_3, \dots a_p$  is, by chapter v, of the form

$$\Sigma . a_0^i (x - a_1 y)^{m_1} (x - a_2 y)^{m_2} \dots (x - a_p y)^{m_p} .$$

$$(a_1 - a_2)^{n_{12}} (a_1 - a_3)^{n_{13}} (a_2 - a_3)^{n_{23}} \dots,$$

where the positive integers, or some of them zeros,

$$m_1, m_2, \dots m_p, n_{12}, n_{13}, n_{23}, \dots$$

satisfy the Diophantine equations

$$\begin{aligned} \omega &= m_1 + m_2 + m_3 + \dots + m_p, \\ i &= m_1 + n_{12} + n_{13} + \dots + n_{1p} \\ &= m_2 + n_{12} + n_{23} + \dots + n_{2p} \\ &= m_3 + n_{13} + n_{23} + \dots + n_{3p} \\ &= \dots \dots \dots \dots \dots \dots \\ &= m_p + n_{1p} + n_{2p} + \dots + n_{p-1, p}, \end{aligned}$$

or is a linear function of such sums for the same values of  $i$  and  $\omega$ ; and all sums of this form are covariants. In particular, those for which the  $m$ 's are all zero are invariants, and those for which the  $n$ 's are all zero are powers of the  $p$ -ic itself.

Now by § 150 (v) the number of simple sets of solutions of these Diophantine equations for  $i$ ,  $\omega$  and the  $m$ 's and  $n$ 's is finite. Every product, such as that under the  $\Sigma$  above, is then a product of powers of elemental products of the same form and with the same properties.

If these elemental products be called  $\eta_1, \eta_2, \dots, \eta_\mu$ , then, exactly as in § 152, these severally are roots of equations of degree  $p!$ , in each of which the coefficient of the first term is unity, and the other coefficients are covariants which are rational integral functions, in the first case of  $\Sigma\eta_1, \Sigma\eta_1^2, \dots, \Sigma\eta_1^{p!}$ , in the second case of  $\Sigma\eta_2, \Sigma\eta_2^2, \dots, \Sigma\eta_2^{p!}$ , &c., &c. Hence, as in § 153, all covariants can be expressed rationally and integrally in terms of covariants included in the limited class

$$\Sigma\eta_1^{\nu_1}\eta_2^{\nu_2}\dots\eta_\mu^{\nu_\mu},$$

where neither of the indices exceeds  $p!$ , and none is in fact so great as  $p!$  unless all the others vanish.

Remark that, the number of irreducible covariants (including invariants) being finite, the number of irreducible seminvariants (including invariants) is also finite. For, if any covariant is a rational integral function of other covariants, the coefficient of the highest power of  $x$  in it is that same rational integral function of those of them which are free from  $x$  (i.e. the invariants among them) and the coefficients of the highest powers of  $x$  in the rest.

Ex. 4. In the case of the cubic the elemental products are

$$a(x-ay)(x-\beta y)(x-\gamma y),$$

$$a^2(\beta-\gamma)(\gamma-a)(a-\beta),$$

and  $a(x-ay)(\beta-\gamma)$ ,  $a(x-\beta y)(\gamma-a)$ ,  $a(x-\gamma y)(a-\beta)$ .

157.] **Several binary quantics.** The proof that all covariants and invariants of a finite number of binary quantics are rational integral functions of a finite number of covariants and invariants of the system is similar.

For a system of a finite number of binary quantics whose leading coefficients are  $a_0, a_0', \dots$ , and whose roots are  $a_1, a_2, \dots, a_p$ , in the case of the first,  $a_1', a_2', \dots, a_p'$ , in the case of the second, and so on, the general expression for covariants

of which all other covariants are linear functions of one degree and order is of the form

$$\Sigma \{ a_0^i a_0'^{i'} \dots \Pi(x - ay) \cdot \Pi(x - a'y) \dots \\ \Pi(a_r - a_s) \cdot \Pi(a_r' - a_s') \dots \Pi(a_r - a_s') \dots \},$$

where  $\Pi(x - ay)$  denotes a product of a number of the differences  $x - a_1y, x - a_2y, \dots, x - a_py$  and their powers,  $\Pi(x - a'y)$  a product of a number of the differences  $x - a_1'y, x - a_2'y, \dots, x - a_p'y$  and their powers,  $\Pi(a_r - a_s), \Pi(a_r' - a_s')$ , &c., products of numbers of differences between two roots of the first, second, &c., quantities and powers of such differences, and  $\Pi(a_r - a_s')$ , &c., products of differences and powers of differences between roots belonging to different quantities of the system. Also, in a product under the  $\Sigma$ , all roots  $a_1, a_2, \dots, a_p$  of the first quantic occur in the same number  $i$  of factors, all roots  $a_1', a_2', \dots, a_p'$  of the second quantic occur in the same number  $i'$  of factors, and so on. The summation  $\Sigma$  consists of  $p! p'! \dots$  terms at most, obtained by permuting the  $p$  roots  $a_1, a_2, \dots, a_p$ , the  $p'$  roots  $a_1', a_2', \dots, a_p'$ , &c., in all possible ways.

The conditions as to degrees in the various  $a$ 's,  $a'$ 's, &c., are expressed by  $p + p' + \dots$  linear Diophantine equations in  $i, i', \dots$  and the exponents of powers of differences; and  $\varpi$ , the order in  $x, y$ , is determined by another sum of the exponents. Now the whole system of  $p + p' + \dots + 1$  equations in  $i, i', \dots, \varpi$  and the exponents has, by § 150 (v), only a finite number of simple sets of solutions. To each simple set of solutions corresponds an elemental product. If these elemental products be  $\omega_1, \omega_2, \omega_3, \dots$ , the general  $\Sigma$  above is, as before, capable of expression in the form  $\Sigma \omega_1^{r_1} \omega_2^{r_2} \omega_3^{r_3} \dots$ .

Also, precisely as before, every elemental product  $\omega_1$  satisfies an equation of finite degree, in no case exceeding  $p! p'! \dots$ , whose coefficients, after the first which is unity, are rational integral functions of a finite number of sums of powers of  $\omega_1$  and the results of permuting among themselves the various roots in  $\omega_1$ . Hence, as in earlier cases, the sum

$$\Sigma \omega_1^{r_1} \omega_2^{r_2} \omega_3^{r_3} \dots$$

can be expressed rationally and integrally in terms of the finite number of like sums in which  $r_1, r_2, r_3, \dots$  do not exceed definite numbers; and these like sums are all rational integral

covariants and invariants. All rational integral covariants and invariants of a system of binary quantics are then rational integral functions of a finite number of such concomitants of the system.

In particular, all invariants of the system are rational integral functions of a finite number of invariants. For if an invariant, free from the variables, be a rational integral function of invariants and covariants, we obtain, upon putting the variables equal to zero in the identity of the invariant and the function, an identity in which all the covariants are replaced by zeros while the invariants alone remain.

## CHAPTER X.

### PROTOMORPHS, ETC.

158.] **Recapitulation.** We have seen in chapter vi that the leading coefficient, that of the highest power of  $x$ , in any covariant of

$$u \equiv (a_0, a_1, a_2, \dots a_p)(x, y)^p$$

is a seminvariant, i.e. is annihilated by

$$\Omega \equiv a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + \dots + pa_{p-1} \frac{d}{da_p},$$

and consequently possesses the half invariant property of being invariantic for such linear substitutions as

$$x = X + mY, \quad y = Y.$$

It has also been seen that conversely

$$S(a_0, a_1, a_2, \dots a_p),$$

any gradient of extent not exceeding  $p$  which is such as to satisfy the differential equation

$$\Omega S = 0,$$

and which is consequently a seminvariant, is the leading coefficient of a covariant whose order  $\varpi$  in  $x, y$  is given in terms of  $p$  and  $i$  and  $w$ , the degree and weight of  $S$ , by the relation

$$\varpi = ip - 2w;$$

in fact, that the covariant in question may be written

$$x^{ip-2w} e^{\frac{y}{x} O} S(a_0, a_1, a_2, \dots a_p),$$

where

$$O \equiv pa_1 \frac{d}{da_0} + (p-1)a_2 \frac{d}{da_1} + \dots + a_p \frac{d}{da_{p-1}}.$$

If  $\omega = 0$ , the seminvariant is an invariant.

Thus covariants, including invariants as a particular case, and seminvariants, also including invariants as a particular case, are equally numerous, and correspond one to another. If any relation or syzygy connects certain covariants, the same syzygy connects their seminvariant leaders, and vice versa.

Another expression for the same covariant, whose leader is the seminvariant  $S(a_0, a_1, a_2, \dots a_p)$ , is

$$(-1)^w y^{i^{p-2}w} e^{\frac{x}{y}\Omega} S(a_p, a_{p-1}, a_{p-2}, \dots a_0),$$

where  $S(a_p, a_{p-1}, a_{p-2}, \dots a_0)$  is the anti-seminvariant obtained by interchanging  $a_0$  and  $a_p$ ,  $a_1$  and  $a_{p-1}$ , &c. in the seminvariant  $S(a_0, a_1, a_2, \dots a_p)$ .

159.] **Elimination of  $x$  between  $u$  and its successive derivatives with regard to  $x$ .** Two interesting conclusions at once follow from the results of chapter vi.

Of these the first is that if in any seminvariant

$$S(a_0, a_1, a_2, \dots a_p)$$

of  $u$  we replace

$a_p$  by  $u$ , i.e. by  $(a_0, a_1, a_2, \dots a_p)(x, y)^p$ , which call  $a_p$ ,

$a_{p-1}$  by  $\frac{1}{p} \frac{du}{dx}$ , i.e. by  $(a_0, a_1, a_2, \dots a_{p-1})(x, y)^{p-1}$ ,  
 which call  $a_{p-1}$ ,

. . . . .

$a_2$  by  $\frac{1}{p(p-1) \dots 3} \frac{d^{p-2}u}{dx^{p-2}}$ , i.e. by  $a_0x^2 + 2a_1xy + a_2y^2$ ,  
 which call  $a_2$ ,

$a_1$  by  $\frac{1}{p(p-1) \dots 3 \cdot 2} \frac{d^{p-1}u}{dx^{p-1}}$ , i.e. by  $a_0x + a_1y$ ,  
 which call  $a_1$ ,

$a_0$  by  $\frac{1}{p(p-1) \dots 3 \cdot 2 \cdot 1} \frac{d^p u}{dx^p}$ , i.e. by  $a_0$ , which call  $a_0$ ,

no  $x$  appears in the result, which is merely the seminvariant itself multiplied by  $y^w$ , where  $w$  is its weight.

The result of substitution is annihilated by

$$a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + 3a_2 \frac{d}{da_3} + \dots + pa_{p-1} \frac{d}{da_p}.$$

Now, writing  $u_r$  for  $\frac{d^r u}{dx^r}$ , for every value of  $r$  from 1 to  $p$  inclusive, we see that this is

$$u_p \frac{d}{du_{p-1}} + u_{p-1} \frac{d}{du_{p-2}} + u_{p-2} \frac{d}{du_{p-3}} + \dots + u_1 \frac{d}{du},$$

which, when it operates on a function of  $u, u_1, u_2, \dots, u_p$ , is merely  $\frac{d}{dx}$ . The annihilation tells us then that the result of substitution is free from  $x$ . What does not vanish when the substitution is made in a seminvariant is clearly  $y^w$  into that seminvariant, for weight is degree in  $x, y$  in the result, and the form of result independent of  $x$  is the form we should get by putting  $x = 0$  first and then making the substitutions.

The intimate connexion of the theory of seminvariants with that of elimination between  $u$  and its  $x$ -derivatives is now apparent. For instance, from § 137 we conclude that, if  $u$  be any rational integral function of order  $p$  in  $x$ , the number of linearly independent rational integral functions of  $u$  and its successive derivatives with regard to  $x$ , which are of degree  $i$  in the coefficients and free from  $x$ , is  $\left(\frac{ip}{2}; i, p\right)$  or  $\left(\frac{ip-1}{2}; i, p\right)$ , according as  $ip$  is even or odd. And again, from known facts as to numbers of irreducible seminvariants of binary quantics of the first few orders we obtain that the numbers of rational integral functions, of rational integral expressions of orders 1, 2, 3, 4, 5 respectively in  $x$  and their successive derivatives with regard to  $x$ , which are free from  $x$  and none of which, in the five cases respectively, can be rationally and integrally expressed in terms of the rest, are, in the five cases respectively, 1, 2, 4, 5, 23.

Once more, from § 128 or from § 100, we gather that any rational integral function of  $u$ , a  $p$ -ic in  $x$ , and its derivatives, which is throughout of degree  $i$  in  $u$  and the derivatives, and in every term of which the sum of the indices  $r$  for factors like  $\frac{d^r u}{dx^r}$  is constant and equal to  $ip - w$ , can be written as the derivative with regard to  $x$  of a rational integral function of  $u$  and its successive derivatives if  $ip - 2w > 0$ .



Ex. 1. Integrate the differential equation

$$u \frac{d^4 u}{dx^4} - \frac{du}{dx} \frac{d^3 u}{dx^3} + \frac{1}{2} \left( \frac{d^2 u}{dx^2} \right)^2 = 0.$$

Ans.  $u = ax^4 + 4bx^3 + 6cx^2 + 4dx + e$ , where  $ae - 4bd + 3c^2 = 0$ .

Ex. 2. If  $u = ax^3 + 3bx^2 + 3cx + d$ , express  $\int u \left( \frac{d^2 u}{dx^2} \right)^4 dx$  rationally and integrally in terms of  $u, \frac{du}{dx}, \frac{d^2 u}{dx^2}, \frac{d^3 u}{dx^3}$ .

Ans. Here  $ip - 2w = 1$ . Assume the most general form and determine the arbitrary coefficients by operating with  $\frac{d}{dx}$ .

160.] Covariants obtained by substitution of  $x$ -derivatives. The second important conclusion referred to at the outset of the preceding article appears to be due to Faa de Bruno (*Am. J.* III). It is that if we make the same substitutions as in the preceding article for  $a_0, a_1, a_2, \dots, a_p$  in

$$S(a_p, a_{p-1}, a_{p-2}, \dots, a_0),$$

the anti-seminvariant obtained by interchanging  $a_0$  and  $a_p, a_1$  and  $a_{p-1}, \&c.$ , in a seminvariant  $S(a_0, a_1, a_2, \dots, a_p)$  of weight  $w$  and degree  $i$ , the result obtained will be the covariant of which that seminvariant is the leader multiplied by  $(-y)^w$ .

Noticing that the substitution of  $a_r$  for  $a_r$  is that of  $y^r A'_r$ , where  $A'_r$  is the  $A_r$  of § 92 with  $\frac{x}{y}$  put for  $m$ , we see by § 93, bearing in mind the isobarism of weight  $ip - w$  of

$$S(a_p, a_{p-1}, a_{p-2}, \dots, a_0),$$

that the result of the substitution is

$$y^{ip-w} e^{\frac{x}{y} \Omega} S(a_p, a_{p-1}, a_{p-2}, \dots, a_0).$$

Now by § 109, since  $(-1)^w S(a_p, a_{p-1}, \dots, a_0)$  is the last coefficient in the covariant which  $S(a_0, a_1, a_2, \dots, a_p)$  leads, the covariant going with  $S$  is

$$(-1)^w y^{ip-2w} e^{\frac{x}{y} \Omega} S(a_p, a_{p-1}, a_{p-2}, \dots, a_0).$$

Of these two expressions the former is  $(-y)^w$  times the latter.

Thus from any seminvariant a mere substitution derives the corresponding covariant.

If the seminvariant be an invariant, the substitution has the effect only of multiplying it by  $(-1)^w y^{ip-w}$ , i.e. by  $(-y)^w$ , since  $ip-2w=0$  for an invariant. This accords with the result of § 159.

161.] A seminvariant is given by certain of its terms. In the substitution of § 159 put  $-\frac{a_1}{a_0}$  for  $x$ , and 1 for  $y$ , i.e. for  $a_0, a_1, a_2, \dots a_p$  substitute

$$a_0' \equiv a_0,$$

$$a_1' \equiv 0,$$

$$a_2' \equiv a_2 - \frac{a_1^2}{a_0},$$

$$a_3' \equiv a_3 - 3 \frac{a_1}{a_0} a_2 + 2 \frac{a_1^3}{a_0^2},$$

$$a_4' \equiv a_4 - 4 \frac{a_1}{a_0} a_3 + 6 \frac{a_1^2}{a_0^2} a_2 - 3 \frac{a_1^4}{a_0^3},$$

. . . . .

$$a_p' \equiv (a_p, a_{p-1}, a_{p-2}, \dots a_0) \left(1, -\frac{a_1}{a_0}\right)^p.$$

The result of § 159 tells us that

$$S(a_0, a_1, a_2, a_3, \dots a_p) = S(a_0', 0, a_2', a_3', \dots a_p'),$$

where  $S$  denotes any seminvariant, or, in particular, invariant.

Thus all rational integral seminvariants are rational integral functions of the  $p$  expressions  $a_0'$  (i.e.  $a_0$ ),  $a_2'$ ,  $a_3'$ ,  $\dots a_p'$ . These expressions are all integral in  $a_1, a_2, \dots a_p$ , but, after the first, are fractional in  $a_0$ . They are seminvariants, fractional after the first, for it is easy to see that they are all annihilated by  $\Omega$ . This will appear in another light presently.

It follows that if we know the terms free from  $a_1$  in any seminvariant or invariant, we know the whole expression of that seminvariant or invariant. To find this whole expression we have merely to write the values of  $a_2', a_3', \dots a_p'$  instead of  $a_2, a_3, \dots a_p$  in the given terms. We shall see presently that this substitution may be effected by differential operation.

We shall also notice later a means of obtaining the terms free from  $a_1$  in *invariants*.

The search for rational integral seminvariants and invariants may be regarded as the search for rational integral homogeneous isobaric functions of  $a_0, a_2', a_3', \dots a_p'$ , which, when the full values of  $a_2', a_3', \dots a_p'$  are substituted in them, are integral in  $a_0$ .

Ex. 3. Given  $a^2d^2 + 4ac^3$ , the terms free from  $b$  in the discriminant of the cubic  $(a, b, c, d)(x, y)^3$  (cf. chap. vi, Ex. 35), obtain the full expression for the discriminant.

$$\begin{aligned} \text{Ex. 4. Verify that } a_0a_2'a_4' - a_0a_3'^2 - a_2'^3 \\ = a_0a_2a_4 + 2a_1a_2a_3 - a_0a_3^2 - a_1^2a_4 - a_2^3. \end{aligned}$$

Ex. 5. No seminvariant has  $a_1$  for a factor.

Ex. 6. The number of linearly independent seminvariants of type  $w, i$  of a  $p$ -ic, whose terms free from  $a_1$  are integral, and whose other terms are integral in  $a_1, a_2, a_3, \dots a_p$  though not necessarily in  $a_0$ , is

$$(w; i, p) - (w-1; i-1, p).$$

Ans. Since  $(w-1; i-1, p)$  is the number of products of type  $w, i$  which involve  $a_1$ , this difference is the number of products of the type of  $a_0, a_2', a_3', \dots a_p'$ .

162.] **Coefficients of quantic deprived of its second term are seminvariants.** If we inspect the expressions for

$$a_0', a_2', a_3', \dots a_p'$$

in the last article, we notice that they are the coefficients of  $X^p, X^{p-2}Y^2, X^{p-3}Y^3, \dots Y^p$  in the result of depriving the  $p$ -ic

$$(a_0, a_1, a_2, \dots a_p)(x, y)^p$$

of its second term by the substitution

$$x = X - \frac{a_1}{a_0} Y, \quad y = Y,$$

of which the modulus is unity.

Now this substitution is one after which a seminvariant as well as an invariant persists in form, for it has the effect of altering all roots by the same addition  $\frac{a_1}{a_0}$ , and leaves the leading coefficient  $a_0$  or  $a_0'$  unaltered.

We thus see clearly the meaning of the identity of the last article

$$S(a_0, a_1, a_2, a_3, \dots a_p) = S(a_0, 0, a_2', a_3', \dots a_p').$$



Let  $S$  be as usual of weight  $w$  and degree  $i$ . Notice that every non-vanishing argument on the right involves  $a_0$  explicitly to the degree  $1-r$ , where  $r$  is its weight. Every product of  $i$  arguments on the right involves then  $a_0$  to the power  $i-w$ , where  $w$  is its whole weight. Consequently, upon multiplying through by  $a_0^{w-i}$  we obtain that

$$a_0^{w-i}S(a_0, a_1, a_2, a_3, \dots a_p) = S(1, 0, A_2, A_3, \dots A_p),$$

which is a rational integral function of  $A_2, A_3, \dots A_p$  only.

We have here the completion of the theorem of § 42. It was there proved that a binary  $p$ -ic cannot have more than  $p$  algebraically independent covariants and invariants including itself, but that if  $p-1$  distinct from the  $p$ -ic itself and independent of it and one another can be found, then all others must be capable of expression in terms of it and them. We have now proved that there are certainly  $p-1$  covariants distinct from the  $p$ -ic, namely, those whose leading coefficients are the seminvariants  $A_2, A_3, \dots A_p$ . These are certainly independent of one another and the  $p$ -ic; for in the series  $a_0, A_2, A_3, \dots A_p$  each involves one of the coefficients  $a_0, a_2, a_3, \dots a_p$  which is absent from all those which precede it. We have also shown that the expression for any rational integral seminvariant or invariant  $S(a_0, a_1, a_2, \dots a_p)$  in terms of  $a_0, A_2, A_3, \dots A_p$  is rational, and is integral in all but the first  $a_0$ , which it involves only in the form of the factor  $a_0^{-w+i}$ .

The expression of the covariant (or invariant) whose leading coefficient is  $S$  in terms of those whose leading coefficients are  $a_0, A_2, A_3, \dots A_p$ , of which the first is the  $p$ -ic itself, follows. The covariant whose leading coefficient is a seminvariant is unique (cf. §§ 111, 112). Now, if  $u$  be the  $p$ -ic, and  $K$  the covariant whose leading coefficient is  $S(a_0, a_1, a_2, \dots a_p)$ , the covariant whose leading coefficient is  $a_0^{w-i}S(a_0, a_1, a_2, \dots a_p)$  is

$$u^{w-i}K.$$

Also, if  $a_2, a_3, \dots a_p$  be the covariants whose leading coefficients are  $A_2, A_3, \dots A_p$ ,

$$S(1, 0, a_2, a_3, \dots a_p)$$

is a covariant. For it is a rational integral function of covariants, and is of constant degree and weight throughout,

and therefore of constant order in  $x, y$ , since the degree and weight of  $A_r$ , the leading coefficient of  $a_r$ , are both  $r$ , and consequently the order  $ip - 2w$  of  $a_r$  is  $r(p-2)$  so that the order of any product of  $a$ 's which occurs is  $w(p-2)$  where  $w$  is the constant weight. This covariant is the one whose leading coefficient is  $S(1, 0, A_2, A_3, \dots, A_p)$ . Hence the identity of seminvariants

$$a_0^{w-i} S(a_0, a_1, a_2, a_3, \dots, a_p) = S(1, 0, A_2, A_3, \dots, A_p)$$

necessitates the identity of covariants

$$u^{w-i} K = S(1, 0, a_2, a_3, \dots, a_p).$$

In other words, any covariant or invariant can be expressed as the result of dividing a rational integral function of  $a_2, a_3, \dots, a_p$  by the power  $u^{w-i}$  of  $u$ , where  $w$  and  $i$  are the weight and degree of the leading coefficient of the covariant in question.

In particular, any invariant  $I(a_0, a_1, a_2, \dots, a_p)$  is equivalent to

$$u^{-\frac{1}{2}i(p-2)} I(1, 0, a_2, a_3, \dots, a_p),$$

for, in the case when  $K$  is an invariant,  $ip = 2w$ .

We have here reasoned for cases when  $w \leq i$ . The student can supply the slight modification of reasoning necessary when  $w < i$ . In such a case the result is best written

$$K = u^{i-w} S(1, 0, a_2, a_3, \dots, a_p).$$

Any covariant, then, whose leading coefficient is of smaller weight than degree, has a power of the quantic for a factor.

164.] **A complete system of protomorphs is not unique.** The seminvariants  $a_0, A_2, A_3, \dots, A_p$ , or

$$a_0, a_0 a_2', a_0^2 a_3', \dots, a_0^{p-1} a_p',$$

do not stand alone among rational integral seminvariants as being a set of  $p$  in terms of which all others can be expressed rationally and integrally but for a power of the first, which when  $w > i$  is a negative power, as factor.

A system of  $p$  seminvariants possessing this property is called a set of *protomorphic seminvariants* or *protomorphs*.

We proceed to see that an allowable system of protomorphs

is composed of  $a_0$  and any set of  $p-1$  rational integral seminvariants  $B_2, B_3, \dots B_p$ , which are such that

$$\begin{array}{cccc} B_2 & \text{is of weight} & 2 & \text{and involves } a_2, \\ B_3 & & 3 & & a_3, \\ \dots & & \dots & & \dots \\ B_p & & p & & a_p; \end{array}$$

necessities which require that no coefficient  $a_r$  of the quantic occurs in a  $B$  with a lower suffix than  $r$ , and that  $a_r$  occurs in  $B_r$  multiplied by a power of  $a_0$  only.

To see this, take any seminvariant, and, if  $a_p$  occur in it, take the expression for  $B_p$ ,

$$B_p = a_0^\lambda a_p + f(a_0, a_1, a_2, \dots a_{p-1}),$$

which gives

$$a_p = a_0^{-\lambda} \{B_p - f(a_0, a_1, a_2, \dots a_{p-1})\},$$

and substitute in the seminvariant this value for  $a_p$ . The seminvariant is then expressed as a rational function of  $B_p$  and  $a_0, a_1, a_2, \dots a_{p-1}$ , integral in all of them but  $a_0$ .

Again, for  $a_{p-1}$  substitute in like manner in terms of  $B_{p-1}$  and  $a_0, a_1, a_2, \dots a_{p-2}$ ; and continue this process as long as possible. We thus obtain an expression for the seminvariant

$$S(a_0, a_1, a_2, \dots a_p) \equiv a_0^{-\mu} F(a_0, a_1, B_2, B_3, \dots B_p).$$

But  $a_1$  cannot, as a matter of fact, enter in  $F$ . To see this operate on both sides with  $\Omega$ , which annihilates  $S$  and  $a_0, B_2, B_3, \dots B_p$ . We obtain

$$0 = a_0^{-\mu} a_0 \frac{d}{da_1} F(a_0, a_1, B_2, B_3, \dots B_p) = 0,$$

i.e. 
$$\frac{dF}{da_1} = 0,$$

the differentiation being partial with regard to  $a_1$  as it occurs explicitly in  $F$ . The conclusion is that it cannot so occur. It is proved then that

$$S(a_0, a_1, a_2, \dots a_p) = a_0^{-\mu} F(a_0, B_2, B_3, \dots B_p),$$

for some rational integral form of  $F$  and for some integral or zero value of  $\mu$ .

Notice the two special conveniences of the protomorphs  $A_2, A_3, \dots A_p$  of § 163. One is that the form of  $F$  for them is at once written down from the form of  $S$ ; and the other

that  $F$  does not involve  $a_0$  explicitly. For these protomorphs weight extent and degree are all equal. For others weight and extent only.

Protomorphs must not be confused with *irreducible* seminvariants. A complete system of irreducible seminvariants and invariants, proved to exist in the preceding chapter, is a system in terms of which all rational integral seminvariants can be rationally and integrally expressed, without the occurrence in any of them of a negative power of  $a_0$  as factor.

165.] **Protomorphs of lowest degrees.** The system of protomorphs which has been most used is a system

$$a_0, C_2, C_3, C_4, \dots C_p$$

in which each is of the lowest possible degree.

Those of even weights 2, 4, 6, ... are of the second degree; viz. the system of § 114

$$C_2 \equiv a_0 a_2 - a_1^2 \equiv A_2,$$

$$C_4 \equiv a_0 a_4 - 4 a_1 a_3 + 3 a_2^2,$$

$$C_6 \equiv a_0 a_6 - 6 a_1 a_5 + 15 a_2 a_4 - 10 a_3^2,$$

$$\dots \dots \dots$$

$$C_{2n} \equiv a_0 a_{2n} - \binom{2n}{1} a_1 a_{2n-1} + \binom{2n}{2} a_2 a_{2n-2} - \dots$$

$$+ (-1)^{n-1} \binom{2n}{n-1} a_{n-1} a_{n+1} + \frac{1}{2} (-1)^n \binom{2n}{n} a_n^2,$$

where  $\binom{2n}{r}$  denotes the number of combinations of  $2n$  things  $r$  together.

For odd weights there are (§ 114) no seminvariants of the second degree. In each case, however, there is a protomorph of degree 3. These can be found by the method of § 114, by determining gradients of degree 3 and the requisite odd weights which  $\Omega$  annihilates.

A readier way of calculating them is afforded by the theorem of Cayley's that, if  $S$  be a seminvariant of degree  $i$  which does not involve  $a_p$ , then

$$\{a_0 \mathfrak{D} - a_1 i\} S,$$

where  $\mathfrak{D}$  denotes the operator

$$a_1 \frac{d}{da_0} + a_2 \frac{d}{da_1} + \dots + a_p \frac{d}{da_{p-1}},$$



is another seminvariant whose degree weight and extent are each one greater than those of  $S$ . The facts as to degree weight and extent are clear. That the gradient educed from  $S$  is annihilated by  $\Omega$ , and is therefore a seminvariant, may be seen as follows.

By the method of § 123 we have the equality of operators

$$\begin{aligned} \Omega\mathfrak{D} - \mathfrak{D}\Omega &= \left( a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + \dots + pa_{p-1} \frac{d}{da_p} \right) \\ &\quad \left( a_1 \frac{d}{da_0} + a_2 \frac{d}{da_1} + \dots + a_p \frac{d}{da_{p-1}} \right) \\ &\quad - \left( a_1 \frac{d}{da_0} + a_2 \frac{d}{da_1} + \dots + a_p \frac{d}{da_{p-1}} \right) \\ &\quad \left( a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + \dots + pa_{p-1} \frac{d}{da_p} \right) \\ &= a_0 \frac{d}{da_0} + 2a_1 \frac{d}{da_1} + \dots + pa_{p-1} \frac{d}{da_{p-1}} \\ &\quad - \left( a_1 \frac{d}{da_1} + 2a_2 \frac{d}{da_2} + \dots + pa_p \frac{d}{da_p} \right) \\ &= \left( a_0 \frac{d}{da_0} + a_1 \frac{d}{da_1} + \dots + a_p \frac{d}{da_p} \right) - (p+1)a_p \frac{d}{da_p}. \end{aligned}$$

Hence, when the operation is on  $S$ , a seminvariant (annihilated by  $\Omega$ ) of degree  $i$ , and by supposition free from  $a_p$ ,

$$\Omega\mathfrak{D}S = iS,$$

whence 
$$\Omega \{ a_0\mathfrak{D} - a_1i \} S = a_0iS - a_0iS - a_1i\Omega S = 0.$$

Consequently  $(a_0\mathfrak{D} - a_1i)S$ , being annihilated by  $\Omega$ , is a seminvariant.

Use of this theorem gives us, from the even weighted or quadratic protomorphs  $C_2, C_4, C_6, \dots$  in succession, the odd weighted or cubic protomorphs  $C_3, C_5, C_7, \dots$  in succession; viz.

$$\begin{aligned} C_3 &\equiv (a_0\mathfrak{D} - 2a_1)C_2 \\ &\equiv a_0^2a_3 - 3a_0a_1a_2 + 2a_1^3 \equiv A_3, \\ C_5 &\equiv (a_0\mathfrak{D} - 2a_1)C_4 \\ &\equiv a_0^2a_5 - 5a_0a_1a_4 + 2a_0a_2a_3 + 8a_1^2a_3 - 6a_1a_2^2, \\ C_7 &\equiv a_0^2a_7 - 7a_0a_1a_6 + 9a_0a_2a_5 - 5a_0a_3a_4 + 12a_1^2a_5 + 20a_1a_3^2 \\ &\quad - 30a_1a_2a_4, \end{aligned}$$

&c., &c.

Ex. 7. The operator  $a_0\mathfrak{D} - a_1i$  may be written

$$(a_0a_2 - a_1^2) \frac{d}{da_1} + (a_0a_3 - a_1a_2) \frac{d}{da_2} + (a_0a_4 - a_1a_3) \frac{d}{da_3} + \dots,$$

in which the last term is  $-a_1a_p \frac{d}{da_p}$ .

Ex. 8. Prove that

$$a_0 \left\{ a_2 \frac{d}{da_1} + 2a_3 \frac{d}{da_2} + \dots + (p-1)a_p \frac{d}{da_{p-1}} \right\} - 2a_1w, \text{ or}$$

$$(a_0a_2 - 2a_1^2) \frac{d}{da_1} + 2(a_0a_3 - 2a_1a_2) \frac{d}{da_2} + 3(a_0a_4 - 2a_1a_3) \frac{d}{da_3} + \dots,$$

is another operator which educes from seminvariants of a  $p$ -ic which do not involve  $a_p$  other seminvariants, and by means of it educe the same series of cubic protomorphs  $C_3, C_5, C_7, \dots$  from the quadratic protomorphs. (*Cayley*.)

Ex. 9. Prove that  $a_0O - a_1(ip - 2w)$  educes seminvariants from all seminvariants of a  $p$ -ic, and derives  $C_3, C_5, C_7, \dots$  from  $C_2, C_4, C_6, \dots$ .

166.] If  $\gamma_2, \gamma_3, \dots, \gamma_p$  be the covariants whose leaders are the protomorphs  $C_2, C_3, \dots, C_p$ , then the expression for a seminvariant  $S$  in terms of them and  $a_0$ ,

$$S = a_0^{-\mu} F(a_0, C_2, C_3, \dots, C_p),$$

where the function  $F$  is rational and integral for a rational integral  $S$ , leads to the expression for the covariant  $K$  whose leader is  $S$ ,

$$K = u^{-\mu} F(u, \gamma_2, \gamma_3, \dots, \gamma_p).$$

This follows by the same argument as in § 163. In fact, in general the expression for a seminvariant or invariant in terms of any system of protomorphs leads to exactly the same expression for the corresponding covariant or the invariant in terms of the covariants led by the protomorphs.

167.] **Seminvariants as integrals of  $\Omega S = 0$ .** Another aspect of the reason for the expressibility of any seminvariant or invariant in terms of  $p$  independent ones should be noticed. A seminvariant is an integral of the differential equation

$$a_0 \frac{dS}{da_1} + 2a_1 \frac{dS}{da_2} + 3a_2 \frac{dS}{da_3} + \dots + pa_{p-1} \frac{dS}{da_p} = 0,$$

which is properly regarded as beginning with a vanishing multiple of  $\frac{dS}{da_0}$ .

Now Lagrange's theory of linear partial differential equations (*Forsyth*, § 184) tells us that when we have  $p$  independent functions  $S$  of  $a_0, a_1, a_2, \dots, a_p$  which satisfy this equation, any other  $S$  which satisfies the equation is a function of those  $p$ .

Now in §§ 161, 163, 164, 165 we have sets of  $p$  independent solutions, viz.

$$\begin{aligned} a_0, a_2', a_3', \dots, a_p'; \\ a_0, A_2, A_3, \dots, A_p; \\ a_0, B_2, B_3, \dots, B_p; \\ a_0, C_2, C_3, \dots, C_p. \end{aligned}$$

We thus have it clearly exhibited that any seminvariant or invariant, even though fractional or irrational, is capable of expression in terms of a set of protomorphs.

168.] Protomorphs for systems of quantics. A seminvariant of the system consisting of two binary quantics

$$\begin{aligned} (a_0, a_1, a_2, \dots, a_p)(x, y)^p, \\ (b_0, b_1, b_2, \dots, b_{p'}) (x, y)^{p'}, \end{aligned}$$

is (§ 115) a solution of the differential equation

$$\begin{aligned} \left( a_0 \frac{dS}{da_1} + 2a_1 \frac{dS}{da_2} + \dots + pa_{p-1} \frac{dS}{da_p} \right) \\ + \left( b_0 \frac{dS}{db_1} + 2b_1 \frac{dS}{db_2} + \dots + p'b_{p'-1} \frac{dS}{db_{p'}} \right) = 0, \end{aligned}$$

in the  $p+p'+2$  variables  $a_0, a_1, a_2, \dots, a_p; b_0, b_1, b_2, \dots, b_{p'}$ . We need  $p+p'+1$  independent solutions of this equation: and these are afforded by a set of  $p$  protomorphs of the  $p$ -ic, a set of  $p'$  protomorphs of the  $p'$ -ic, and the one additional  $a_0b_1 - a_1b_0$ . All seminvariants of the system can then be expressed in terms of these  $p+p'+1$  seminvariants.

Moreover, we can easily prove a theorem due to Clebsch, that the expression for any rational integral seminvariant of the system can be expressed in terms of  $a_0b_1 - a_1b_0$  and sets of protomorphs  $a_0, B_2, B_3, \dots, B_p; b_0, B_2', B_3', \dots, B_{p'}$  in a form

which is rational, and is integral except as regards  $a_0$  and  $b_0$ . As in § 164, we reduce it to the form

$$a_0^{-\mu} b_0^{-\mu'} F(a_0, a_1, B_2, B_3, \dots B_p; b_0, b_1, B_2', B_3', \dots B_p'),$$

where  $F$  is rational and integral in its arguments. We also, upon expressing the annihilation by  $\Sigma\Omega$ , obtain that

$$\left(a_0 \frac{d}{da_1} + b_0 \frac{d}{db_1}\right) F = 0;$$

and this tells us that  $F$  involves  $a_1, b_1$  in the connexion  $a_0 b_1 - a_1 b_0$  only.

And, quite generally, rational integral seminvariants (including, of course, invariants) of any number of binary quantics are rational functions of sets of protomorphs of those quantics severally, and of the leaders

$$a_0 b_1 - a_1 b_0, a_0 c_1 - a_1 c_0, \dots$$

of the Jacobians of one of the quantics and the rest, which are integral except as to powers of  $a_0, b_0, c_0, \dots$ , the protomorphs that are the leaders of the quantics themselves.

169.] **Protomorphs applied to the analysis of irreducible systems. The Cubic.** There is a method due to Cayley for finding the complete system of irreducible seminvariants and invariants of a binary quantic, and therefore the system of irreducible covariants and invariants, from a system of protomorphs, which is simple for the cases of the cubic and the quartic.

It is of very little consequence whether we start from the system of protomorphs  $a_0, A_2, A_3, \dots A_p$  of § 163, or the system  $a_0, C_2, C_3, \dots C_p$  of § 165. For the cubic these systems are the same, since  $A_2 \equiv C_2$  and  $A_3 \equiv C_3$ .

By § 163 any seminvariant of degree  $i$  and weight  $w$  of the cubic  $(a, b, c, d)(x, y)^3$  is of the form

$$a^{-w+i} F(A_2, A_3),$$

where  $F(A_2, A_3)$  is rational and integral.

Here, if  $i > w$ , the seminvariant is integrally expressed, and has the positive power  $a^{i-w}$  of  $a$  as a factor. It is then a rational integral function of  $a, A_2, A_3$ , and is not irreducible, unless it be only  $a$  itself.

If  $i = w$  the seminvariant is a rational integral function  $F(A_2, A_3)$  of  $A_2$  and  $A_3$ , and is not irreducible unless it be either  $A_2$  or  $A_3$  itself.

Any irreducible seminvariant, other than  $a, A_2, A_3$ , must then be the result of dividing some rational integral function of  $A_2$  and  $A_3$  by a power of  $a$ . Such a function, being divisible by  $a$ , must vanish when  $a = 0$ , i.e. when we put for  $A_2, A_3$  the values

$$A_2' = -b^2, \quad A_3' = 2b^3.$$

Now the one rational integral function of these which vanishes is

$$A_3'^2 + 4A_2'^3.$$

Any other can only vanish in consequence of having this for a factor. We are thus led to form

$$\begin{aligned} A_3^2 + 4A_2^3 &\equiv a^2\{a^2d^2 - 6abcd + 4ac^3 + 4b^3d - 3b^2c^2\} \\ &\equiv a^2\Delta, \end{aligned}$$

and to conclude that  $\Delta$  is a seminvariant—it is of course the discriminant, a full invariant. It is found as one whose weight exceeds its degree, so that it is not a rational integral function of  $a, A_2, A_3$ .

We are also led to conclude that any other seminvariant whose weight exceeds its degree is given by the rejection of an  $a$  factor from a function  $F(A_2, A_3)$  of which

$$A_3^2 + 4A_2^3 \equiv a^2\Delta$$

is a factor, and consequently that it has  $\Delta$  for a factor, and is not irreducible, but a rational integral function of  $a, A_2, A_3, \Delta$ .

Thus all rational integral seminvariants of the cubic are rational integral functions of some or all of  $a, A_2, A_3, \Delta$ , which alone are irreducible, and are connected by the syzygy

$$A_3^2 + 4A_2^3 = a^2\Delta.$$

These are the results of § 141.

170.] **Irreducible system for the quartic.** Consider now the quartic  $(a, b, c, d, e)(x, y)^4$ ; and take the protomorphs of § 165,

$a, C_2 \equiv ac - b^2, C_3 \equiv a^2d - 3abc + 2b^3, C_4 \equiv ae - 4bd + 3c^2$ , the terms free from  $a$  in which are

$$0, C_2' \equiv -b^2, C_3' \equiv 2b^3, C_4' \equiv -4bd + 3c^2.$$

As in the last article we are led to a seminvariant (not now an invariant)

$$D \equiv a^2 d^2 - 6abcd + 4ac^3 + 4b^3d - 3b^2c^2,$$

which may, so far as we know at present, turn out to be irreducible, though that it was irreducible in the case of the cubic, for which there were three protomorphs only, affords no reason why it should be now that there are four. The terms free from  $a$  in  $D$  are

$$D' \equiv 4b^3d - 3b^2c^2.$$

Any seminvariant not a rational integral function of these five will, by § 165, be the result of rejecting a factor which is a power of  $a$  from a rational integral function of  $a, C_2, C_3, C_4$ ; and such a rational integral function may present itself in the form

$$F(a, C_2, C_3, C_4, D),$$

where  $F$  is rational and integral in its arguments. The result of putting  $a = 0$  in this is

$$F(0, C_2', C_3', C_4', D'),$$

and must vanish identically, since  $F$ , expressed in terms of  $a, b, c, d, e$ , has  $a$  for a factor.

Now a result, distinct from  $C_3'^2 + 4C_2'^3$  which led to  $D$ , of eliminating  $b, c, d$  from

$$C_2' \equiv -b^2, \quad C_3' \equiv 2b^3, \quad C_4' \equiv -4bd + 3c^2, \quad D' \equiv 4b^3d - 3b^2c^2,$$

is

$$C_2' C_4' - D' = 0,$$

and this leads to

$$\begin{aligned} C_2 C_4 - D &\equiv a(ace + 2bcd - ad^2 - b^2e - c^3) \\ &\equiv aJ, \end{aligned}$$

which shows two things: (1) that there is a new seminvariant  $J$ , an invariant, in fact, which may prove to be irreducible; and (2) that  $D$  is not irreducible, but is equal to  $C_2 C_4 - aJ$ .

We have now further to look for rational integral functions of  $a, C_2, C_3, C_4, J$  which have  $a$  for a factor, so that the same functions of

$$\begin{aligned} 0, \quad C_2' \equiv -b^2, \quad C_3' \equiv 2b^3, \quad C_4' \equiv -4bd + 3c^2, \\ J' \equiv 2bcd - b^2e - c^3 \end{aligned}$$

vanish identically.

But there are no such new functions. For  $C_4'$  involves a letter  $c$  which does not occur in  $C_2'$  or  $C_3'$ , so that no relation connects it with them; and  $J'$  involves  $e$  which does not occur in either  $C_2'$ ,  $C_3'$  or  $C_4'$ , so that it again is independent of the preceding.

Consequently  $a, C_2, C_3, C_4, J$  is the complete system of irreducible seminvariants and invariants of the quartic. Of these  $C_4$  is what in previous chapters we have called  $I$ . The result is that of § 145.

The one syzygy which connects members of the irreducible system (cf. § 145) is also exhibited. We have, as above,

$$C_3^2 + 4C_2^3 = a^2D,$$

and 
$$C_2C_4 - D = aJ,$$

from which, after elimination of the reducible  $D$ , there results

$$a^3J - a^2C_2C_4 + C_3^2 + 4C_2^3 = 0.$$

171.] The method is not suited for extended application to higher binary quantities. It may be pursued in dealing with the quintic and the sextic, but the labour is enormous owing to the number of eliminations and the length and complexity of the functions dealt with. Moreover serious theoretical difficulties present themselves, and without guidance by a knowledge of the results to be obtained those results could hardly be thus arrived at with certainty.

One general fact should be mentioned. The protomorphs  $a_0, C_2, C_3, C_4, \dots C_p$  are in all cases irreducible; for each involves its most advanced letter from among  $a_0, a_1, a_2, \dots a_p$  in the first degree only, and is the seminvariant of lowest degree which exists for its own weight. The same cannot be said of the other system of protomorphs  $a_0, A_2, A_3, \dots A_p$ , which after the third are of higher degrees than the lowest possible.

172.] **A seminvariant arranged by powers of  $a_1$ .** When in a seminvariant, or, in particular, invariant, the terms free from  $a_1$  are known, the whole is known.

This has been proved in § 161. We have only to replace in the given terms  $a_2, a_3, \dots a_p$  by the  $a_2', a_3', \dots a_p'$  of that article.

Any gradient in  $a_0, a_2', a_3', \dots a_p'$  is a seminvariant, but not necessarily an integral seminvariant.

A seminvariant which is integral may be expressed in its integral form, when its terms free from  $a_1$  are known, by aid of differentiations only in virtue of the following.

Let  $Q_i$  be the given terms free from  $a_1$ , and suppose the whole seminvariant to be

$$S \equiv Q_i + a_1 Q_{i-1} + a_1^2 Q_{i-2} + \dots + a_1^m Q_{i-m},$$

where  $Q_i, Q_{i-1}, Q_{i-2}, \dots Q_{i-m}$  are all free from  $a_1$ , and where  $m$  is some number not exceeding  $i$ , the degree of  $S$ .

If we write

$$\begin{aligned} \Omega &\equiv a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + \left( 3a_2 \frac{d}{da_3} + 4a_3 \frac{d}{da_4} + \dots + pa_{p-1} \frac{d}{da_p} \right) \\ &\equiv a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + \omega, \end{aligned}$$

where  $\omega$  does not involve either  $a_1$  or  $\frac{d}{da_1}$ , then

$$\Omega(a_1^\mu Q) = a_1^{\mu-1} \cdot \mu a_0 Q + a_1^\mu \cdot \omega Q + a_1^{\mu+1} \cdot 2 \frac{d}{da_2} Q.$$

Hence, arranging the seminvariant condition  $\Omega S = 0$  by powers of  $a_1$ , and expressing that the parts of  $\Omega S$  with different powers of  $a_1$  for factors must vanish separately, we have the succession of facts

$$\begin{aligned} a_0 Q_{i-1} + \omega Q_i &= 0, \\ 2a_0 Q_{i-2} + \omega Q_{i-1} + 2 \frac{d}{da_2} Q_i &= 0, \\ 3a_0 Q_{i-3} + \omega Q_{i-2} + 2 \frac{d}{da_2} Q_{i-1} &= 0, \\ \dots & \\ \omega Q_{i-m} + 2 \frac{d}{da_2} Q_{i-m+1} &= 0, \\ 2 \frac{d}{da_2} Q_{i-m} &= 0, \end{aligned}$$

of which all but the last two suffice to determine from  $Q_i$  the expressions for  $Q_{i-1}, Q_{i-2}, \dots Q_{i-m}$  in succession by operations with  $\omega$ , i.e. by differentiations.



The last equation tells us that the terms multiplying the highest power of  $a_1$  which occurs in a seminvariant are free from  $a_2$ .

Ex. 10. State the corresponding facts with regard to anti-seminvariants, derived from their annihilator  $O$ .

173.] A simpler method of determining the whole seminvariant  $S$ , when its terms  $Q_i$  free from  $a_1$  are known, is given as follows.

If  $G$  be any gradient whatever, the following is an identity,

$$\Omega \left\{ 1 - \frac{a_1}{a_0} \Omega + \frac{1}{1 \cdot 2} \frac{a_1^2}{a_0^2} \Omega^2 - \frac{1}{1 \cdot 2 \cdot 3} \frac{a_1^3}{a_0^3} \Omega^3 + \dots \right\} G = 0.$$

For the left-hand member is

$$\begin{aligned} & \Omega G \\ & - \frac{a_1}{a_0} \Omega^2 G - \Omega G \\ & + \frac{1}{1 \cdot 2} \frac{a_1^2}{a_0^2} \Omega^3 G + \frac{a_1}{a_0} \Omega^2 G \\ & - \frac{1}{1 \cdot 2 \cdot 3} \frac{a_1^3}{a_0^3} \Omega^4 G - \frac{1}{1 \cdot 2} \frac{a_1^2}{a_0^2} \Omega^3 G \\ & + \dots \end{aligned}$$

of which the terms cancel against one another up to a certain point, and after that point vanish.

Now take  $Q_i$  for the gradient  $G$ . We obtain that

$$\left( 1 - \frac{a_1}{a_0} \Omega + \frac{1}{1 \cdot 2} \frac{a_1^2}{a_0^2} \Omega^2 - \frac{1}{1 \cdot 2 \cdot 3} \frac{a_1^3}{a_0^3} \Omega^3 + \dots \right) Q_i$$

is annihilated by  $\Omega$ , and is consequently a seminvariant if it do not vanish. But it does not vanish since its terms  $Q_i$  free from  $a_1$  do not.

Its form is apparently fractional, but cannot be really so. It must, in fact, be exactly

$$Q_i + a_1 Q_{i-1} + a_1^2 Q_{i-2} + \dots + a_1^m Q_{i-m};$$

for otherwise the difference of the two would be divisible by  $a_1$  and annihilated by  $\Omega$ . Now this is impossible, since no seminvariant can have  $a_1$  for a factor, seminvariants being

functions of  $a_0, a_2 - \frac{a_1^2}{a_0}, a_3 - 3\frac{a_1}{a_0}a_2 + 2\frac{a_1^3}{a_0^2}, \dots$ , which are the non-vanishing and independent  $a_0, a_2, a_3, \dots$  when  $a_1 = 0$ .

The full expression for a seminvariant is then found by operating on its terms free from  $a_1$  with

$$1 - \frac{a_1}{a_0}\Omega + \frac{1}{1 \cdot 2} \frac{a_1^2}{a_0^2}\Omega^2 - \frac{1}{1 \cdot 2 \cdot 3} \frac{a_1^3}{a_0^3}\Omega^3 + \dots$$

Ex. 11. Prove that this operator annihilates any gradient with  $a_1$  for a factor, and produces from any other gradient a seminvariant, not necessarily integral, which is also produced from the terms in it which are free from  $a_1$ .

Ex. 12. If  $G$  be any gradient,  $\Omega(a_1 G)$  involves as many arbitraries as  $G$ .

174.] **Annihilator of terms free from  $a_1$  in invariants.** There does not seem to be a simple general method for finding the terms free from  $a_1$  in integral *seminvariants* of given type. The case is different, however, with regard to *invariants*, which are at once seminvariants and anti-seminvariants. The terms free from  $a_1$  in an invariant have an annihilator, which suffices to determine them. The fact is due to Cayley.

Consider the two annihilators,

$$\Omega = a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + 3a_2 \frac{d}{da_3} + \dots + pa_{p-1} \frac{d}{da_p},$$

$$O = pa_1 \frac{d}{da_0} + (p-1)a_2 \frac{d}{da_1} + (p-2)a_3 \frac{d}{da_2} + \dots + a_p \frac{d}{da_{p-1}},$$

of invariants of a binary  $p$ -ic. Eliminating  $\frac{d}{da_1}$  we have

$$\begin{aligned} a_0 O - (p-1)a_2 \Omega &\equiv a_1 \left\{ pa_0 \frac{d}{da_0} - 2(p-1)a_2 \frac{d}{da_2} \right\} \\ &+ a_0 \left\{ (p-2)a_3 \frac{d}{da_2} + (p-3)a_4 \frac{d}{da_3} + \dots + a_p \frac{d}{da_{p-1}} \right\} \\ &- (p-1)a_2 \left\{ 3a_2 \frac{d}{da_3} + 4a_3 \frac{d}{da_4} + \dots + pa_{p-1} \frac{d}{da_p} \right\} \\ &\equiv a_1 \phi + \psi, \text{ say,} \end{aligned}$$

where  $\phi$  and  $\psi$  do not involve  $a_1$  or  $\frac{d}{da_1}$ .

Now let an invariant, arranged by powers of  $a_1$  as in the last two articles, be written

$$I \equiv R_i + a_1 R_{i-1} + a_1^2 R_{i-2} + \dots + a_1^m R_{i-m}.$$

We find that, since  $\Omega$  and  $O$ , and therefore  $a_0 O - (p-1) a_2 \Omega$ , annihilate  $I$ ,

$$\begin{aligned} \psi R_i &= 0, \\ \psi R_{i-1} + \phi R_i &= 0, \\ \psi R_{i-2} + \phi R_{i-1} &= 0, \\ &\vdots \\ &\vdots \\ &\vdots \\ \phi R_{i-m} &= 0. \end{aligned}$$

Of these the first is the result important for our purpose; but before examining it we notice that the last, viz.

$$\left\{ p a_0 \frac{d}{d a_0} - 2(p-1) a_2 \frac{d}{d a_2} \right\} R_{i-m} = 0,$$

tells us that, since  $\frac{d}{d a_2} R_{i-m} = 0$  by § 172, we must also have  $\frac{d}{d a_0} R_{i-m} = 0$ , i.e. that the terms which multiply the highest power of  $a_1$  which occurs in any invariant are free from both  $a_0$  and  $a_2$ .

The more important conclusion, gathered from the first of the above equalities, is that the terms free from  $a_1$  in an invariant of a binary  $p$ -ic have the annihilator

$$\begin{aligned} \psi \equiv a_0 \left\{ (p-2) a_3 \frac{d}{d a_2} + (p-3) a_4 \frac{d}{d a_3} + \dots + a_p \frac{d}{d a_{p-1}} \right\} \\ - (p-1) a_2 \left\{ 3 a_2 \frac{d}{d a_3} + 4 a_3 \frac{d}{d a_4} + \dots + p a_{p-1} \frac{d}{d a_p} \right\}. \end{aligned}$$

We need to know conversely that all gradients in  $a_0, a_2, a_3, \dots, a_p$ , for which  $ip = 2w$ , and which have  $\psi$  for an annihilator, are the terms free from  $a_1$  in invariants. When this is known we are sure that we have a means, by expressing the annihilation by  $\psi$ , of finding the numerical multipliers in those terms of invariants of a given degree, and thence, in either of the ways of §§ 161, 172, 173, the complete expressions of the invariants.

We shall encounter a proof of this converse proposition in the next chapter (cf. § 186). It amounts to proving that all

gradients for which  $ip = 2w$  in  $a_0, a_2', a_3', \dots a_p'$ , the coefficients in the quantic deprived of its second term as in § 162, which are annihilated by  $\psi'$ , the result of accenting the letters  $a_2, a_3, \dots a_p$  in  $\psi$ , are invariants, i.e. are annihilated by  $O$  as well as by  $\Omega$ , which latter must annihilate them since it annihilates  $a_0, a_2', a_3', a_p'$ . Note that no expression fractional in  $a_0$  can be annihilated by  $O$ .

It is for the present left to the student to arrive at the proposition for himself by the sequence of theorems of the following four examples. He will see from Ex. 16 that if  $G'$  satisfy  $\psi' G' = 0$  it must certainly satisfy  $OG' = 0$ , provided  $ip - 2w = 0$ , as is the case.

Ex. 13. Prove that, if by the substitution  $x = X + mY, y = Y$  the quantic  $(a_0, a_1, a_2, \dots a_p)(x, y)^p$  be transformed into

$(a_0, a_1, a_2, \dots a_p)(X, Y)^p$ , so that (§ 92)

$$a_0 = a_0,$$

$$a_1 = a_1 + a_0 m,$$

$$a_2 = a_2 + 2a_1 m + a_0 m^2, \text{ \&c. \&c.,}$$

then

$$\frac{d}{da_0} = \frac{d}{da_0} + m \frac{d}{da_1} + m^2 \frac{d}{da_2} + \dots + m^p \frac{d}{da_p},$$

and, generally,

$$\frac{d}{da_r} = \frac{1}{r!} \left\{ 1.2.3 \dots r \frac{d}{da_r} + 2.3 \dots (r+1) m \frac{d}{da_{r+1}} + \dots + (p-r+1)(p-r+2) \dots p m^{p-r} \frac{d}{da_p} \right\},$$

the last case of which is

$$\frac{d}{da_p} = \frac{d}{da_p};$$

the operations on the left being all upon a function of  $a_0, a_1, a_2, \dots a_p$ , and those on the right all on the function of  $a_0, a_1, a_2, \dots a_p$  to which it is equal.

Ex. 14. Hence show that the operators

$$i \equiv a_0 \frac{d}{da_0} + a_1 \frac{d}{da_1} + \dots + a_p \frac{d}{da_p},$$

$$\Omega \equiv a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + \dots + p a_{p-1} \frac{d}{da_p},$$

$$w - \frac{a_1}{a_0} \Omega \equiv a_1 \frac{d}{da_1} + 2a_2 \frac{d}{da_2} + \dots + pa_p \frac{d}{da_p} - \frac{a_1}{a_0} \left\{ a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + \dots + pa_{p-1} \frac{d}{da_p} \right\},$$

$$O - \frac{a_1}{a_0} (ip - 2w) - \frac{a_1^2}{a_0^2} \Omega \equiv pa_1 \frac{d}{da_0} + (p-1)a_2 \frac{d}{da_1} + \dots + a_p \frac{d}{da_{p-1}} - \frac{a_1}{a_0} \left\{ pa_0 \frac{d}{da_0} + (p-2)a_1 \frac{d}{da_1} + (p-4)a_2 \frac{d}{da_2} + \dots - pa_p \frac{d}{da_p} \right\} - \frac{a_1^2}{a_0^2} \left\{ a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + \dots + pa_{p-1} \frac{d}{da_p} \right\},$$

transform into operators of like forms.

Ex. 15. Show that the only necessary modification of the above when the operation is on a seminvariant, and when  $m$  is the non-constant  $-\frac{a_1}{a_0}$ , and  $a_0, a_1, a_2, \dots, a_p$  are consequently the  $a_0, 0, a_2', a_3', a_p'$  of § 162, is that the undetermined  $\frac{d}{da_1'}$  is to be taken as defined by the persistence in form of  $\Omega$ .

Ex. 16. By this and the fourth of Ex. 14 prove that the effect of operating with  $\psi'$ , i.e.

$$a_0 \left\{ (p-2)a_3' \frac{d}{da_2'} + (p-3)a_4' \frac{d}{da_3'} + \dots + a_{p-1}' \frac{d}{da_{p-1}'} \right\} - (p-1)a_2' \left\{ 3a_2' \frac{d}{da_3'} + 4a_3' \frac{d}{da_4'} + \dots + pa_{p-1}' \frac{d}{da_p'} \right\},$$

on a gradient in  $a_0, a_2', a_3', \dots, a_p'$  is the same as that of operating on the equivalent function of  $a_0, a_1, a_2, \dots, a_p$  with  $a_0 O - a_1(ip - 2w)$ , since  $\Omega$  annihilates it.

175.] Taking  $p = 5$ , i.e. the case of the quintic

$$(a, b, c, d, e, f)(x, y)^5,$$

the annihilator  $\psi$  is

$$3ad\partial_e + (2ae - 12c^2)\partial_d + (af - 16cd)\partial_c - 20ce\partial_f.$$

An invariant of the quintic of degree  $i$ , and consequently of weight  $\frac{5i}{2}$ , if such exist, can then be found as follows. Write down the most general gradient of the type in  $a, c, d, e, f$ .

Operate on it with the annihilator above, and equate to zero the coefficients of the various terms in the result. If the equations can be satisfied by values of the arbitrary coefficients in the assumed gradient which are not all zero, we obtain as many linearly independent invariants of the type as there are coefficients left arbitrary. If they cannot be so satisfied there is no invariant of the type.

The terms free from  $b$  in the invariant are thus found. If  $R_i$  be the said terms, the whole expression for the invariant of which  $R_i$  is part is by § 173

$$e^{-\beta\Omega}R_i,$$

where, after the operations are performed,  $\beta$  is to be replaced by  $\frac{b}{a}$ .

In this way the invariants of degrees 4, 8, 12, 18 of the quintic may, with much labour in the last two cases, be calculated.

For the sextic  $(a, b, c, d, e, f, g)(x, y)^6$  the annihilator  $\psi$  of terms free from  $b$  in invariants is

$$4ad\partial_c + (3ae - 15c^2)\partial_a + (2af - 20cd)\partial_e + (ag - 25ce)\partial_f - 30cf\partial_g,$$

by means of which the invariants of degrees 2, 4, 6, 10, 15 may be found.

And similarly for higher quantics in succession.

Ex. 17. Integrate by Lagrange's method the differential equation for the case of the cubic

$$\psi G \equiv (ad\partial_c - 6c^2\partial_a)G = 0,$$

and thus show, remembering  $3i = 2w$ , that an invariant of the cubic is necessarily a power of  $a^2d'^2 + 4ac'^3$ , where

$$c' = c - \frac{b^2}{a}, \quad d' = d - \frac{3bc}{a} + \frac{2b^3}{a^2}.$$

Ex. 18. Integrate the differential equation for the case of the quartic

$$\psi G \equiv \{2ad\partial_c + (ae - 9c^2)\partial_a - 12cd\partial_e\} G = 0,$$

and show that invariants of the quartic are functions of the invariants

$$ae' + 3c'^2, \quad ac'e' - ad'^2 - c'^3.$$

176.] **Seminvariants arranged by powers of their most advanced\* letter.** The present is a convenient place for a theorem or two not directly connected with the rest of the chapter.

Take a seminvariant  $S$  of extent  $p$ . We are not necessarily regarding it as a seminvariant of a  $p$ -ic in particular. It is equally one of course of any binary quantic of order not less than  $p$ , whose first  $p+1$  coefficients (after rejection of their binomial factors) are the  $a_0, a_1, a_2, \dots, a_p$  involved in  $S$ .

Arrange  $S$  according to powers of  $a_p$ , its most advanced letter, and write it

$$S \equiv a_p^n P_0 + a_p^{n-1} P_1 + a_p^{n-2} P_2 + \dots + P_n,$$

where suffixes of  $P$ 's do not of course, as they did in § 172, &c., indicate degree.

Express the annihilation of  $S$  by  $\Omega$ , i. e. by

$$a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + \dots + (p-1)a_{p-2} \frac{d}{da_{p-1}} + pa_{p-1} \frac{d}{da_p} \cdot$$

The terms involving different powers of  $a_p$  in  $\Omega S$  must vanish separately, for the vanishing is identical. Hence

$$\begin{aligned} \Omega P_0 &= 0, \\ \Omega P_1 + n p a_{p-1} P_0 &= 0, \\ \Omega P_2 + (n-1) p a_{p-1} P_1 &= 0, \\ \cdot &\cdot \cdot \cdot \cdot \cdot \cdot \cdot \cdot \\ \cdot &\cdot \cdot \cdot \cdot \cdot \cdot \cdot \cdot \\ \Omega P_n + p a_{p-1} P_{n-1} &= 0. \end{aligned}$$

From these identities we draw, among others, the following conclusions.

(1)  $P_0$ , the function of  $a_0, a_1, a_2, \dots, a_{p-1}$  which is the coefficient of the highest power of  $a_p$ , the most advanced letter which occurs, in a seminvariant  $S$ , is itself a seminvariant.

(2) When  $P_n$ , which consists of the terms free from  $a_p$ , the most advanced letter in a seminvariant, is known, the rest of the seminvariant can be found by aid of a succession of operations with  $\Omega$  and divisions by multiples of  $a_{p-1}$ , i. e. by aid of differentiations and elementary algebraical processes only.

Ex. 19. In the case of an invariant, prove that also when the terms free from  $a_0$  are known the whole invariant can be written down by aid of differentiations and elementary algebraical processes only.

Ans. Consider the annihilation by  $O$  as we have that by  $\Omega$ .

177.] **Seminvariants as seminvariants of seminvariants.** Another interesting conclusion which can be drawn from the identities of the preceding article is that any seminvariant of a seminvariant of a binary quantic, regarded as itself a binary quantic in  $a_p : 1$ , where  $a_p$  is the most advanced letter involved, is a seminvariant of the original quantic.

If  $f(P_0, P_1, P_2, \dots, P_n)$

be any function of the  $P$ 's,

$$\begin{aligned} \Omega f(P_0, P_1, P_2, \dots, P_n) &= \frac{df}{dI_0} \Omega P_0 + \frac{df}{dP_1} \Omega P_1 + \frac{df}{dP_2} \Omega P_2 + \dots + \frac{df}{dP_n} \Omega P_n \\ &= -pa_{p-1} \left\{ nP_0 \frac{d}{dP_1} + (n-1)P_1 \frac{d}{dP_2} + \dots \right. \\ &\quad \left. + P_{n-1} \frac{d}{dP_n} \right\} f(P_0, P_1, P_2, \dots, P_n), \end{aligned}$$

by the identities proved.

Now the seminvariant  $S$  of the last article is

$$(P_0, P_1, P_2, \dots, P_n) (a_p, 1)^n,$$

and if we write this

$$(P_0, P_1', P_2', \dots, P_n) (a_p, 1)^n,$$

we have generally

$$P_r = \frac{n(n-1)\dots(n-r+1)}{r!} P_r'.$$

Now, with this change of notation,

$$\begin{aligned} nP_0 \frac{d}{dP_1} + (n-1)P_1 \frac{d}{dP_2} + (n-2)P_2 \frac{d}{dP_3} + \dots + P_{n-1} \frac{d}{dP_n} \\ \equiv P_0 \frac{d}{dP_1'} + 2P_1' \frac{d}{dP_2'} + 3P_2' \frac{d}{dP_3'} + \dots + nP_{n-1}' \frac{d}{dP_n'}, \end{aligned}$$

which is of the form of  $\Omega$ , and annihilates only functions of the  $P$ 's which are seminvariants of  $S$  looked upon as a quantic in  $a_p : 1$ .



Such seminvariants are then also annihilated by  $\Omega$ , i.e. they are seminvariants of the quantic whose coefficients are  $a_0, a_1, a_2, a_3, \dots$ .

This includes as a very particular case the result (1) of the preceding article.

178.] **Seminvariants derived by differentiation of seminvariants.** One more fact with regard to any seminvariant and its most advanced letter  $a_p$  may be mentioned.

We immediately prove the alternant identity

$$\Omega \frac{d}{da_p} - \frac{d}{da_p} \Omega = 0, \text{ or } = -(p+1) \frac{d}{da_{p+1}},$$

according as  $\Omega$  does or does not extend beyond  $a_p$ .

If then  $S$  is a seminvariant which does not extend beyond  $a_p$

$$\Omega \frac{dS}{da_p} = 0,$$

i.e.  $\frac{dS}{da_p}$  is a seminvariant.

By repetition it follows that  $\frac{d^2S}{da_p^2}, \frac{d^3S}{da_p^3}, \dots$  are seminvariants.

This again includes as a particular case the result (1) of § 176.

If  $a_r$  be a letter short of the last  $a_p$  which occurs in a seminvariant,

$$\frac{d}{da_r} S = -\frac{1}{r} \Omega \frac{dS}{da_{r-1}}.$$

The theorems of these last articles are partly Cayley's and partly Sylvester's.

179.] **Passage from discriminant to discriminant.** To Cayley is due an application of § 176 (2) to the determination of the discriminant of a binary  $p$ -ic from that of a binary  $(p-1)$ -ic. Let us apply it to find the discriminant of a cubic from that of a quadratic.

The discriminant of  $(a, b, c, d)(x, y)^3$  is an invariant which, when we put  $d = 0$  in it, becomes the discriminant of

$$x(ax^2 + 3bxy + 3cy^2),$$

which is a numerical multiple of the product of the squares of differences between 0,  $\alpha, \beta$ , where  $\alpha, \beta$  are the roots of

$ax^2 + 3bxy + 3cy^2$ , with the factor  $a^4$  necessitated by its known degree. Now this is a numerical multiple of  $a^4(a\beta)^2(a-\beta)^2$ , i.e. of

$$c^2(4ac - 3b^2).$$

These then are the terms free from the most advanced coefficient  $d$  in the discriminant of the cubic. The other terms are determined from them as an example of § 176.

180.] **Operators which generate seminvariants.** Seminvariants and invariants can be derived from gradients which are not seminvariants by operations involving differentiation and elementary algebraical processes only. In fact, *all* seminvariants can be thus obtained. The idea is Hilbert's.

It was proved in § 128 that, if  $G$  be any gradient in  $a_0, a_1, a_2, \dots, a_p$ , or some of them, for which  $\eta \equiv ip - 2w$  is positive,

$$\left\{ 1 - \frac{1}{1 \cdot \eta} \Omega O + \frac{1}{1 \cdot 2 \cdot \eta(\eta+1)} \Omega O^2 \Omega - \frac{1}{1 \cdot 2 \cdot 3 \cdot \eta(\eta+1)(\eta+2)} \Omega O^3 \Omega^2 + \dots \right\} G = 0,$$

and in § 100 was proved the really equivalent theorem that in the same case

$$\left\{ 1 - \frac{\Omega O}{1^2} + \frac{\Omega^2 O^2}{1^2 \cdot 2^2} - \frac{\Omega^3 O^3}{1^2 \cdot 2^2 \cdot 3^2} + \dots \right\} G = 0.$$

Now for  $G$  put  $\Omega F$ , where  $F$  is any gradient of degree  $i$  and weight  $w+1$  in  $a_0, a_1, a_2, \dots, a_p$  or some of them. The two theorems tell us that

$$\Omega \left\{ 1 - \frac{1}{1 \cdot \eta} O \Omega + \frac{1}{1 \cdot 2 \cdot \eta(\eta+1)} O^2 \Omega^2 - \frac{1}{1 \cdot 2 \cdot 3 \cdot \eta(\eta+1)(\eta+2)} O^3 \Omega^3 + \dots \right\} F = 0$$

and  $\Omega \left\{ 1 - \frac{O \Omega}{1^2} + \frac{\Omega O^2 \Omega}{1^2 \cdot 2^2} - \frac{\Omega^2 O^3 \Omega}{1^2 \cdot 2^2 \cdot 3^2} + \dots \right\} F = 0.$

Now write  $w$  instead of  $w+1$ , so that  $F$  is any gradient of weight  $w$  and degree  $i$  in  $a_0, a_1, a_2, \dots, a_p$ , or some of them, for

which  $\eta \equiv ip - 2w$  is greater than  $-2$ , i.e.  $\ngtr -1$ . We have to put  $\eta + 2$  for  $\eta$ , and deduce that

$$\left\{ 1 - \frac{1}{1 \cdot (\eta + 2)} O\Omega + \frac{1}{1 \cdot 2 \cdot (\eta + 2)(\eta + 3)} O^2\Omega^2 - \frac{1}{1 \cdot 2 \cdot 3 \cdot (\eta + 2)(\eta + 3)(\eta + 4)} O^3\Omega^3 - \dots \right\} F,$$

or its equivalent

$$\left\{ 1 - \frac{O\Omega}{1^2} + \frac{\Omega O^2\Omega}{1^2 \cdot 2^2} - \frac{\Omega^2 O^3\Omega}{1^2 \cdot 2^2 \cdot 3^2} + \dots \right\} F,$$

is annihilated by  $\Omega$ , so that it is either zero or a seminvariant.

In particular, taking  $\eta \equiv ip - 2w = 0$ , so that the first expression is

$$\left\{ 1 - \frac{O\Omega}{1 \cdot 2} + \frac{O^2\Omega^2}{1 \cdot 2^2 \cdot 3} - \frac{O^3\Omega^3}{1 \cdot 2^2 \cdot 3^2 \cdot 4} + \dots \right\} F,$$

this or the second expression is either zero or an invariant.

If  $\eta \equiv ip - 2w = -1$  for  $F$ , the first expression is

$$\left\{ 1 - \frac{O\Omega}{1^2} + \frac{O^2\Omega^2}{1^2 \cdot 2^2} - \frac{O^3\Omega^3}{1^2 \cdot 2^2 \cdot 3^2} + \dots \right\} F,$$

and this, or its equivalent the second expression, is necessarily zero, for there are no seminvariants (cf. § 112) for which  $ip - 2w = -1$ .

For the sake of unity of statement let us adopt the second general form, though the former is in most cases the one best adapted for actual calculation. The general conclusion arrived at is that if we write down any gradient or sum of gradients whatever, with arbitrary multipliers, and arbitrary degrees and weights subject to  $ip - 2w \ngtr -1$ , the result of operating on that sum with

$$1 - \frac{O\Omega}{1^2} + \frac{\Omega O^2\Omega}{1^2 \cdot 2^2} - \frac{\Omega^2 O^3\Omega}{1^2 \cdot 2^2 \cdot 3^2} + \dots$$

is a sum of seminvariants and invariants, except for cases when it vanishes, as it must in particular for degree weights subject to  $ip - 2w = -1$ .

181.] We thus obtain all seminvariants and invariants whatever. To see this it suffices to observe that the result of

operating on any seminvariant  $S$  is to produce  $S$  itself, for  $\Omega S = 0$ , so that  $O\Omega S = 0$ ,  $\Omega O^2\Omega S = 0$ , &c.

If, then, we operate on the most general gradient of weight  $w$ , degree  $i$ , and extent in no term exceeding  $p$ , which accordingly contains  $(w; i, p)$  arbitraries, we obtain a result with  $(w; i, p) - (w-1; i, p)$  arbitraries, which is the most general seminvariant of the type in question. There is always the requirement  $ip - 2w \leq -1$ .

We may distinguish between those gradients of type  $w, i, p$  from which the operator

$$1 - \frac{O\Omega}{1 \cdot (\eta+2)} + \frac{O^2\Omega^2}{1 \cdot 2 \cdot (\eta+2)(\eta+3)} - \frac{O^3\Omega^3}{1 \cdot 2 \cdot 3 \cdot (\eta+2)(\eta+3)(\eta+4)} + \dots,$$

or its equivalent, produces seminvariants, and those which it annihilates. The latter are those gradients  $F$  of the type which are of the form  $OF'$ . (Note that if  $ip - 2(ip-w)$ , i. e.  $2w - ip$ , were positive, all gradients  $F$  would be of the form  $OF'$ , by the duality of  $\Omega$  and  $O$  and the fact that when  $ip - 2w$  is positive all gradients are of the form  $\Omega F''$ . But we are attending to cases in which  $ip - 2w \leq -1$ , for which, except in the one case where  $\leq$  is replaced by  $=$ , there is no such expression in general possible.)

This we can see as follows. First, if the result of operation vanishes,  $F$  is of the form  $OF'$ , for the expression of the vanishing may be written

$$F = O \left\{ \frac{\Omega}{1 \cdot (\eta+2)} - \frac{O\Omega^2}{1 \cdot 2 \cdot (\eta+2)(\eta+3)} + \dots \right\} F',$$

which is of the form in question. Secondly, the operator must annihilate an  $OF'$ . For

$$\begin{aligned} & \left\{ 1 - \frac{O\Omega}{1 \cdot (\eta+2)} + \frac{O^2\Omega^2}{1 \cdot 2 \cdot (\eta+2)(\eta+3)} - \dots \right\} OF' \\ &= O \left\{ 1 - \frac{\Omega O}{1 \cdot (\eta+2)} + \frac{O\Omega^2 O}{1 \cdot 2 \cdot (\eta+2)(\eta+3)} - \dots \right\} F', \\ &= O \left\{ 1 - \frac{\Omega O}{1 \cdot (\eta+2)} + \frac{\Omega O^2 \Omega}{1 \cdot 2 \cdot (\eta+2)(\eta+3)} - \dots \right\} F', \end{aligned}$$

since  $O\Omega \cdot \Omega O = \Omega O \cdot O\Omega$ ,  $O^2\Omega^2 \cdot \Omega O = \Omega O \cdot O^2\Omega^2, \dots$  by § 125, Ex. 4. Now this must vanish, for  $\eta + 2$  is the  $\eta$  of  $F'$  and is positive, so that the expression is  $O$  operating on a vanishing result, by the first relation of § 180.

The same conclusions may be drawn from the second form of the operator.

182.] To determine seminvariants and invariants by this method we naturally operate on the simplest gradients we can choose, i.e. on single products of letters chosen from among  $a_0, a_1, a_2, \dots a_p$ . Unfortunately no simple rule presents itself as to what products can and what cannot be written in the form  $OF'$ , i.e. what products lead to seminvariants or invariants and what to zeroes.

In the next chapter we shall, however, see that when we are not limited to a particular extent  $p$  a like method can be employed with perfect definiteness, and we can assign an exact system of products to which there is a one to one correspondence of seminvariants.

Ex. 20. Obtain the invariant  $ace + 2bcd - ad^2 - b^2e - c^3$  of a quartic by operating on a single term of it with

$$1 - \frac{O\Omega}{1.2} + \frac{O^2\Omega^2}{1.2^2.3} - \frac{O^3\Omega^3}{1.2^2.3^2.4} + \dots$$

Ex. 21. If  $F = OF'$ , where  $F$  is of the type  $w, i, p$  and  $ip - 2w \nless - 1$ , prove that

$$F' = \Omega \left\{ \frac{1}{1^2} - \frac{\Omega O}{1^2.2^2} + \frac{\Omega^2 O^2}{1^2.2^2.3^2} - \dots \right\} F.$$

Hence if  $F'$  be general of type  $w - 1, i, p$ , so as to involve  $(w - 1; i, p)$  arbitrary coefficients,  $F$  or  $OF'$  must also be a sum of  $(w - 1; i, p)$  independent multiples of linearly independent gradients.

Ex. 22. The gradients of type  $w, i, p, ip - 2w \nless - 1$ , which are of the form  $OF'$ , are linear functions of the coefficients of  $x^{ip-w}$  in products of order  $ip - w + 1$  in  $x$  of  $i$  quantities chosen from

$$\begin{aligned} & a_p, \\ & a_{p-1}x + a_p, \\ & a_{p-2}x^2 + 2a_{p-1}x + a_p, \\ & \dots \dots \dots \\ & a_0x^p + pa_1x^{p-1} + \dots + a_p. \end{aligned}$$

Ex. 23. If  $I$ , any invariant of a binary  $p$ -ic, can be expressed in the form

$$A_1 I_1 + A_2 I_2 + A_3 I_3 + \dots,$$

where  $I_1, I_2, I_3, \dots$  are invariants, prove that it may be expressed in the form

$$A_1' I_1 + A_2' I_2 + A_3' I_3 + \dots,$$

where  $A_1', A_2', A_3', \dots$  are also invariants. (*Hilbert.*)

*Ans.* Operate with either form of generator of invariants, remembering that  $\Omega$  and  $O$  annihilate  $I, I_1, I_2, I_3, \dots$

## CHAPTER XI.

### FURTHER THEORY OF SEMINVARIANTS. THE BINARY QUANTIC OF INFINITE ORDER.

183.]  $\Omega$  expressed by means of roots. We commence this chapter by a consideration of the expression of seminvariants by means of the roots of a quantic, and more particularly by means of the sums of like powers of the roots, which might well have been given at a much earlier stage.

It was seen in § 91 that a seminvariant of

$$(a_0, a_1, a_2, \dots a_p)(x, 1)^p,$$

in virtue of its having

$$\Omega \equiv a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + \dots + pa_{p-1} \frac{d}{da_p} \quad \dots(1)$$

for an annihilator, is a function of  $a_0$  and the differences between roots. It is accordingly annihilated by

$$\Sigma \left( \frac{d}{da} \right) \equiv \frac{d}{da_1} + \frac{d}{da_2} + \dots + \frac{d}{da_p}, \quad \dots(2)$$

where  $a_1, a_2, \dots a_p$  are the roots. Moreover, any function of the roots which has this last for an annihilator is a function of their differences, by the ordinary theory of linear partial differential equations.

We can see as follows, what is thus suggested, that the effects on any function of the coefficients, which is therefore a function of  $a_0$  and the roots, of the operators (1) and (2) are identical, but for sign.

If

$$f(x) \equiv (a_0, a_1, a_2, \dots a_p)(x, 1)^p \equiv a_0(x-a_1)(x-a_2)\dots(x-a_p),$$

$$\frac{d}{da_1} f(x) \equiv -a_0(x-a_2)(x-a_3)\dots(x-a_p) \equiv -\frac{f(x)}{x-a_1}.$$

Therefore  $\Sigma \left( \frac{d}{da} \right) f(x) \equiv -\Sigma \frac{f(x)}{x-a} \equiv -f'(x).$

Now in this identity equate coefficients of the same powers of  $x$  on the left and right. It follows that

$$\begin{aligned}\Sigma\left(\frac{d}{da}\right)a_0 &= 0, \\ \Sigma\left(\frac{d}{da}\right)a_1 &= -a_0, \\ \Sigma\left(\frac{d}{da}\right)a_2 &= -2a_1, \\ &\dots\dots\dots \\ \Sigma\left(\frac{d}{da}\right)a_p &= -pa_{p-1}.\end{aligned}$$

Consequently, when operating on any function of

$$\begin{aligned}a_0, a_1, a_2, \dots a_p, \\ \Sigma\left(\frac{d}{da}\right) &\equiv \Sigma\left(\frac{d}{da}\right)a_0 \cdot \frac{d}{da_0} \\ &\quad + \Sigma\left(\frac{d}{da}\right)a_1 \cdot \frac{d}{da_1} + \dots + \Sigma\left(\frac{d}{da}\right)a_p \cdot \frac{d}{da_p} \\ &\equiv -a_0 \frac{d}{da_1} - 2a_1 \frac{d}{da_2} - \dots - pa_{p-1} \frac{d}{da_p} \\ &\equiv -\Omega.\end{aligned}$$

184.]  $\Omega$  expressed by means of sums of powers of roots. Now by Newton's formulæ for symmetric functions (Burnside and Panton, § 126),  $a_1, a_2, \dots a_p$  can be expressed, rationally and integrally, in terms of  $a_0$  and  $s_1, s_2, \dots s_p$ , the sums of the first, second, ...  $p$ th powers of the roots. Thus any rational integral function of  $a_0, a_1, a_2, \dots a_p$  may be expressed as a rational integral function of  $a_0$  and  $s_1, s_2, \dots s_p$ . Let us find the expression for  $\Sigma\left(\frac{d}{da}\right)$  which is suitable for operating with on functions so expressed.

We have at once  $\frac{d}{da_1} s_r = r a_1^{r-1}$ ,

so that  $\Sigma\left(\frac{d}{da}\right) s_r = r s_{r-1}$ .

In particular  $\Sigma\left(\frac{d}{da}\right) s_1 = p = s_0$ , say.



Also  $\Sigma \left( \frac{d}{da} \right) a_0 = 0$ . Thus, the operation being on a function of  $a_0$  and  $s_1, s_2, \dots, s_p$ ,

$$\begin{aligned} \Sigma \left( \frac{d}{da} \right) &\equiv \Sigma \left( \frac{d}{da} \right) a_0 \cdot \frac{d}{da_0} + \Sigma \left( \frac{d}{da} \right) s_1 \cdot \frac{d}{ds_1} \\ &\quad + \Sigma \left( \frac{d}{da} \right) s_2 \cdot \frac{d}{ds_2} + \dots + \Sigma \left( \frac{d}{da} \right) s_p \cdot \frac{d}{ds_p} \\ &\equiv s_0 \frac{d}{ds_1} + 2 s_1 \frac{d}{ds_2} + \dots + p s_{p-1} \frac{d}{ds_p}, \end{aligned}$$

where  $s_0 = p$ .

We conclude that

$$\begin{aligned} \Omega &\equiv a_0 \frac{d}{da_1} + 2 a_1 \frac{d}{da_2} + \dots + p a_{p-1} \frac{d}{da_p} \\ &\equiv - \left\{ s_0 \frac{d}{ds_1} + 2 s_1 \frac{d}{ds_2} + \dots + p s_{p-1} \frac{d}{ds_p} \right\}, \end{aligned}$$

i.e. that the  $\Omega$  operator is identical in form, but for sign, when expressed in form for operation on a function of  $a_0$  and the  $s$ 's, as when expressed in form for operation on the equivalent function of  $a_0, a_1, a_2, \dots, a_p$ . Note however the absence of  $a_0$  from the  $s$ -form of  $\Omega$ .

We gather then that, when a function of  $a_0, a_1, a_2, \dots, a_p$  is a seminvariant, so is the same function of  $s_0$ , i.e.  $p$ , and  $s_1, s_2, \dots, s_p$ . The latter function when expressed in terms of the  $a$ 's will of course have a negative power of  $a_0$  as factor, since the  $s$ 's are functions of the ratios of the  $a$ 's to  $a_0$ .

The idea of this duality appears to be due to M. Roberts.

Ex. 1. If a homogeneous isobaric function of degree  $i$ , in the coefficients, and weight  $w$ , be called  $\phi_1$  when expressed in terms of  $a_0, a_1, a_2, \dots, a_p$ , and  $\phi_2$  when expressed in terms of  $a_0$  and the roots,  $\phi_3$  when expressed in terms of  $a_0$  and  $s_1, s_2, \dots, s_p$ , and  $\phi$  when no particular expression is necessarily implied, prove that

$$i\phi = \left( a_0 \frac{d}{da_0} + a_1 \frac{d}{da_1} + \dots + a_p \frac{d}{da_p} \right) \phi_1 = a_0 \frac{d}{da_0} \phi_2 = a_0 \frac{d}{da_0} \phi_3.$$

Ex. 2. With the same notation

$$\begin{aligned} w\phi &= \left( a_1 \frac{d}{da_1} + 2 a_2 \frac{d}{da_2} + \dots + p a_p \frac{d}{da_p} \right) \phi_1 = \Sigma \left( a \frac{d}{da} \right) \phi_2 \\ &= \left( s_1 \frac{d}{ds_1} + 2 s_2 \frac{d}{ds_2} + \dots + p s_p \frac{d}{ds_p} \right) \phi_3. \end{aligned}$$

~~Ex. 1~~  
 $a_0$  appears  
 4 or 5 of  
 as fact

Ex. 3. With the same notation

$$\begin{aligned} \left\{ O - ip \frac{a_1}{a_0} \right\} \phi &= \left\{ \left[ pa_1 \frac{d}{da_0} + (p-1)a_2 \frac{d}{da_1} + \dots + a_p \frac{d}{da_{p-1}} \right] \right. \\ &\quad \left. - p \frac{a_1}{a_0} \left[ a_0 \frac{d}{da_0} + a_1 \frac{d}{da_1} + \dots + a_p \frac{d}{da_p} \right] \right\} \phi_1 \\ &= \Sigma \left( a^2 \frac{d}{da} \right) \phi_2 = \left\{ s_2 \frac{d}{ds_1} + 2s_3 \frac{d}{ds_2} + \dots + ps_{p+1} \frac{d}{ds_p} \right\} \phi_3, \end{aligned}$$

which by Ex. 1 may also be written

$$\begin{aligned} O\phi &= \left[ pa_1 \frac{d}{da_0} + (p-1)a_2 \frac{d}{da_1} + \dots + a_p \frac{d}{da_{p-1}} \right] \phi_1 \\ &= \left\{ \Sigma \left( a^2 \frac{d}{da} \right) - \Sigma(a) a_0 \frac{d}{da_0} \right\} \phi_2 \\ &= \left\{ -s_1 a_0 \frac{d}{da_0} + s_2 \frac{d}{ds_1} + 2s_3 \frac{d}{ds_2} + \dots + ps_{p+1} \frac{d}{ds_p} \right\} \phi_3. \end{aligned}$$

*Ans.* Prove the  $\phi_1, \phi_2$  equality by considering that, as in § 183, the effect of  $-O$  is that of  $\Sigma \left( \frac{d}{d\frac{1}{a}} \right)$  on the function expressed in terms of  $a_p$  and  $a_1, a_2, \dots, a_p$ .

Ex. 4. By means of the expressions for the operators in terms of  $a_0$  and roots prove the known equivalence of an operator and a multiplier

$$\Omega O - O \Omega = ip - 2w.$$

Ex. 5. By Leibnitz' theorem prove (§ 125, Ex. 1) that

$$\begin{aligned} \Omega^r O - O \Omega^r &\equiv (-1)^r \left\{ \left( \Sigma \frac{d}{da} \right)^r \left[ \Sigma \left( a^2 \frac{d}{da} \right) - \Sigma(a) a_0 \frac{d}{da_0} \right] \right. \\ &\quad \left. - \left[ \Sigma \left( a^2 \frac{d}{da} \right) - \Sigma(a) a_0 \frac{d}{da_0} \right] \left( \Sigma \frac{d}{da} \right)^r \right\} \\ &= r(ip - 2w + r - 1) \Omega^{r-1}. \end{aligned}$$

Ex. 6. By means of the equivalence of

$$\Sigma \left( \frac{d}{da} \right) \text{ and } s_0 \frac{d}{ds_1} + 2s_1 \frac{d}{ds_2} + \dots + ps_{p-1} \frac{d}{ds_p},$$

prove that, if  $s_{p+1}, s_{p+2}, \dots$  be regarded as functions of  $s_1, s_2, \dots, s_p$ ,

$$- \Omega s_{p+r} = \left( s_0 \frac{d}{ds_1} + 2s_1 \frac{d}{ds_2} + \dots + ps_{p-1} \frac{d}{ds_p} \right) s_{p+r} = (p+r) s_{p+r-1}.$$

Ex. 7. By means of Ex. 3 prove that, whether  $m$  be less than equal to or greater than  $p$ ,

$$ms_{m+1} = \left( s_2 \frac{d}{ds_1} + 2s_3 \frac{d}{ds_2} + \dots + ps_{p+1} \frac{d}{ds_p} \right) s_m = 0 s_m.$$

Ex. 8. Also, even when  $m$  exceeds  $p$ ,

$$ms_{m-1} = \left( s_0 \frac{d}{ds_1} + 2s_1 \frac{d}{ds_2} + \dots + ps_{p-1} \frac{d}{ds_p} \right) s_m = -\Omega s_m.$$

185.] It must be borne in mind that two distinct things may be meant by the expression of a function of  $a_0, a_1, a_2, \dots, a_p$  in terms of  $a_0$  and the sums of the powers of the roots. We may mean the expression in terms of  $a_0$  and  $s_1, s_2, s_3, \dots, s_p$  only. This expression is in all cases unique, and, when the function of  $a_0, a_1, a_2, \dots, a_p$  is rational and integral and of weight not greater than  $p$ , is the only expression. When, however, the weight exceeds  $p$ , there will be, as a rule, also other expressions involving  $s_{p+1}, s_{p+2}, \dots$  or some of them as well as lower sums. The above articles contemplate the unique expression obtained from any correct expression by giving in it to  $s_{p+1}, s_{p+2}, \dots$  their values in terms of  $s_1, s_2, \dots, s_p$ .

We shall have occasion presently to consider the binary quantic of infinite order, in which the series  $a_0, a_1, a_2, a_3, \dots$  of coefficients is unending. In its case the distinction does not arise.

Ex. 9. Prove that, if a homogeneous isobaric function  $\phi$  of  $a_0, a_1, a_2, \dots, a_p$  be expressed in any manner in terms of  $a_0$  and  $s_1, s_2, \dots, s_p, s_{p+1}, \dots$ , and when so expressed be called  $\phi_4$ ,

$$(1) \Omega \phi = - \left\{ s_0 \frac{d}{ds_1} + 2s_1 \frac{d}{ds_2} + \dots \text{to } \infty \right\} \phi_4,$$

$$(2) i\phi = a_0 \frac{d}{da_0} \phi_4,$$

$$(3) w\phi = \left\{ s_1 \frac{d}{ds_1} + 2s_2 \frac{d}{ds_2} + \dots \text{to } \infty \right\} \phi_4,$$

$$(4) \left\{ O - ip \frac{a_1}{a_0} \right\} \phi = \left\{ s_2 \frac{d}{ds_1} + 2s_3 \frac{d}{ds_2} + \dots \text{to } \infty \right\} \phi_4.$$

186.] Completion of § 174. It is interesting to gather from § 183, and the examples which follow § 184, the following conclusions

$$-\Omega = \Sigma \left( \frac{d}{da} \right),$$

$$w - \frac{a_1}{a_0} \Omega = \Sigma \left\{ \left( a + \frac{a_1}{a_0} \right) \frac{d}{da} \right\},$$

$$O - \frac{a_1}{a_0} (ip - 2w) - \frac{a_1^2}{a_0^2} \Omega = \Sigma \left\{ \left( a + \frac{a_1}{a_0} \right)^2 \frac{d}{da} \right\},$$

which, if we put  $a'$  for  $a + \frac{a_1}{a_0}$  and similarly as to all the roots, i. e. if we transform  $(a_0, a_1, a_2, \dots a_p)(x, 1)^p$  into

$$(a_0, 0, a_2', \dots a_p')(x + \frac{a_1}{a_0}, 1)^p,$$

a form without its second term, may be written

$$-\Omega = \Sigma \left( \frac{d}{da'} \right) = -\Omega',$$

$$w - \frac{a_1}{a_0} \Omega = \Sigma \left( a' \frac{d}{da'} \right) = w',$$

$$O - \frac{a_1}{a_0} (ip - 2w) - \frac{a_1^2}{a_0^2} \Omega = \Sigma \left( a'^2 \frac{d}{da'} \right) = O',$$

the remainders of the right-hand sides vanishing since  $a_1' = 0$ .

We must not lose sight here of the tacit assumption made that the function operated on can be expressed in terms of  $a_0$  and  $a_1 + \frac{a_1}{a_0}$ ,  $a_2 + \frac{a_1}{a_0}$ , ...  $a_p + \frac{a_1}{a_0}$ , which we have called  $a_1', a_2', \dots a_p'$ . The functions which can be so expressed are, we know, seminvariants. The second equation really exhibits the fact anew. The weights  $w$  and  $w'$  are equal. Consequently that equation gives  $\Omega = 0$ , i. e. that the function operated on is a seminvariant.

In these equalities  $\Omega'$ ,  $O'$  and  $w'$  (regarded as an operator) are the same operators in  $a_0, a_1' (= 0), a_2', \dots a_p'$  as  $\Omega, O$  and  $w$  are in  $a_0, a_1, a_2, \dots a_p$ . They contain the symbol  $\frac{d}{da_1}$ , whose

meaning is not obvious, but is really defined by the first of the equalities.

This symbol may be eliminated by subtracting  $(p-1)a_2'$  times  $\Omega'$  from  $a_0$  times  $O'$ . We have, in fact, since  $a_2' = a_2 - \frac{a_1^2}{a_0}$ ,

$$a_0 O' - (p-1)a_2' \Omega' = a_0 O - a_1(ip-2w) - \left\{ (p-1)a_2 - (p-2)\frac{a_1^2}{a_0} \right\} \Omega.$$

We may take this in connexion with § 174. The operator on the left is the  $\psi'$  of that article. We have, in fact, here before us the materials for the proof of the converse proposition there stated, that every gradient in  $a_0, a_2', a_3', \dots, a_p'$  for which  $ip-2w=0$  and which has  $\psi'$  for an annihilator is an invariant, i.e. is annihilated by  $O$  as well as by  $\Omega$ , which last must annihilate it as it does any function of  $a_0, a_2', a_3', \dots, a_p'$ . We remember that the facts of being annihilated by  $\Omega$ , and being of properly connected degree and weight, were not sufficient to assure us of its invariancy in default of evidence either that it had  $O$  for an annihilator or that it was integral in  $a_0$ . We now see that as  $\Omega$  and  $\psi'$  annihilate it, and as the relation  $ip=2w$  holds, the annihilation by  $a_0 O$ , and therefore by  $O$ , follows.

Ex. 10. Prove that  $a_0 O - a_1(ip-2w)$  cannot annihilate any function which is fractional in  $a_0$  and for which  $ip-2w$  is zero or positive.

Ans. The terms of highest degree in  $a_0^{-1}$  in the expression of the annihilation of  $P a_0^{-\mu} + Q a_0^{-(\mu-1)} + \dots + T$  must vanish. This gives  $P=0$ . So  $Q=0$ , &c.

Ex. 11. Any gradient in  $a_0, a_1, a_2, \dots, a_p$  for which  $ip-2w$  is positive and which is annihilated by  $a_0 O - a_1(ip-2w)$ , is the product of a power of  $a_0$  and a gradient which it annihilates and for which

$$ip-2w=0.$$

Ex. 12. Hence  $\psi'$  only annihilates gradients in  $a_0, a_2', a_3', \dots, a_p'$  which are invariants or invariants multiplied by powers of  $a_0$ .

187.] Partial differentiation with regard to  $s_1, s_2, \dots, s_p$ . Let us replace  $a_0, p a_1, \frac{p(p-1)}{1 \cdot 2} a_2, \dots, a_p$  by  $c_0, c_1, c_2, \dots, c_p$ , and also replace  $x:1$  by  $1:y$ , thus considering the  $p$ -ic without binomial coefficients

$$c_0 + c_1 y + c_2 y^2 + \dots + c_p y^p \equiv c_0 (1 - a_1 y) (1 - a_2 y) \dots (1 - a_p y).$$

Taking logarithms we have at once

$$\begin{aligned} \log (c_0 + c_1 y + c_2 y^2 + \dots + c_p y^p) \\ = \log c_0 - s_1 y - s_2 \frac{y^2}{2} - s_3 \frac{y^3}{3} - \dots - s_p \frac{y^p}{p} - s_{p+1} \frac{y^{p+1}}{p+1} - \dots \end{aligned}$$

Now regard  $c_1, c_2, \dots, c_p$ , and also  $s_{p+1}, s_{p+2}, \dots$  as functions of  $c_0$  and  $s_1, s_2, \dots, s_p$ . Partial differentiation with regard to  $s_r$ , where  $r < 1$  and  $\succ p$ , gives us the identity

$$\begin{aligned} \frac{dc_1}{ds_r} y + \frac{dc_2}{ds_r} y^2 + \dots + \frac{dc_p}{ds_r} y^p = (c_0 + c_1 y + c_2 y^2 + \dots + c_p y^p) \\ \left\{ -\frac{y^r}{r} - \frac{ds_{p+1}}{ds_r} \frac{y^{p+1}}{p+1} - \frac{ds_{p+2}}{ds_r} \frac{y^{p+2}}{p+2} - \dots \right\} \end{aligned}$$

which holds for all values of  $y$ . Equating corresponding coefficients on the two sides we have

$$\frac{dc_m}{ds_r} = 0, \text{ if } m < r, \text{ and}$$

$$\frac{dc_m}{ds_r} = -\frac{1}{r} c_{m-r}, \text{ if } m \text{ lie between } r \text{ and } p \text{ inclusive.}$$

The other equations, given by the terms in  $y^{p+1}, y^{p+2}, \dots$ , determine for us  $\frac{ds_{p+1}}{ds_r}, \frac{ds_{p+2}}{ds_r}, \dots$

Hence, if the operations on the right be on a function of  $c_0, c_1, c_2, \dots, c_p$ , and that on the left be on the equivalent of that function in terms of  $c_0$  and  $s_1, s_2, \dots, s_p$ ,

$$\begin{aligned} \frac{d}{ds_r} = \frac{dc_0}{ds_r} \cdot \frac{d}{dc_0} + \frac{dc_1}{ds_r} \cdot \frac{d}{dc_1} + \frac{dc_2}{ds_r} \cdot \frac{d}{dc_2} + \dots + \frac{dc_p}{ds_r} \cdot \frac{d}{dc_p} \\ = -\frac{1}{r} \left\{ c_0 \frac{d}{dc_r} + c_1 \frac{d}{dc_{r+1}} + \dots + c_{p-r} \frac{d}{dc_p} \right\}. \end{aligned}$$

In particular, taking  $r = 1$ , we have

$$\frac{d}{ds_1} = - \left\{ c_0 \frac{d}{dc_1} + c_1 \frac{d}{dc_2} + c_2 \frac{d}{dc_3} + \dots + c_{p-1} \frac{d}{dc_p} \right\}.$$

Let us here revert to our first notation, replacing

$$c_0, c_1, c_2, \dots, c_p \text{ by } a_0, pa_1, \frac{p(p-1)}{1 \cdot 2} a_2, \dots, a_p,$$

so that  $s_1$ , &c., are the sums of the powers of the roots of

$$(a_0, a_1, a_2, \dots a_p)(x, 1)^p = 0.$$

The result obtained is that

$$\frac{d}{ds_1} = - \left\{ \frac{1}{p} a_0 \frac{d}{da_1} + \frac{2}{p-1} a_1 \frac{d}{da_2} + \frac{3}{p-2} a_2 \frac{d}{da_3} + \dots + \frac{p}{1} a_{p-1} \frac{d}{da_p} \right\}.$$

188.]  $\Omega$  as an annihilator of non-unitary symmetric functions. A more instructive conclusion is, however, derived by replacing  $c_0, c_1, c_2, \dots c_p$  by  $b_0, \frac{b_1}{1!}, \frac{b_2}{2!}, \frac{b_3}{3!}, \dots$ , so that the equation of which  $s_1, s_2, \dots s_p$  are the first  $p$  sums of the powers of the roots is

$$b_0 x^p + \frac{b_1}{1!} x^{p-1} + \frac{b_2}{2!} x^{p-2} + \dots + \frac{b_p}{p!} = 0.$$

With this notation we obtain that

$$- \frac{d}{ds_1} = b_0 \frac{d}{db_1} + 2 b_1 \frac{d}{db_2} + 3 b_2 \frac{d}{db_3} + \dots + p b_{p-1} \frac{d}{db_p},$$

whose right-hand side is of the well-known form of  $\Omega$ .

Our conclusion hence is that the seminvariants of

$$a_0 x^p + p a_1 x^{p-1} y + \frac{p(p-1)}{1.2} a_2 x^{p-2} y^2 + \dots + a_p y^p$$

are identical, but for the factor  $a_0^i$  where  $i$  is the degree in each case, with those symmetric functions of the roots of

$$a_0 x^p + \frac{a_1}{1!} x^{p-1} + \frac{a_2}{2!} x^{p-2} + \dots + \frac{a_p}{p!} = 0$$

which when expressed in terms of  $s_1, s_2, \dots s_p$  are free from  $s_1$ .

Reference is made to works on the Theory of Equations for the fact that a symmetric function

$$\Sigma . a_1^l a_2^m a_3^n \dots,$$

of which  $l+m+n+\dots$  is the weight, and the greatest of  $l, m, n, \dots$  is the least number which can be taken for  $i$  that

upon multiplication by  $a_0^i$  it may become integral in the coefficients, may be written as a rational integral function of

$$\begin{aligned} & s_l, s_m, s_n, \dots, \\ & s_{l+m}, s_{l+n}, s_{m+n}, \dots, \\ & s_{l+m+n}, \dots, \\ & \dots \end{aligned}$$

so that, if none of  $l, m, n, \dots$  is unity, and if the weight  $l + m + n + \dots$  does not exceed  $p$ ,  $s_1$  does not occur, and a seminvariant is thus obtained.

If, however, either of  $l, m, n, \dots$  is unity no seminvariant is given.

If, even though this be not the case, the weight  $l + m + n + \dots$  exceeds  $p$ , then, though the expression in terms of  $a_0$  and  $s_1, s_2, s_3, \dots, s_{l+m+n+\dots}$  does not involve  $s_1$ , the same cannot be said necessarily or as a rule of the expression in terms of  $a_0$  and  $s_1, s_2, \dots, s_p$ , for  $s_{p+1}$ , &c., expressed in terms of  $s_1, s_2, \dots, s_p$ , are not free from  $s_1$  by § 187.

When  $p$  is infinite the case of  $l + m + n + \dots$  exceeding  $p$  does not arise. Thus in this limiting case  $s_1$  does not occur in the symmetric function  $\Sigma. a_1^l a_2^m a_3^n \dots$  unless one or more of  $l, m, n, \dots$  is unity, and does otherwise.

The seminvariants of  $(a_0, a_1, a_2, \dots)(x, y)^p$ , when  $p$  is infinite, are then what are called the 'non-unitary' symmetric functions of  $a_1, a_2, a_3, \dots$ , where

$$a_0 + \frac{a_1}{1!}y + \frac{a_2}{2!}y^2 + \dots \text{ to } \infty \equiv a_0(1 - a_1y)(1 - a_2y)(1 - a_3y) \dots,$$

each multiplied by  $a_0^i$ , where  $i$  is the degree, i.e. is the greatest of the indices  $l, m, n, \dots$  in the typical product of roots summed, or any greater number.

It is from this point of view that MacMahon has discussed the concomitants of the binary quantic of infinite order.

It will of course be remembered that a seminvariant, of a quantic of any order  $(a_0, a_1, a_2, \dots, a_p)(x, y)^p$ , which is only of extent  $r$ , i.e. which involves only  $a_0, a_1, a_2, \dots, a_r$ , is equally a seminvariant of each of the lower quantics

$$(a_0, a_1, a_2, \dots, a_r)(x, y)^r, (a_0, a_1, a_2, \dots, a_r, a_{r+1})(x, y)^{r+1}, \dots$$



Thus, in particular, when we have a seminvariant of a binary quantic of infinite order, we have in it a seminvariant of a binary quantic whose order is the extent of the seminvariant and one of every order higher than this extent. If, in fact,

$$a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + 3a_2 \frac{d}{da_3} + \dots \text{ to } \infty$$

annihilates  $S(a_0, a_1, a_2, \dots a_r)$ , then equally do

$$a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + 3a_2 \frac{d}{da_3} + \dots + ra_{r-1} \frac{d}{da_r},$$

$$a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + 3a_2 \frac{d}{da_3} + \dots + ra_{r-1} \frac{d}{da_r} + (r+1)a_r \frac{d}{da_{r+1}},$$

&c., &c.

Ex. 13. Prove that, for any positive integral value of  $p$ ,  
 $(a_0, a_1, a_2, \dots a_p) (x, 1)^p$

$$= \left( a_0 + \frac{a_1}{1} \frac{d}{dx} + \frac{a_2}{2!} \frac{d^2}{dx^2} + \frac{a_3}{3!} \frac{d^3}{dx^3} + \dots \text{ to } \infty \right) x^p.$$

Ex. 14. Prove that all the coefficients but that of  $t$  in the expansion in powers of  $t$  of

$$\log \left( a_0 + \frac{a_1}{1} t + \frac{a_2}{2!} t^2 + \frac{a_3}{3!} t^3 + \dots \right) - \log a_0$$

are seminvariants in the letters  $a_0, a_1, a_2, a_3, \dots$ , fractional in  $a_0$ .

189.] **Generating Functions. Perpetuants.** Complete tables of symmetric functions have been calculated up to the weight 14: for weights 1 to 10 by Meyer Hirsch (cf. Notes to Salmon's *Higher Algebra*): for weight 11 by Faa de Bruno (cf. his *Formes Binaires*): for weights 12 and 14 by Durfee (*Am. Journal*, Vols. V, IX): and for weight 13 by MacMahon (*Am. Journal*, Vol. VI). Thus a complete set of seminvariants up to weight 14, of which all seminvariants whatever up to that weight are linear functions, is known.

It has been seen (§ 135) that the number of linearly independent seminvariants of weight  $w$  degree  $i$  and extent  $p$  or less is the coefficient of  $x^w$  in the development of

$$\frac{(1-x^{p+1})(1-x^{p+2}) \dots (1-x^{p+i})}{(1-x^2)(1-x^3) \dots (1-x^i)}.$$

Here make  $p = \infty$ . It follows that the whole number of linearly independent seminvariants of weight  $w$  and degree  $i$  of the quantic of infinite order, or of a quantic of order not less than the weight  $w$ , is

$$\text{co. } x^w \text{ in developement of } \frac{1}{(1-x^2)(1-x^3)\dots(1-x^i)}.$$

We may want also the number of linearly independent (or aszygetic) seminvariants of degree-weight  $i, w$  which are aszygetic with seminvariants of the same weight and lower degrees multiplied by powers of  $a_0$ . This number may be found by subtracting from the number of weight  $w$  and degree  $i$  the number of weight  $w$  and degree  $i-1$ . Thus it is

$$\begin{aligned} \text{co. } x^w \text{ in developement of } & \frac{1}{(1-x^2)(1-x^3)\dots(1-x^{i-1})(1-x^i)} \\ & - \frac{1}{(1-x^2)(1-x^3)\dots(1-x^{i-1})} \\ = \text{ " " " } & \frac{x^i}{(1-x^2)(1-x^3)\dots(1-x^i)}. \end{aligned}$$

The same generating function is given by MacMahon's theory of non-unitary symmetric functions. The non-unitary symmetric functions which give such seminvariants are of the form  $\Sigma . a_1^l a_2^m a_3^n \dots$ , where one at least of the indices  $l, m, n, \dots$  is  $i$ , and where, as in general, none of them is unity or greater than  $i$ , and their sum is the weight. Now the type-products of weight  $w$ , whose summations give such symmetric functions, are in number equal to the number of ways in which  $w-i$  may be made up of  $i-1$  or fewer numbers chosen from  $2, 3, \dots, i$ , i. e. to the coefficient of  $x^{w-i}$  in the expansion of the product

$$(1+x^2+x^4+\dots)(1+x^3+x^6+\dots)\dots(1+x^i+x^{2i}+\dots),$$

i. e. to the coefficient of  $x^w$  in  $x^i$  times this product, which is the developement of

$$\frac{x^i}{(1-x^2)(1-x^3)\dots(1-x^i)}.$$

The problem of the enumeration of the irreducible seminvariants of the binary quantic of infinite order is one which admits of solution. Indeed it has been solved by MacMahon by an analysis of non-unitary partitions, and his conclusions

have been fully confirmed symbolically by Stroh. Irreducible seminvariants of the quantic of infinite order are called *perpetuants*, the name being Sylvester's. A perpetuant is a seminvariant which cannot be expressed rationally and integrally in terms of other perpetuants of lower degree. Of the first degree there is one perpetuant  $a_0$ . Of the second degree there is one perpetuant of each even weight, viz.  $a_0a_2 - a_1^2$ ,  $a_0a_4 - 4a_1a_3 + 3a_2^2$ ,  $a_0a_6 - 6a_1a_5 + 15a_2a_4 - 10a_3^2$ , .... For any higher degree  $i$  the number of perpetuants of weight  $w$  is the coefficient of  $x^w$  in the developement of the generating function

$$\frac{x^{2^i-1}-1}{(1-x^2)(1-x^3)\dots(1-x^i)}.$$

The mistaken idea must not be entertained that when we know the perpetuants of extent  $p$  or less, i. e. the irreducible seminvariants of extent  $p$  or less of the quantic of infinite order, we know the irreducible seminvariants or ground forms of a  $p$ -ic. This is not the case. There may be seminvariants of extent  $p$  or less, which are not capable of rational integral expression in terms of lower seminvariants of extent  $p$  or less, but which are in terms of seminvariants of lower degree and extents some of which exceed  $p$ . One instance which we have met with will suffice to illustrate this. We have found (§ 169) that the seminvariant  $(ad - bc)^2 - 4(ac - b^2)(bd - c^2)$  is irreducible when we are confined to extent 3, being an irreducible invariant of the cubic. But (§ 170) when we proceed to extent 4 it is no longer irreducible, being capable of being written

$$(ac - b^2)(ae - 4bd + 3c^2) - a(ace + 2bcd - ad^2 - b^2e - c^3).$$

It is an irreducible invariant of the cubic, but is reducible for quantics of higher order, and so is not a perpetuant.

For a synopsis of Stroh's method of investigation (*Mathematische Annalen*, Vol. XXXVI), see Exx. 35 to 42 at the end of the present chapter.

The theory of perpetuants has been recently completed by MacMahon, who has investigated expressions for them in the notation of partitions. (*Proc. Lond. Math. Soc.* 1895.)

190.] **Reciprocity.** By Hermite's law of reciprocity there must be a strictly correlative theory to much of the above in

which the ideas of degree and extent are interchanged. It concerns seminvariants of infinite degree, or of degree not less than weight, just as the above theory concerns seminvariants of a quantic of infinite order, or of order not less than the weight of the seminvariants in question.

The number of aszygetic seminvariants, for which  $i$  is not less than  $w$ , of a  $p$ -ic is thus the coefficient of  $x^w$  in the developement of

$$\frac{1}{(1-x^2)(1-x^3)\dots(1-x^p)};$$

and the number of aszygetic seminvariants of a  $p$ -ic, for which  $i$  is not less than  $w$  and which are really of extent  $p$ , so as not to belong to a  $(p-1)$ -ic equally, is the coefficient of  $x^w$  in the developement of

$$\frac{x^p}{(1-x^2)(1-x^3)\dots(1-x^p)}.$$

These facts may be independently arrived at. By § 163 the seminvariants in question are rational integral functions of  $A_2, A_3, \dots, A_p$ , i. e.  $a_0 a_2', a_0^2 a_3', \dots, a_0^{p-1} a_p'$ , raised to the requisite excess of degree over weight by the power  $a_0^{i-w}$  of  $a_0$  as factor.

The theory dual to that of perpetuants is not so obvious.

191.] **Power ending products.** It has been seen in the last article that the

$$(w; \infty, p) - (w-1; \infty, p)$$

aszygetic seminvariants of a  $p$ -ic whose weight is  $w$  and whose degree is a definite number not less than  $w$  have a one to one correspondence with the

$$(w; \infty, p) - (w-1; \infty, p)$$

products of weight  $w$  of  $a_0, a_2, a_3, \dots, a_p$ , i. e. with the 'non-unitary' partitions of  $w$ .

It is by reciprocity suggested as probable that there is a system of

$$(w; i, \infty) - (w-1; i, \infty)$$

partitions of  $w$  into  $i$  or fewer parts, i. e. a system of this number of products of weight  $w$  of  $i$  of  $a_0, a_1, a_2, a_3, \dots, a_p$ , to which there is a one to one correspondence of the

$$(w; i, \infty) - (w-1; i, \infty)$$

asyzygetic seminvariants of weight  $w$  and degree  $i$  of the  $p$ -ic, when  $p$  is infinite or not less than  $w$ .

It is also suggested that those partitions of  $w$ , or those products, are the partitions or products which are exhibited in § 130 by aid of Ferrers' diagrams as the reciprocals of non-unitary partitions or products.

Now if we write down the diagram of a non-unitary product, we see that the absence of  $a_1$  in the product is exhibited by the fact that the first two columns at least in the diagram contain equal numbers of dots.

This tells us that in the reciprocal product the letter of highest suffix which occurs is present to a higher power than the first.

Products of this class are called by MacMahon and Cayley *power ending products* or *power enders*. Let us adopt the notation  $a, b, c, \dots$  of alphabetical sequence instead of the notation  $a_0, a_1, a_2, \dots$  of numerical sequence. *Power enders* are those products of some of  $a, b, c, \dots$  which, when their factors are alphabetically arranged from left to right, end in a higher power than the first. Thus  $a^2, ab^2, abd^3, c^3, \dots$  are power enders, while  $a, a^2b, ab^2d, c, \dots$  are not.

The whole number of products of weight  $w$  of  $i$  of  $a, b, c, \dots$  is  $(w; i, \infty)$ , and the whole number of non-power enders is the number of products which can be derived from products of degree  $i$  and weight  $w-1$  by replacing the last letter in each, once only, by the next more advanced letter, i.e. is  $(w-1; i, \infty)$ . The number of power enders of the type is then

$$(w; i, \infty) - (w-1; i, \infty).$$

We shall see that there is the expected one to one correspondence of these products with a complete system of  $(w; i, \infty) - (w-1; i, \infty)$  asyzygetic seminvariants of weight  $w$  and degree  $i$ ; in fact, that the latter complete system may be derived, one from one, by differential operations on the former complete system.

192.] **An annihilator of all gradients.** Let us refer back to § 180, and proceed to the limit when  $p$ , the order of the quantic there under consideration, or the extent of  $\Omega$ , is infinite. Remember, too, that though we consider the quantic

of infinite order we deal with gradients of finite weight, involving consequently only a finite series of the letters  $a, b, c, d, \dots$ .

We have now

$$\Omega \equiv a\partial_b + 2b\partial_c + 3c\partial_a + \dots \text{ to } \infty,$$

where  $\partial_k$  as usual denotes  $\frac{d}{dk}$ . We have also

$$O \equiv pb\partial_a + (p-1)c\partial_b + (p-2)d\partial_c + \dots,$$

where  $p$  is infinite,

$$\begin{aligned} &\equiv p\{b\partial_a + c\partial_b + d\partial_c + \dots\} - \{c\partial_b + 2d\partial_c + 3e\partial_a + \dots\} \\ &\equiv p\mathfrak{S} - \phi, \text{ say,} \end{aligned}$$

where the result of operating with  $\phi$  on a gradient of finite extent vanishes in comparison with that of operating with the infinite  $p\mathfrak{S}$ .

We have also  $\eta = ip - 2w$ ,

which is infinite, and consequently always positive,  $w$  being finite.

The limiting form for  $p$  infinite taken by the operator

$$1 - \frac{1}{1 \cdot (\eta + 2)} O \Omega + \frac{1}{1 \cdot 2 \cdot (\eta + 2)(\eta + 3)} O^2 \Omega^2 - \dots$$

is hence at once seen to be

$$1 - \frac{1}{i} \mathfrak{S} \Omega + \frac{1}{i^2} \frac{\mathfrak{S}^2 \Omega^2}{1 \cdot 2} - \frac{1}{i^3} \cdot \frac{\mathfrak{S}^3 \Omega^3}{1 \cdot 2 \cdot 3} + \dots$$

This then, by § 180, operating on any gradient in  $a, b, c, d, \dots$ , and in particular of course on any single product, produces either zero or a seminvariant of the quantic of infinite order, or, as is the same thing, of a quantic of order not less than  $w$ , the weight of the gradient.

193.] It is really most convincing and easiest to prove this independently, and not deduce it as the limit of something else. The student will have no difficulty in proving by the method of §§ 123, &c., that,  $G$  being any gradient of degree  $i$ ,

$$\begin{aligned} (\Omega \mathfrak{S} - \mathfrak{S} \Omega) G &= iG, \\ (\Omega \mathfrak{S}^2 - \mathfrak{S}^2 \Omega) \Omega G &= 2i \mathfrak{S} \Omega G, \\ (\Omega \mathfrak{S}^3 - \mathfrak{S}^3 \Omega) \Omega^2 G &= 3i \mathfrak{S}^2 \Omega^2 G, \\ &\text{\&c., \&c.,} \end{aligned}$$

and by addition of properly chosen multiples of these we obtain the identity

$$\left\{ 1 - \frac{1}{i} \cdot \frac{\Omega \mathcal{S}}{1} + \frac{1}{i^2} \cdot \frac{\Omega \mathcal{S}^2 \Omega}{1 \cdot 2} - \frac{1}{i^3} \cdot \frac{\Omega \mathcal{S}^3 \Omega^2}{1 \cdot 2 \cdot 3} + \dots \right\} G = 0,$$

in which the series practically terminates, since  $\Omega^{w+1} G = 0$ , so that no doubt arising from questions of convergency presents itself.

This tells us first that any gradient  $G$  is of the form  $\Omega F$ , when we allow the extent of  $F$  to be greater than that of  $G$ , and  $\Omega$  to be non-terminating. The limitation imposed by the requirement of  $ip - 2w$  to be positive in § 128 does not of course arise,  $p$  being now infinite.

The result we need follows by putting  $\Omega F$ , where  $F$  is any gradient of degree  $i$ , for  $G$ . This gives us that

$$\Omega \left\{ 1 - \frac{1}{i} \frac{\mathcal{S} \Omega}{1} + \frac{1}{i^2} \frac{\mathcal{S}^2 \Omega^2}{1 \cdot 2} - \frac{1}{i^3} \frac{\mathcal{S}^3 \Omega^3}{1 \cdot 2 \cdot 3} + \dots \right\} F = 0,$$

so that, as in the last article,

$$1 - \frac{1}{i} \frac{\mathcal{S} \Omega}{1} + \frac{1}{i^2} \frac{\mathcal{S}^2 \Omega^2}{1 \cdot 2} - \frac{1}{i^3} \frac{\mathcal{S}^3 \Omega^3}{1 \cdot 2 \cdot 3} + \dots$$

produces from any gradient of degree  $i$  a seminvariant or zero.

Ex. 15. Prove in like manner that,  $\phi$  being as in § 192,

$$\begin{aligned} (\Omega \phi - \phi \Omega) G &= 2wG, \\ (\Omega \phi^2 - \phi^2 \Omega) \Omega G &= 2(2w-1)\phi \Omega G, \\ (\Omega \phi^3 - \phi^3 \Omega) \Omega^2 G &= 3(2w-2)\phi^2 \Omega^2 G, \\ &\text{\&c. \&c.}, \end{aligned}$$

and hence that

$$\begin{aligned} 1 - \frac{1}{2w-2} \frac{\phi \Omega}{1} + \frac{1}{(2w-2)(2w-3)} \frac{\phi^2 \Omega^2}{1 \cdot 2} \\ - \frac{1}{(2w-2)(2w-3)(2w-4)} \frac{\phi^3 \Omega^3}{1 \cdot 2 \cdot 3} - \dots, \text{ to } w+1 \text{ terms,} \end{aligned}$$

produces a seminvariant or a zero from every gradient of weight 2 or more.

194.] Two generators of all seminvariants. One to one correspondence of seminvariants and power ends. Now a gradient  $F$  which

$$1 - \frac{1}{i} \frac{\mathcal{S} \Omega}{1} + \frac{1}{i^2} \frac{\mathcal{S}^2 \Omega^2}{1 \cdot 2} - \dots$$

annihilates is of the form  $\mathfrak{S}F'$ , for the fact of annihilation gives us

$$F = \mathfrak{S} \left\{ \frac{1}{i} \Omega - \frac{1}{i^2} \frac{\mathfrak{S} \Omega^2}{1 \cdot 2} + \dots \right\} F.$$

If then we can be sure that no power ending product can be of the form  $\mathfrak{S}F'$ , we shall be sure that the result of operating on any power ending product is a seminvariant and not a zero.

Now this is the case. Take any gradient  $F'$  whatever, and let  $a_r$  be the most advanced letter which occurs in it, so that

$$F' = A + a_r B + a_r^2 C + \dots,$$

where  $A, B, C, \dots$  are all free from  $a_r$ , and  $B, C, \dots$  do not all vanish. It follows that

$$\begin{aligned} \mathfrak{S}F' &= \left( \dots + a_{r+1} \frac{d}{da_r} + \dots \right) F' \\ &= a_{r+1} \{ B + 2a_r C + 3a_r^2 D + \dots \} + \text{terms free from } a_{r+1}. \end{aligned}$$

Thus,  $B, C, D, \dots$  not all vanishing,  $\mathfrak{S}F'$  contains necessarily a non-power ending term or terms; namely a term or terms ending in the first power  $a_{r+1}$ .

$$\text{Thus} \quad 1 - \frac{1}{i} \frac{\mathfrak{S} \Omega}{1} + \frac{1}{i^2} \frac{\mathfrak{S}^2 \Omega^2}{1 \cdot 2} - \dots,$$

which write

$$1 - \mathfrak{S}X,$$

generates seminvariants from all power enders.

Moreover, from the complete system of

$$(w; i, \infty) - (w-1; i, \infty)$$

power enders of degree  $i$  and weight  $w$  it generates a complete system of  $(w; i, \infty) - (w-1; i, \infty)$  seminvariants of that degree and weight. For, if possible, let the seminvariants

$$S_1 = (1 - \mathfrak{S}X)P_1, S_2 = (1 - \mathfrak{S}X)P_2, S_3 = (1 - \mathfrak{S}X)P_3, \dots,$$

generated from the complete system of power enders of degree  $i$  and weight  $w$ , be connected by a linear relation

$$\lambda_1 S_1 + \lambda_2 S_2 + \lambda_3 S_3 + \dots = 0.$$

This would necessitate that

$$\lambda_1(1 - \mathfrak{S}X)P_1 + \lambda_2(1 - \mathfrak{S}X)P_2 + \lambda_3(1 - \mathfrak{S}X)P_3 + \dots = 0,$$



i.e. that  $(1 - \mathfrak{S}X)(\lambda_1 P_1 + \lambda_2 P_2 + \lambda_3 P_3 + \dots) = 0,$

or  $\lambda_1 P_1 + \lambda_2 P_2 + \lambda_3 P_3 + \dots = \mathfrak{S} \cdot X (\lambda_1 P_1 + \lambda_2 P_2 + \lambda_3 P_3 + \dots),$

i.e., by the above, that

$$\lambda_1 P_1 + \lambda_2 P_2 + \lambda_3 P_3 + \dots$$

involve at least one non-power ending product. But it does not.

It is then completely established that there is a one to one correspondence between a complete system of power ends and a complete system of seminvariants, the latter complete system for any degree  $i$  being generated from the former by operation with

$$1 - \frac{1}{i} \frac{\mathfrak{S} \Omega}{1} + \frac{1}{i^2} \frac{\mathfrak{S}^2 \Omega^2}{1 \cdot 2} - \frac{1}{i^3} \frac{\mathfrak{S}^3 \Omega^3}{1 \cdot 2 \cdot 3} + \dots$$

The theorem of § 193, Ex. 15 would lead to the same conclusion as to one to one correspondence, and afford an alternative generator of all seminvariants from power ends.

### ADDITIONAL EXAMPLES (MISCELLANEOUS).

Ex. 16. A non-unitary symmetric function of the roots of an equation of order  $p$ , i.e. one whose expression in terms of  $s_1, s_2, \dots, s_p$  does not involve  $s_1$ , or, if  $p$  be not less than the weight  $l + m + n + \dots$ , one  $\Sigma(a_1^l a_2^m a_3^n \dots)$  in which none of  $l, m, n, \dots$  is unity, has its full expression in terms of the coefficients determinate when the non-unitary part of that expression, i.e. the part of it free from the unitary coefficient  $a_1$ , is known, just as the full expression of a seminvariant is determinate from its non-unitary portion. (*MacMahon.*)

Ex. 17. Prove that

$$\frac{1}{2} a^2 \partial_b + ab \partial_c + (ac + \frac{1}{2} b^2) \partial_d + (ad + bc) \partial_e + (ae + bd + \frac{1}{2} c^2) \partial_f + \dots$$

and

$$2 \cdot \frac{1}{2} a^2 \partial_b + 3ab \partial_c + 4(ac + \frac{1}{2} b^2) \partial_d + 5(ad + bc) \partial_e + 6(ae + bd + \frac{1}{2} c^2) \partial_f + \dots$$

generate seminvariants from seminvariants. (*MacMahon.*)

*Ans.* Form the alternants with the infinitely continued  $\Omega$ .

Ex. 18. If  $G$  be any gradient,

$$\left(1 - \frac{b}{a} \Omega + \frac{1}{1 \cdot 2} \frac{c}{a} \Omega^2 - \frac{1}{1 \cdot 2 \cdot 3} \frac{d}{a} \Omega^3 + \dots\right) G,$$

where  $\Omega$  is infinitely continued, is either a seminvariant, not necessarily integral, or zero.

Ex. 19. Among the seminvariants thus derived from gradients  $G$  all integral seminvariants of the type of  $G$  are included.

*Ans.* For the result of operating on  $S$  is to produce  $S$ .

Ex. 20. The coefficients of powers of  $x$  in the expansion of

$$\left( a - b \frac{ix}{1} + c \frac{i^2 x^2}{1.2} - d \frac{i^3 x^3}{1.2.3} + \dots \right) \left( a + b \frac{x}{1} + c \frac{x^2}{1.2} + d \frac{x^3}{1.2.3} + \dots \right)^i,$$

where  $i$  is any positive integer, are all seminvariants, except such as are zero.

Ex. 21. If  $\mathfrak{D}$  be the infinitely continued operator defined in § 192, and if  $S_1, S_2$  be two seminvariants of degrees  $i_1, i_2$  respectively, then  $i_2 S_2 \mathfrak{D} S_1 - i_1 S_1 \mathfrak{D} S_2$  is a seminvariant. (*D'Ocagne.*)

Ex. 22. Prove that  $\Omega$  annihilates the product

$$\left\{ a + b(x-y) + \frac{c}{1.2} (x-y)^2 + \dots \right\} \left\{ a + b(y-z) + \dots \right\} \\ \dots \left\{ a + b(u-v) + \dots \right\} \left\{ a + b(v-x) + \dots \right\},$$

where  $x, y, z, \dots u, v$  are arbitrary, and the series in brackets extend to infinity; and hence that all non-vanishing coefficients in the product expanded by powers and products of  $x, y, z, \dots u, v$  are seminvariants.

(*S. Roberts.*)

Ex. 23. If  $S$  be a seminvariant of  $(a, b, c, d, \dots) (x, y)^\infty$ , prove that  $(2w \mathfrak{D} - i\phi)S$ , where the notation is that of § 192, is another seminvariant of the same degree, and weight one higher. (*Cayley.*)

*Ans.* Cf. § 165, and the example 8 which follows it.

Ex. 24. If in any seminvariant of  $(a, b, 1, c, 1.2, d, 1.2.3, \dots) (x, y)^\infty$ ,  $a, b, c, d, \dots$  are replaced by  $b, c, d, e, \dots$  the result gives the terms free from  $a$  in a seminvariant of the same degree and higher weight. (*MacMahon.*)

Ex. 25. A seminvariant of  $(a, b, c, d, e, \dots) (x, y)^\infty$  is a seminvariant of the system

$$\begin{aligned} & (a, b, 1, c, 1.2, d, 1.2.3, \dots) (x, y)^\infty \\ & (b, c, 1, d, 1.2, e, 1.2.3, \dots) (x, y)^\infty \\ & (c, d, 1, e, 1.2, \dots) (x, y)^\infty \\ & \text{\&c., \&c.} \quad (\textit{MacMahon.}) \end{aligned}$$

Ex. 26. The seminvariant of weight and degree 3 of

$$(a_0, a_1, a_2, \dots a_p) (x, y)^p$$

is a multiple of  $a_0^3$  times the sum of the cubes of the roots of the equation

$$a_0 x^p + \frac{a_1}{1} x^{p-1} + \frac{a_2}{1.2} x^{p-2} + \dots + \frac{a_p}{p!} = 0.$$

Ex. 27. If  $G$  be any gradient of weight  $w$  degree  $i$  and extent  $p$ , and if  $\sigma$  be the sum of its numerical coefficients, then, whatever  $x$  be,

$$x^{-w} \left( \frac{d}{da_0} + x \frac{d}{da_1} + x^2 \frac{d}{da_2} + \dots + x^p \frac{d}{da_p} \right)^i G = i! \sigma.$$

Ex. 28. Hence any seminvariant of weight  $w$  and degree  $i$  is annihilated by the operator which is the coefficient of  $x^w$  in the expansion of

$$(\partial_a + x \partial_b + x^2 \partial_c + x^3 \partial_d + \dots)^i.$$

Ex. 29. Referring to § 174, Ex. 13 for the notation, show that, if  $\delta_r$ , denote  $r! (p-r)! \frac{d}{da_r}$ , and  $\delta'_r$  denote  $r! (p-r)! \frac{d}{da_r}$ ,  $\delta_p, \delta_{p-1}, \dots, \delta_0$  are the same functions of  $\delta'_p, \delta'_{p-1}, \dots, \delta'_0$  and  $m$  as  $a_0, a_1, \dots, a_p$  are of  $a_0, a_1, \dots, a_p$  and  $m$ .

Ex. 30. In the same notation  $\delta'_p, \delta'_{p-1}, \dots, \delta'_0$  are the same functions of  $\delta_p, \delta_{p-1}, \dots, \delta_0$  and  $-m$  as  $a_0, a_1, \dots, a_p$  are of  $a_0, a_1, \dots, a_p$  and  $m$ .

Ex. 31. If, upon the substitution of  $X+mY$ ,  $Y$  for  $x, y$ ,

$$(a_0, a_1, a_2, \dots, a_p) (x, y)^p$$

become

$$(a_0, a_1, a_2, \dots, a_p) (X, Y)^p,$$

then

$$(\delta_p, -\delta_{p-1}, \delta_{p-2}, \dots, (-1)^p \delta_0) (x, y)^p$$

becomes

$$(\delta'_p, -\delta'_{p-1}, \delta'_{p-2}, \dots, (-1)^p \delta'_0) (X, Y)^p. \quad (\text{Sylvester.})$$

Ex. 32. Any seminvariant of  $(a_0, a_1, a_2, \dots, a_p) (x, y)^p$  becomes, when in it  $\delta_p, -\delta_{p-1}, \delta_{p-2}, \dots, (-1)^p \delta_0$  are put for  $a_0, a_1, a_2, \dots, a_p$ , an operator which has the same effect on any function of  $a_0, a_1, a_2, \dots, a_p$  as the result of replacing in it  $\delta_p, \delta_{p-1}, \dots$  by  $\delta'_p, \delta'_{p-1}, \dots$  has on the equivalent function of  $a_0, a_1, a_2, \dots, a_p$ , and may be called a seminvariant operator. (*Sylvester.*)

Ex. 33. More generally, any operator obtained by writing down a seminvariant of the two quantities

$$(a_0, a_1, a_2, \dots, a_p) (x, y)^p, (\delta_p, -\delta_{p-1}, \delta_{p-2}, \dots, (-1)^p \delta_0) (x, y)^p,$$

the symbols  $\delta$  being written last in every term, is a seminvariant operator. (*Sylvester.*)

Ex. 34. Hence obtain the results (cf. § 186) that

$$i, \Omega, w - \frac{a_1}{a_0} \Omega, O - \frac{a_1}{a_0} (pi - 2w) - \frac{a_1^2}{a_0^2} \Omega$$

are four seminvariant operators.

Ex. 35. If  $a_1, a_2, \dots, a_i$  are  $i$  symbols, and

$$(a_1 - a_2)^{n_{12}} (a_1 - a_3)^{n_{13}} (a_2 - a_3)^{n_{23}} \dots$$

any product of  $w$  differences between pairs of them which is such that not more than  $p$  factors involve any one of the symbols, and if the product is expanded and multiplied by  $a_0^i$ , and in the result  $\frac{a_1}{a_0}$  is put for every one of the first powers  $a_1, a_2, \dots, a_i$ ,  $\frac{a_2}{a_0}$  for every one of  $a_1^2, a_2^2, \dots, a_i^2$ , and generally  $\frac{a_r}{a_0}$  for every one of  $a_1^r, a_2^r, \dots, a_p^r$ , the result is a seminvariant of  $(a_0, a_1, a_2, \dots, a_p)$   $(x, y)^p$  of degree  $i$  and weight  $w$ ; and all seminvariants are linear functions of seminvariants which can be thus expressed. (*Stroh.*)

*Ans.* Such a function is annihilated by  $\frac{d}{da_1} + \frac{d}{da_2} + \dots + \frac{d}{da_i}$ , and a seminvariant is annihilated by  $\Omega$ . Now both these operators are expressed by

$$\begin{aligned} \frac{\partial}{\partial a_1} + \frac{\partial}{\partial a_2} + \dots + \frac{\partial}{\partial a_i} + 2 \left\{ a_1 \frac{\partial}{\partial (a_1^2)} + \dots + a_i \frac{\partial}{\partial (a_i^2)} \right\} \\ + 3 \left\{ a_1^2 \frac{\partial}{\partial (a_1^3)} + \dots \right\} + \dots + p \left\{ a_1^{p-1} \frac{\partial}{\partial (a_1^p)} + \dots \right\}. \end{aligned}$$

Or again, the functions lead hyperdeterminants (§ 60).

Ex. 36. If  $\lambda_1, \lambda_2, \dots, \lambda_i$  be  $i$  quantities whose sum is zero, then, after expansion and substitution as in the last example,

$$a_0^i (\lambda_1 a_1 + \lambda_2 a_2 + \dots + \lambda_i a_i)^w$$

is a seminvariant of degree  $i$  and weight  $w$ , provided that  $w$  do not exceed  $p$ . (*Stroh.*)

Ex. 37. If  $p$  be infinite, or not less than  $w$ , and if  $e_1 (= 0), e_2, e_3, \dots, e_i$  be the elementary symmetric functions

$$\Sigma(\lambda) (= 0), \Sigma(\lambda_1 \lambda_2), \Sigma(\lambda_1 \lambda_2 \lambda_3), \dots, \lambda_1 \lambda_2 \dots \lambda_i,$$

then, when the function of the last example is expanded and expressed in terms of powers and products of  $e_2, e_3, \dots, e_p$ , and substitution for the  $a$ 's and their powers made as before, the coefficients of the various products of  $e$ 's are a complete system of  $(w; \infty, i) - (w-1; \infty, i)$  linearly independent seminvariants of weight  $w$  and degree  $i$ . (*Stroh.*)

Ex. 38. If the numbers of powers and products of  $e_2, e_3, \dots, e_i$  in the sum of Ex. 37 be diminished as much as possible by means of the relations in  $e_2, e_3, \dots, e_i$  any one of which expresses that in some way  $\lambda_1 + \lambda_2 + \dots + \lambda_i$  is a sum of two sums

$$\lambda_1 + \dots + \lambda_\phi, \lambda_{\phi+1} + \dots + \lambda_i$$

each of which vanishes, the coefficients of powers and products of the  $e$ 's which remain are non-perpetuant seminvariants, and the number of

perpetuants of degree  $i$  and weight  $w$  is the number of powers and products which have disappeared. (*Stroh.*)

*Ans.* Seminvariants which do not disappear are reducible in terms of seminvariants of lower degree, and others which are not syzygetic with these are not so reducible.

Ex. 39. Perpetuants of degree  $i$  and weight  $w$  are just as numerous as products of  $e_2, e_3, \dots e_i$  and powers of them which when multiplied

by 
$$\Pi \lambda \cdot \Pi (\lambda_1 + \lambda_2) \cdot \Pi (\lambda_1 + \lambda_2 + \lambda_3) \dots \Pi (\lambda_1 + \lambda_2 + \dots + \lambda_\nu)$$

are raised to weight  $w$  in  $e$ -suffixes, i.e. to dimensions  $w$  in the  $\lambda$ 's. Here

$$\Pi \lambda = e_i, \Pi (\lambda_1 + \lambda_2) = (\lambda_1 + \lambda_2) (\lambda_1 + \lambda_3) \dots (\lambda_2 + \lambda_3) \dots (\lambda_{i-1} + \lambda_i),$$

&c., and  $\nu$  is  $\frac{1}{2}i$  or  $\frac{1}{2}(i-1)$  according as  $i$  is even or odd. (If  $i$  is even, two conjugate sums  $\lambda_1 + \lambda_2 + \dots + \lambda_\nu, \lambda_{\nu+1} + \lambda_{\nu+2} + \dots + \lambda_i$  are not both written in the last product.) (*Stroh.*)

Ex. 40. The product  $\Pi \lambda \cdot \Pi (\lambda_1 + \lambda_2) \dots \Pi (\lambda_1 + \lambda_2 + \dots + \lambda_\nu)$  is of weight  $2^{i-1} - 1$ , whether  $i$  be even or odd. Consequently the weight of the lowest perpetuant of degree  $i$  is  $2^{i-1} - 1$ . (*Stroh.*)

*Ans.* For instance,  $i$  even gives the weight

$$i + \frac{i(i-1)}{1.2} + \dots + \frac{1}{2} \frac{i(i-1) \dots \left(\frac{i}{2} + 1\right)}{1.2 \dots \frac{i}{2}} = \frac{1}{2} 2^i - 1.$$

Ex. 41. The number of perpetuants of a higher weight  $w$  than this, and of degree  $i$ , is the number of solutions in positive integral and zero values of  $\mu_2, \mu_3, \mu_4, \dots$  of

$$2^{i-1} - 1 + 2\mu_2 + 3\mu_3 + 4\mu_4 + \dots + i\mu_i = w.$$

Ex. 42. Deduce the generating function for perpetuants (§ 189), viz.

$$\frac{x^{2^{i-1}-1}}{(1-x^2)(1-x^3) \dots (1-x^i)}.$$

## CHAPTER XII.

### CANONICAL FORMS, ETC.

195.] WHEN a binary quantic  $(a_0, a_1, a_2, \dots a_p) (x, y)^p$  is transformed by the linear substitution

$$x = lX + mY, y = l'X + m'Y,$$

four constants  $l, m, l', m'$  are introduced whose values may be assigned at will. These may be so chosen that the form of the transformed quantic is simplified by the absence of certain of its coefficients, or by relations among certain coefficients. The quantic is thus reduced to a simpler form without any loss of generality.

For instance, we know perfectly well that by giving  $l, m, l', m'$  the values  $1, -\frac{a_1}{a_0}, 0, 1$  the quantic is transformed into one wanting its second term. The quantic without a second term is then not a special one, but is in effect just as general as one with its second term present. Any binary quantic can be so expressed by means of a linear transformation.

Moreover, it is to be noticed that the deprivation of a quantic of its second term is not something which can be done by linear transformation in one way only, but that there is a wide class of linear substitutions any one of which will effect the purpose. In fact, if by the general linear substitution  $(a_0, a_1, a_2, \dots a_p) (x, y)^p$  be transformed into  $(A_0, A_1, A_2, \dots A_p) (X, Y)^p$ , we see at once that

$$A_1 = a_0 l^{p-1} m + a_1 (l^{p-1} m' + \overline{p-1} l^{p-2} m l') + \dots + a_p l^{p-1} m',$$

in which we may give to  $l, m, l'$  any values we please, and obtain, by solution of an equation of the first degree, a value of  $m'$  which, going with those values of  $l, m, l'$ , will make

$A_1$  vanish. The usual way of depriving a quantic of its second term is then only the simplest of many ways.

We see, in fact, that by proper choice of the four quantities  $l, m, l', m'$  we may in general impose four conditions on the coefficients in a binary quantic, and still have a form to which the quantic can be reduced by a linear substitution without losing its generality. These may not be any four conditions we choose, for the equations in  $l, m, l', m'$  which express four conditions may not prove to be consistent with one another. In particular, for instance, we can never make four separate coefficients in the quantic vanish. For to express this we should have to make  $l, m, l', m'$  satisfy four homogeneous equations, i.e. to choose the ratios  $\frac{l}{m}, \frac{l'}{m}, \frac{m'}{m}$ , three quantities, so as to satisfy four independent equations, which cannot be done.

196.] **Definition of canonical forms.** Now the binary  $p$ -ic contains  $p+1$  coefficients. Taking 4 from this number, we see that no binary  $p$ -ic with less than  $p-3$  perfectly arbitrary coefficients can be equivalent to a perfectly general binary  $p$ -ic subjected to linear transformation, but that there is a certain presumption in favour of one which has  $p-3$  perfectly arbitrary coefficients, or whose coefficients involve  $p-3$  perfectly arbitrary quantities, being equivalent to the general binary  $p$ -ic, which presumption must, however, in every case be tested before it can be stated as a certainty.

A form of binary  $p$ -ic whose coefficients involve  $p-3$  perfectly arbitrary quantities, and which is proved to be a form to which the general binary  $p$ -ic can be reduced by a linear substitution, is called a *Canonical Form* of the binary  $p$ -ic. There may be different forms for the same value of  $p$  which have equal claims to the name canonical, but in practice, for the cubic, quartic, &c., respectively, one canonical form is chosen because of symmetry of shape and convenience of treatment, and often spoken of as *the* canonical form.

The case  $p=2$  of the quadratic stands by itself in that  $p-3$  is negative. Of course the three coefficients of a binary quadratic cannot be subjected to more than three conditions. To each of the simple forms  $X^2+Y^2, XY$  a quadratic can be

reduced in an infinite number of ways, since one of  $l, m, l', m'$  is arbitrary. Indeed, the general binary quadratic

$$ax^2 + 2bxy + cy^2$$

can be given the form of any quadratic  $a'X^2 + 2b'XY + c'Y^2$  whatever whose discriminant  $a'c' - b'^2$  does not vanish. The like fact is true as to quadratics in any number of variables.

197.] Canonical forms have here been defined for binary quantics only. For quantics in more variables than two the definition is similar. A form of  $q$ -ary  $p$ -ic which is a simplest form to which linear transformation can reduce the general  $q$ -ary  $p$ -ic is a canonical form of the  $q$ -ary  $p$ -ic, one form being regarded as more simple than another when of its coefficients a smaller number are arbitrary, or, as is the same thing, when its coefficients are known functions of a smaller number of arbitrary quantities.

The number of coefficients in the  $q$ -ary  $p$ -ic being easily seen to be

$$\frac{(p+1)(p+2)\dots(p+q-1)}{(q-1)!},$$

and the number of constants in the general scheme of linear substitution being  $q^2$ , the number of perfectly arbitrary coefficients left in a canonical form will be

$$\frac{(p+1)(p+2)\dots(p+q-1)}{(q-1)!} - q^2,$$

when the degree  $p$  is sufficiently great for this to be positive.

198.] The knowledge of invariants and covariants both aids, and is aided by the determination of canonical forms of quantics. On the one hand, as we shall illustrate by examples, invariants and covariants supply information as to forms which are canonical and the reduction of general quantics to those forms, and on the other, since invariants and covariants of a quantic have relations to one another, expressed by homogeneous isobaric syzygies, which hold however the quantic be linearly transformed, it suffices, in order to discover those relations, to consider the quantic and the invariants and covariants in simpler forms which they can assume without loss of generality.



Geometrical interpretation of invariants and covariants is also greatly assisted by the simplification afforded by canonical forms.

199.] Canonical form of binary cubic. The binary cubic

$$ax^3 + 3bx^2y + 3cxy^2 + dy^3$$

can be expressed in the canonical form

$$X^3 + Y^3.$$

In other words, constants  $\lambda, \mu, \lambda', \mu'$  can be found such that

$$ax^3 + 3bx^2y + 3cxy^2 + dy^3 \equiv (\lambda x + \mu y)^3 + (\lambda' x + \mu' y)^3$$

is an identity.

A presumption in favour of this is afforded by the fact that the identification of coefficients of  $x^3, x^2y, xy^2, y^3$  on the left and right gives four equations for the determination of  $\lambda, \mu, \lambda', \mu'$ ; but we have to be sure that the four equations are consistent and independent and can actually be solved. This will first be proved without any reference to the invariant theory.

With a change of notation, we have to see that  $p, q, a, \beta$  can be found so as to make

$$ax^3 + 3bx^2y + 3cxy^2 + dy^3 \equiv p(x + ay)^3 + q(x + \beta y)^3,$$

i. e. so as to make simultaneously

$$p + q = a,$$

$$pa + q\beta = b,$$

$$pa^2 + q\beta^2 = c,$$

$$pa^3 + q\beta^3 = d.$$

The first three of these are consistent for the determination of  $p, q$  if

$$\begin{vmatrix} 1, & 1, & a \\ a, & \beta, & b \\ a^2, & \beta^2, & c \end{vmatrix} = 0,$$

and the values of  $pa, q\beta$  which satisfy the second and third also satisfy the fourth if

$$\begin{vmatrix} 1, & 1, & b \\ a, & \beta, & c \\ a^2, & \beta^2, & d \end{vmatrix} = 0.$$

We have thus two equations for the determination of  $a, \beta$ . We may write them

$$Pa + Qb + Rc = 0,$$

and

$$Pb + Qc + Rd = 0,$$

where also  $P + Qa + Ra^2 = 0$ , and  $P + Q\beta + R\beta^2 = 0$ . These are made consistent by taking for  $a$  and  $\beta$  the two roots of the quadratic

$$\begin{vmatrix} a, b, c \\ b, c, d \\ 1, a, a^2 \end{vmatrix} = 0.$$

Having thus found  $a$  and  $\beta$ , any two of the first set of four equations suffice to determine  $p$  and  $q$ . Thus the possibility of reducing the cubic to the canonical form  $X^3 + Y^3$  is proved, and the means of doing it, by solution of quadratic and linear equations, afforded.

The student should notice that there is failure to effect what is desired when  $a, b, c, d$  have such specially connected values that the quadratic for  $a$  and  $\beta$  has equal roots, i.e. when

$$(ad - bc)^2 - 4(ac - b^2)(bd - c^2) = 0,$$

that is to say, when the discriminant of the cubic vanishes, so that the cubic has a square factor. The canonical form for cubics with a square factor is not  $X^3 + Y^3$  but  $X^2Y$ .

This leads to the general remark that a canonical form of the general quantic of any type is one to which a quantic of that type can be reduced when general, but not necessarily one to which every special quantic of that type can be reduced.

Ex. 1. Verify that

$$(pq)^2 (a - \beta)^6 = (ad - bc)^2 - 4(ac - b^2)(bd - c^2).$$

Ex. 2. A binary cubic with general coefficients can be linearly transformed into any other.

*Ans.* Through  $X^3 + Y^3$  as an intermediary.

200.] The reduction of the cubic to the form

$$p(x + ay)^3 + q(x + \beta y)^3$$

and thence to its canonical form  $X^3 + Y^3$  is most easily effected by means of its one quadratic covariant, the Hessian

$$(ac - b^2)x^2 + (ad - bc)xy + (bd - c^2)y^2.$$

Regard the cubic

$$ax^3 + 3bx^2y + 3cxy^2 + dy^3$$

as the transformed form obtained by the substitution

$$x' = x + \alpha y, \quad y' = y + \beta y,$$

whose modulus is  $\beta - \alpha$ , from the form  $px'^3 + qy'^3$ .

The Hessian of the transformed is the Hessian of the untransformed multiplied by the square of the modulus. Thus

$$\begin{aligned} (ac - b^2)x^2 + (ad - bc)xy + (bd - c^2)y^2 &= (\beta - \alpha)^2 pq x' y' \\ &= pq (\alpha - \beta)^2 (x + \alpha y)(x + \beta y). \end{aligned}$$

Consequently if the Hessian

$$(ac - b^2)x^2 + (ad - bc)xy + (bd - c^2)y^2$$

be broken up into factors

$$(ac - b^2)(x + \alpha y)(x + \beta y),$$

the cubic must have the form

$$p(x + \alpha y)^3 + q(x + \beta y)^3,$$

in which  $p$  and  $q$  may be found by the equations

$$p + q = a, \quad pa + q\beta = b.$$

Thus  $p^{\frac{1}{3}}(x + \alpha y)$  and  $q^{\frac{1}{3}}(x + \beta y)$ , the  $X$  and  $Y$  of the canonical form, are found.

The determination of the canonical form of the binary cubic effects the solution of a cubic equation. (Cf. § 11, Exx. 14, 15.) For it reduces the cubic equation to the form

$$X^3 + Y^3 = 0,$$

i.e. to the three linear equations

$$X + Y = 0, \quad X + \omega Y = 0, \quad X + \omega^2 Y = 0.$$

The student is advised to illustrate this by an example, e.g. to solve  $x^3 - 3x^2 \tan \alpha - 3x + \tan \alpha = 0$ .

201.] **Concomitants of cubic in canonical form.** We have seen in § 169 and elsewhere that the cubic

$$u \equiv ax^3 + 3bx^2y + 3cxy^2 + dy^3$$

has, besides its Hessian

$$H \equiv (ac - b^2)x^2 + (ad - bc)xy + (bd - c^2)y^2,$$

a cubicovariant

$$G \equiv (a^2d - 3abc + 2b^3)x^3 + 3(abd - 2ac^2 + b^2c)x^2y \\ + 3(2b^2d - acd - bc^2)xy^2 + (3bcd - ad^2 - 2c^3)y^3,$$

and one invariant, its discriminant

$$\Delta \equiv (ad - bc)^2 - 4(ac - b^2)(bd - c^2).$$

The same functions of the coefficients and variables in the canonical forms are

$$X^3 + Y^3,$$

$$XY,$$

$$X^3 - Y^3,$$

1.

Now let  $M'$  be the modulus of the substitution which expresses  $X$  and  $Y$  in terms of  $x$  and  $y$ , so that in the two notations of § 199

$$M' = \lambda\mu' - \lambda'\mu = (pq)^3(\beta - \alpha).$$

We remember from chapters ii, iii, that the index of the power of the modulus, which has to multiply an invariant or covariant of a binary quantic to produce the equivalent of the same invariant or covariant of the transformed quantic, is equal to the weight of the invariant or of the leading coefficient in the covariant. Thus the information given by invariant algebra as to a binary cubic and its canonical form is presented in the four identities

$$u = X^3 + Y^3,$$

$$H = M'^2 \cdot XY,$$

$$G = M'^3(X^3 - Y^3),$$

$$\Delta = M'^6.$$

Of these the last tells us at once that the modulus of the substitution which expresses  $X$  and  $Y$  in terms of  $x$  and  $y$ , i.e. the reciprocal (§ 23) of the modulus of that which expresses  $x$  and  $y$  in terms of  $X$  and  $Y$ , is equal to the sixth root of the discriminant.

We have also in a clear form before us the fact that

$u, H, G, \Delta$ , though irreducible in that none of them can be expressed rationally and integrally in terms of the rest, are not independent, but are connected by the syzygy (cf. § 169)

$$u^2\Delta = G^2 + 4H^3,$$

which is obtained by eliminating  $X, Y$  and  $M'$ . Moreover, no other syzygy connects them, for there is no other way of eliminating those three quantities.

It is of interest to notice that we have also readily given by these identities the values of the  $p, q, a, \beta$  of § 199. We have

$$u = X^3 + Y^3 = p(x + \alpha y)^3 + q(x + \beta y)^3,$$

$$\Delta^{-\frac{1}{2}}G = X^3 - Y^3 = p(x + \alpha y)^3 - q(x + \beta y)^3.$$

Thus, taking the full expressions for  $u$  and  $G$ , and attending only to the equalities of the coefficients of  $x^3$  and  $x^2y$ ,

$$p + q = a,$$

$$pa + q\beta = b,$$

$$p - q = \Delta^{-\frac{1}{2}}(a^2d - 3abc + 2b^3),$$

$$pa - q\beta = \Delta^{-\frac{1}{2}}(abd - 2ac^2 + b^2c);$$

whence

$$2p = a + \Delta^{-\frac{1}{2}}(a^2d - 3abc + 2b^3),$$

$$2q = a - \Delta^{-\frac{1}{2}}(a^2d - 3abc + 2b^3),$$

$$2pa = b + \Delta^{-\frac{1}{2}}(abd - 2ac^2 + b^2c),$$

$$2q\beta = b - \Delta^{-\frac{1}{2}}(abd - 2ac^2 + b^2c).$$

We have also  $X^3$  and  $Y^3$  themselves; viz.

$$\frac{1}{2}(u + \Delta^{-\frac{1}{2}}G) \text{ and } \frac{1}{2}(u - \Delta^{-\frac{1}{2}}G).$$

The solutions of the cubic equation

$$(a, b, c, d)(x, y)^3 = 0$$

in  $x : y$  are then given by the three linear equations

$$(u + \Delta^{-\frac{1}{2}}G)^{\frac{1}{3}} + (u - \Delta^{-\frac{1}{2}}G)^{\frac{1}{3}} = 0,$$

$$(u + \Delta^{-\frac{1}{2}}G)^{\frac{1}{3}} + \omega(u - \Delta^{-\frac{1}{2}}G)^{\frac{1}{3}} = 0,$$

$$(u + \Delta^{-\frac{1}{2}}G)^{\frac{1}{3}} + \omega^2(u - \Delta^{-\frac{1}{2}}G)^{\frac{1}{3}} = 0.$$

202.] **Geometry of concomitants of cubic.** Geometrically, taking  $(a, b, c, d)(x, y)^3 = 0$  to represent three straight lines

through a point, the reduction of the cubic to its canonical form is the reference to the lines which form the Hessian. As examples of geometrical information yielded by the canonical form the following are left to the student.

Ex. 3. The cubicovariant of a pencil of three lines  $L, M, N$  represents the pencil  $L', M', N'$  which consists of the harmonic conjugate of  $L$  with regard to  $M$  and  $N$ , that of  $M$  with regard to  $N$  and  $L$ , and that of  $N$  with regard to  $L$  and  $M$ .

*Ans.* It suffices to prove that  $X - Y$  and  $X + Y$  are harmonic with regard to  $X + \omega Y$  and  $X + \omega^2 Y$ .

Ex. 4.  $L, L'; M, M'; N, N'$  are pairs of an involution, of which  $H$  the Hessian represents the double lines.

By means of the expressions for  $u, H, G, \Delta$  in terms of  $X, Y, M'$  in § 201 we may readily prove the following theorem due to Cayley.

Ex. 5. The Hessian, cubicovariant, and discriminant of  $ku + k'G$  are respectively

$$(k^2 - k'^2 \Delta)H, (k^2 - k'^2 \Delta)(kG + k' \Delta u), (k^2 - k'^2 \Delta)^2 \Delta.$$

Ex. 6. If  $L', M', N'$  are the harmonic conjugates of  $L$  with regard to  $M$  and  $N$ , of  $M$  with regard to  $N$  and  $L$ , and of  $N$  with regard to  $L$  and  $M$ , then  $L, M, N$  are respectively the harmonic conjugates of  $L'$  with regard to  $M'$  and  $N'$ , of  $M'$  with regard to  $N'$  and  $L'$ , and of  $N'$  with regard to  $L'$  and  $M'$ .

203.] **Canonical reduction of binary  $(2n-1)$ -ic.** The proposition of § 199 is a case of a general one due, like most of the rest of the theory of canonical forms, to Sylvester. This is that a general binary quantic of odd order  $2n-1$  is a sum of  $n$   $(2n-1)$ th powers of linear forms.

As indicating the likelihood of this, we notice that the sum

$$(\lambda_1 x + \mu_1 y)^{2n-1} + (\lambda_2 x + \mu_2 y)^{2n-1} + \dots + (\lambda_n x + \mu_n y)^{2n-1},$$

or its equivalent

$$p_1 (x + a_1 y)^{2n-1} + p_2 (x + a_2 y)^{2n-1} + \dots + p_n (x + a_n y)^{2n-1},$$

is a binary  $(2n-1)$ -ic with no obvious connexion among its coefficients, which coefficients are functions of  $2n$  constants that may be chosen at will, this number  $2n$  being exactly that of coefficients in the general binary  $(2n-1)$ -ic

$$(a_0, a_1, a_2, \dots, a_{2n-1})(x, y)^{2n-1}.$$

We have to see, however, that values of the  $2n$   $a$ 's and  $p$ 's, to adopt the second notation, actually exist, which make the sum and the quantic identical. We shall prove that the  $a$ 's are the roots of an equation of degree  $n$ , and so do exist, though general expressions cannot algebraically be found for them when  $n$  exceeds 4, and that, when the  $a$ 's are known, the  $p$ 's are determinate by solution of equations of the first degree.

For the identity

$$(a_0, a_1, a_2, \dots, a_{2n-1}) (x, y)^{2n-1} \equiv p_1(x + a_1y)^{2n-1} + p_2(x + a_2y)^{2n-1} + \dots + p_n(x + a_ny)^{2n-1}$$

to hold, we must have simultaneously

$$\begin{aligned} p_1 &+ p_2 &+ \dots + p_n &= a_0, \\ p_1 a_1 &+ p_2 a_2 &+ \dots + p_n a_n &= a_1, \\ p_1 a_1^2 &+ p_2 a_2^2 &+ \dots + p_n a_n^2 &= a_2, \\ \dots &\dots &\dots &\dots \\ p_1 a_1^{2n-1} &+ p_2 a_2^{2n-1} &+ \dots + p_n a_n^{2n-1} &= a_{2n-1}. \end{aligned}$$

To prove that values of the letters on the left exist which satisfy these  $2n$  equations it will suffice to show that there is a recurring series, with scale of relation of the  $n$ th degree, of which  $a_0, a_1, a_2, \dots, a_{2n-1}$  are the first  $2n$  terms. Now  $n$  quantities  $q_1, q_2, \dots, q_n$  can at once be found to satisfy the  $n$  equations of the first degree

$$\begin{aligned} a_n &+ q_1 a_{n-1} + q_2 a_{n-2} + \dots + q_n a_0 &= 0, \\ a_{n+1} &+ q_1 a_n + q_2 a_{n-1} + \dots + q_n a_1 &= 0, \\ a_{n+2} &+ q_1 a_{n+1} + q_2 a_n + \dots + q_n a_2 &= 0, \\ \dots &\dots &\dots &\dots \\ a_{2n-1} &+ q_1 a_{2n-2} + q_2 a_{2n-3} + \dots + q_n a_{n-1} &= 0; \end{aligned}$$

and thus a scale of relation

$$a^n + q_1 a^{n-1} + q_2 a^{n-2} + \dots + q_n = 0$$

of a recurring series to which  $a_0, a_1, a_2, \dots, a_{2n-1}$  in order belong is determined.

By the ordinary theory of recurring series the  $n$  roots of this scale of relation are the  $a_1, a_2, \dots, a_n$  required; and, these being known, the solution of any  $n$  of the  $2n$  equations whose

right-hand sides are  $a_0, a_1, a_2, \dots, a_{2n-1}$  give uniquely the values of  $p_1, p_2, p_3, \dots, p_n$ .

The equation whose roots are  $a_1, a_2, \dots, a_n$  has the form

$$\begin{vmatrix} a_n & , & a_{n-1} & , & a_{n-2} & , & \dots & a_0 \\ a_{n+1} & , & a_n & , & a_{n-1} & , & \dots & a_1 \\ a_{n+2} & , & a_{n+1} & , & a_n & , & \dots & a_2 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ a_{2n-1} & , & a_{2n-2} & , & a_{2n-3} & , & \dots & a_{n-1} \\ a^n & , & a^{n-1} & , & a^{n-2} & , & \dots & 1 \end{vmatrix} = 0,$$

as is at once seen by elimination of  $q_1, q_2, \dots, q_n$ .

For the quintic  $(a, b, c, d, e, f) (x, y)^5$ , and the septic  $(a, b, c, d, e, f, g, h) (x, y)^7$ ,  $n$  has the values 3 and 4 respectively. Thus the reduction of the quintic to a sum of three fifth powers, and that of the septic to a sum of four seventh powers, can actually be effected algebraically. For quantities of higher odd orders the actual reduction would depend on the solution of equations in  $a$  of degrees above the fourth. For such higher cases the quintic is proved to have an equivalent expression as a sum of powers, but the algebraic reduction to the form is not effected.

204.] **Case of canonizing equation having equal roots.** There is failure to effect the required reduction when the coefficients in the  $(2n-1)$ -ic are so specially connected that the equation of the  $n$ th degree in  $a$  has equal roots.

The condition for such equality of roots is the vanishing of the discriminant of the  $n$ -ic in  $a$ . This is of degree  $2(n-1)$  in the coefficients of the  $n$ -ic, which themselves are of degree  $n$  in  $a_0, a_1, a_2, \dots, a_{2n-1}$ . The condition is then the vanishing of a function of degree  $2n(n-1)$  in the coefficients of the  $(2n-1)$ -ic. This function is an invariant, being the discriminant of what will presently be exhibited as a covariant. For the case of the quintic  $n = 3$ , and the invariant is of degree 12.

Let us discuss the failure for the case of the quintic. The equations which express that  $a, b, c, d, e, f$  form a recurring series whose scale of relation is  $(1 - ax)(1 - \beta x)^2$  are not those



of § 203, with  $n = 3$ , but

$$\begin{aligned} p + q &= a, \\ pa + (q + r)\beta &= b, \\ pa^2 + (q + 2r)\beta^2 &= c, \\ pa^3 + (q + 3r)\beta^3 &= d, \\ pa^4 + (q + 4r)\beta^4 &= e, \\ pa^5 + (q + 5r)\beta^5 &= f, \end{aligned}$$

of which any three determine  $p, q, r$ .

Now these give the quintic the form

$$p(x + ay)^5 + q(x + \beta y)^5 + 5r\beta y(x + \beta y)^4,$$

i. e.

$$p(x + ay)^5 + q(x + \beta y)^5 + \frac{5r\beta}{\alpha - \beta} \{(x + ay) - (x + \beta y)\} (x + \beta y)^4,$$

whose form is

$$p(x + ay)^5 + 5r'(x + ay)(x + \beta y)^4 + q'(x + \beta y)^5.$$

Thus the canonical form of a quintic which is special in that its invariant of the twelfth degree above vanishes is most simply written

$$X^5 + 5\lambda XY^4 + Y^5,$$

in which three consecutive coefficients are wanting.

When  $a, \beta, \gamma$  are all equal, it is easy to see that the degenerate form is

$$(x + ay)^3 \{p(x + ay)^2 + 5q(x + ay)y + 10ry^2\},$$

so that the quintic has a perfect cube for a factor.

205.] **Canonical forms of quintic, septic, &c.** In the identity

$$(a, b, c, d, e, f)(x, y)^5 \equiv p(x + ay)^5 + q(x + \beta y)^5 + r(x + \gamma y)^5,$$

we may write  $X$  for  $p^{\frac{1}{5}}(x + ay)$  and  $Y$  for  $q^{\frac{1}{5}}(x + \beta y)$ , and consequently  $\lambda X + \mu Y$  for  $r^{\frac{1}{5}}(x + \gamma y)$ , where  $\lambda$  and  $\mu$  are constants. We thus have as a canonical form of the general binary quintic

$$X^5 + Y^5 + (\lambda X + \mu Y)^5,$$

which involves two free constants only. More symmetrically we may write it

$$X^5 + Y^5 + Z^5,$$

where  $X, Y, Z$  are connected by a linear relation without constant term; or again, we may write it

$$\lambda'x'^5 + \mu'y'^5 + z'^5,$$

where

$$x' + y' + z' = 0.$$

In like manner a canonical form of the general binary septic is

$$X^7 + Y^7 + (\lambda X + \mu Y)^7 + (\lambda' X + \mu' Y)^7;$$

and similarly for binary quantics of higher odd orders. —

206.] **Canonizants.** In § 200 it was seen that the  $x + ay, x + \beta y$  of the cubic have for their product multiplied by a function of the coefficients a certain covariant, the Hessian, which may be written in either of the forms

$$\begin{vmatrix} ax + by, & bx + cy \\ bx + cy, & cx + dy \end{vmatrix},$$

$$\begin{vmatrix} a, & b, & c \\ b, & c, & d \\ y^2, & -xy, & x^2 \end{vmatrix}.$$

There are corresponding facts for the quintic, septic, ...  $(2n-1)$ -ic.

The covariant whose factors are

$$x + a_1y, x + a_2y, \dots x + a_ny$$

is not, after the cubic, the Hessian, but its form is analogous to either of the forms of the Hessian of the cubic here written down.

Regard the equation whose roots are  $a_1, a_2, \dots a_n$  which has been exhibited in § 203; and remember that, if  $a_1, a_2, \dots a_n$  are the roots of

$$x^n + q_1x^{n-1} + q_2x^{n-2} + \dots + q_n = 0,$$

then  $x + a_1y, x + a_2y, \dots x + a_ny$  are the factors of

$$x^n - q_1x^{n-1}y + q_2x^{n-2}y^2 - \dots + (-1)^n q_ny^n.$$

We at once gather, altering the arrangement of rows in the canonizing determinant, that

$$\begin{vmatrix} x^n & , & -x^{n-1}y, & x^{n-2}y^2, & -x^{n-3}y^3, & \dots & (-1)^n y^n \\ a_n & , & a_{n-1} & , & a_{n-2} & , & a_{n-3} & , & \dots & a_0 \\ a_{n+1}, & a_n & , & a_{n-1} & , & a_{n-2} & , & \dots & a_1 \\ a_{n+2}, & a_{n+1} & , & a_n & , & a_{n-1} & , & \dots & a_2 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ a_{2n-1}, & a_{2n-2}, & a_{2n-3} & , & a_{2n-4} & , & \dots & a_{n-1} \end{vmatrix}$$

$$\equiv (x + a_1 y) (x + a_2 y) \dots (x + a_n y) \begin{vmatrix} a_{n-1} & , & a_{n-2} & , & a_{n-3} & , & \dots & a_0 \\ a_n & , & a_{n-1} & , & a_{n-2} & , & \dots & a_1 \\ a_{n+1} & , & a_n & , & a_{n-1} & , & \dots & a_2 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ a_{2n-2}, & a_{2n-3}, & a_{2n-4}, & \dots & a_{n-1} \end{vmatrix}$$

For in p  
y = a  
by adding  
sum w  
eg is 0 +  
is a factor.  
: Det of is y  
T. (x + a<sub>1</sub>y) - (x  
- to find T part  
+ the T  
= |a<sub>n-1</sub>  
s  
m-2

Accordingly the determination of the

$$x + a_1 y, x + a_2 y, \dots, x + a_n y$$

of the canonical expression of the  $(2n - 1)$ -ic is effected by the breaking up of the  $n$ -ic which is the determinant on the left into its  $n$  factors.

To reduce the determinant to its other form, we best proceed by multiplying it according to the ordinary rule by another determinant of the same number of rows and columns, viz.

$$\begin{vmatrix} y, & x, & 0, & 0, & \dots & 0, & 0 \\ 0, & y, & x, & 0, & \dots & 0, & 0 \\ 0, & 0, & y, & x, & \dots & 0, & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0, & 0, & 0, & 0, & \dots & y, & x \\ 0, & 0, & 0, & 0, & \dots & 0, & 1 \end{vmatrix},$$

whose value is  $y^n$ . Combining rows with rows the product is

$$\begin{vmatrix} 0 & , & 0 & , & 0 & , & \dots & 0 & , & (-1)^n y^n \\ a_n y + a_{n-1} x & , & a_{n-1} y + a_{n-2} x & , & a_{n-2} y + a_{n-3} x & , & \dots & a_1 y + a_0 x & , & a_0 \\ a_{n+1} y + a_n x & , & a_n y + a_{n-1} x & , & a_{n-1} y + a_{n-2} x & , & \dots & a_2 y + a_1 x & , & a_1 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ a_{2n-1} y + a_{2n-2} x, & a_{2n-2} y + a_{2n-3} x, & a_{2n-3} y + a_{2n-4} x, & \dots & a_n y + a_{n-1} x, & & & & & a_{n-1} \end{vmatrix},$$

i.e. rearranging columns, is

$$(-1)^{\frac{1}{2}n(n-1)}y^n \begin{vmatrix} a_0x + a_1y, & a_1x + a_2y, & \dots, & a_{n-1}x + a_ny \\ a_1x + a_2y, & a_2x + a_3y, & \dots, & a_nx + a_{n+1}y \\ \cdot & \cdot & \cdot & \cdot \\ a_{n-1}x + a_ny, & a_nx + a_{n+1}y, & \dots, & a_{2n-2}x + a_{2n-1}y \end{vmatrix}.$$

The omission of the factor  $y^n$  from each side now establishes the identity, but for sign at most, of this last determinant with the first.

The determinant is a covariant, viz. the catalecticant of the  $(2n-2)$ th emanant (§ 56, cf. also § 17, Ex. 20). In the last written form of determinant the convention of § 71 as to sign and numerical multiple will be seen to have been adopted. The covariant, from the property here developed in connexion with canonical forms, is called the *canonizant* of the  $(2n-1)$ -ic.

207.] To realize the conclusion by particularization let us restate it for the quintic only. To reduce the quintic

$$(a, b, c, d, e, f)(x, y)^5$$

to its canonical form  $X^5 + Y^5 + Z^5$ , form the canonizant

$$\begin{vmatrix} ax+by, & bx+cy, & cx+dy \\ bx+cy, & cx+dy, & dx+ey \\ cx+dy, & dx+ey, & ex+fy \end{vmatrix},$$

and break it up into three linear factors  $\lambda_1x + \mu_1y$ ,  $\lambda_2x + \mu_2y$ ,  $\lambda_3x + \mu_3y$ .  $X, Y, Z$  respectively are multiples of these. To determine the multiples assume them arbitrarily. Then, by equating the coefficients of  $x^5$ ,  $5x^4y$ ,  $10x^2y^2$  in  $X^5 + Y^5 + Z^5$  to  $a, b, c$  respectively, we obtain three equations of the first degree for their determination.

And in like manner for the septic, nonic, &c.

The failing case when the coefficients are so specially connected that the canonizant has a square factor has been considered in § 204.

Ex. 7. If the canonizant of a quintic is a perfect cube the quintic can be reduced to the form  $(A_0, A_1, A_2, 0, 0, 0)(X, Y)^5$ ; and all invariants vanish.

Ans. Cf. § 204, and § 28, Ex. 5.



Now the necessary and sufficient condition for this is that, reversing the order of the columns,

$$\begin{vmatrix} a_0, a_1, a_2, \dots, a_n \\ a_1, a_2, a_3, \dots, a_{n+1} \\ a_2, a_3, a_4, \dots, a_{n+2} \\ \cdot \cdot \cdot \cdot \cdot \cdot \cdot \cdot \\ a_n, a_{n+1}, a_{n+2}, \dots, a_{2n} \end{vmatrix} = 0,$$

i. e. that the invariant defined as the *catalecticant* (§ 17 Examples) vanish.

Ex. 9. The binary quartic  $(a, b, c, d, e) (x, y)^4$  will be a sum of two fourth powers if

$$J \equiv ace + 2bcd - ad^2 - b^2e - c^3 = 0.$$

Ex. 10. The binary sextic  $(a, b, c, d, e, f, g) (x, y)^6$  will be a sum of three sixth powers if

$$\begin{vmatrix} a, b, c, d \\ b, c, d, e \\ c, d, e, f \\ d, e, f, g \end{vmatrix} = 0.$$

209.] **Catalecticants are invariants.** It is instructive to notice that what we have before us affords a proof that the catalecticant of a binary quantic of even order is an invariant. Its vanishing expresses the necessary and sufficient condition that the quantic may have a special property, that of being a sum of  $n$   $2n$ th powers, which is entirely independent of any linear transformation. If, in fact, the most general linear substitution possible in  $(a_0, a_1, a_2, \dots, a_{2n}) (x, y)^{2n}$  transforms that  $2n$ -ic into  $(A_0, A_1, A_2, \dots, A_{2n}) (X, Y)^{2n}$ , the vanishing of the same function of  $A_0, A_1, A_2, \dots, A_{2n}$  expresses the necessary and sufficient condition for the same special property. Moreover, the  $A$ 's being of the first degree in the  $a$ 's, the degree in the  $a$ 's of the catalecticants of the original and transformed  $2n$ -ics are the same. The one, then, can only differ from the other by a factor involving merely the constants of the substitution. The catalecticant is therefore an invariant by the definition (§ 3). That the factor is a power of the modulus has been established in general in § 23.

The student is advised to establish that the canonizant

of § 206 is a covariant of a binary  $(2n-1)$ -ic by similar reasoning.

210.] In § 208 it is proved that if the catalecticant vanish the  $2n$ -ic is a sum of  $n$   $2n$ th powers, and that conversely if a  $2n$ -ic is a sum of  $n$   $2n$ th powers its catalecticant vanishes. The latter fact is well exhibited as follows. For brevity of writing the case of the quartic alone is taken.

The catalecticant of  $p(x+ay)^4 + q(x+\beta y)^4$  is

$$\begin{vmatrix} p+q & , & pa+q\beta & , & pa^2+q\beta^2 \\ pa+q\beta & , & pa^2+q\beta^2 & , & pa^3+q\beta^3 \\ pa^2+q\beta^2 & , & pa^3+q\beta^3 & , & pa^4+q\beta^4 \end{vmatrix}.$$

Now this is a sum of  $8 (= 2^3)$  determinants, of which the first is

$$\begin{vmatrix} p & , & pa & , & pa^2 \\ pa & , & pa^2 & , & pa^3 \\ pa^2 & , & pa^3 & , & pa^4 \end{vmatrix},$$

the other seven being obtained from this one by replacing  $p$  and  $a$  by  $q$  and  $\beta$  in one or more of its columns.

But in every one of these eight determinants there are either two  $(p, a)$  columns at least or two  $(q, \beta)$  columns at least. Moreover, taking one which contains two  $(p, a)$  columns, we notice that the constituents in one of these two columns are either  $a$  or  $a^2$  times those in the other of the two, so that the determinant is either  $a$  or  $a^2$  times one with two columns identical, and therefore vanishes. Similarly every one of the eight which has two  $(q, \beta)$  columns vanishes. All the eight then vanish, and consequently their sum the catalecticant vanishes.

211.] **Canonical form of quartic.** We now proceed to show that the general binary quartic may be reduced to the canonical form

$$X^4 + Y^4 + 6mX^2Y^2,$$

in favour of which there is a presumption as the form contains one  $(= 4-3, \text{ cf. } \S 196)$  free coefficient  $m$ .

We have to see that  $p, q, a, \beta$  and  $\mu$  can be found so as to make

$$\begin{aligned} (a, b, c, d, e)(x, y)^4 \\ \equiv p(x+ay)^4 + q(x+\beta y)^4 + 6\mu(x+ay)^2(x+\beta y)^2 \end{aligned}$$

an identity. It is not enough to notice that the number of free constants on the right is equal to the number of coefficients on the left.

Knowledge that a quartic equation has four roots is assumed. Thus  $(a, b, c, d, e)(x, y)^4$  can be broken up into linear factors, and consequently into two quadratic factors in three ways, corresponding to the arrangements (12, 34), (13, 24), (14, 23) of the linear factors. Let one of the quadratic factorizations be

$$(a'x^2 + 2b'xy + c'y^2)(a''x^2 + 2b''xy + c''y^2).$$

What we have to prove will be established if we can find  $p', q', p'', q'', \alpha, \beta$  so that simultaneously

$$\begin{aligned} a'x^2 + 2b'xy + c'y^2 &\equiv p'(x + \alpha y)^2 + q'(x + \beta y)^2, \\ a''x^2 + 2b''xy + c''y^2 &\equiv p''(x + \alpha y)^2 + q''(x + \beta y)^2. \end{aligned}$$

The six equations for finding the constants on the right, that this may be the case, are

$$\begin{aligned} p' + q' &= a', & p'' + q'' &= a'', \\ p'\alpha + q'\beta &= b', & p''\alpha + q''\beta &= b'', \\ p'\alpha^2 + q'\beta^2 &= c', & p''\alpha^2 + q''\beta^2 &= c''. \end{aligned}$$

Now of these the first three are consistent for the determination of  $p', q'$  if

$$c' - b'(a + \beta) + a'\alpha\beta = 0,$$

as we see by eliminating  $p'$  and  $q'$ , and the last three are consistent for the determination of  $p'', q''$  if

$$c'' - b''(a + \beta) + a''\alpha\beta = 0;$$

and these two conditions are satisfied by taking

$$\frac{1}{a'b'' - a''b'} = \frac{\alpha + \beta}{a'c'' - a''c'} = \frac{\alpha\beta}{b'c'' - b''c'},$$

i.e. by taking for  $\alpha, \beta$  the roots of the quadratic

$$(a'b'' - a''b')\alpha^2 - (a'c'' - a''c')\alpha + b'c'' - b''c' = 0,$$

which are finite and unequal if  $a', b', c', a'', b'', c''$  are unconnected, i.e. if the quartic is general.

We see then that the required reduction is possible, and possible in three distinct ways, one corresponding to each way of breaking up the quartic into two quadratic factors.



The quartic in the form

$$p(x+ay)^4 + q(x+\beta y)^4 + 6\mu(x+ay)^2(x+\beta y)^2,$$

into which it is now shown capable of being thrown, is given the canonical form

$$X^4 + Y^4 + 6mX^2Y^2$$

by taking for  $X$  either  $\pm p^{\frac{1}{4}}(x+ay)$  or  $\pm \sqrt{-1}p^{\frac{1}{4}}(x+ay)$ , and for  $Y$  either  $\pm q^{\frac{1}{4}}(x+\beta y)$  or  $\pm \sqrt{-1}q^{\frac{1}{4}}(x+\beta y)$ . It is thus seen that  $m$  may have either of the two values  $\pm (pq)^{-\frac{1}{2}}\mu$ . Each of the three reductions above produces then two varieties of the canonical form differing only in the sign of  $m$ . The equation to be found for the determination of  $m$  should consequently prove to be a cubic in  $m^2$ .

We thus encounter a striking difference between the quartic, and other quantics of even order, and the cubic, and quantics of odd order, in the matter of canonical forms. The reduction of the cubic to its canonical form  $X^3 + Y^3$  is unique. On the other hand, the reduction of the quartic to its canonical form  $X^4 + Y^4 + 6mX^2Y^2$  is sixfold.

212.] We now proceed to exhibit the information given by invariant algebra with reference to the general binary quartic

$$u \equiv (a, b, c, d, e)(x, y)^4$$

and its canonical form

$$X^4 + Y^4 + 6mX^2Y^2 \equiv (1, 0, m, 0, 1)(X, Y)^4.$$

Suppose that  $X$  and  $Y$ , expressed in terms of  $x$  and  $y$ , are  $\lambda x + \mu y$  and  $\lambda'x + \mu'y$  respectively. Let  $M'$  denote  $\lambda\mu' - \lambda'\mu$ , so that  $M'$  is the modulus of the substitution which reduces the canonical form to the general, and consequently  $M'^{-1}$  the modulus of that which reduces the general to the canonical.

By § 170 the irreducible concomitants of the quartic are, including itself, five in number. They are

(1) the quartic itself

$$u \equiv (a, b, c, d, e)(x, y)^4,$$

(2, 3) its two invariants

$$I \equiv ae - 4bd + 3c^2,$$

$$J \equiv ace + 2bcd - ad^2 - b^2e - c^3,$$

of which the latter is its catalecticant,

(4) its quartic covariant, or Hessian,

$$H \equiv \begin{vmatrix} ax^2 + 2bxy + cy^2, & bx^2 + 2cxy + dy^2 \\ bx^2 + 2cxy + dy^2, & cx^2 + 2dxy + ey^2 \end{vmatrix},$$

of which the seminvariant leader is  $ac - b^2$ , and

(5) a sextic covariant of which the seminvariant leader is  $a^2d - 3abc + 2b^3$ , which, written at length by the method of § 110, is

$$\begin{aligned} G \equiv & (a^2d - 3abc + 2b^3)x^6 + (a^2e + 2abd - 9ac^2 + 6b^2c)x^5y \\ & + (5abe - 15acd + 10b^2d)x^4y^2 + (10b^2e - 10ad^2)x^3y^3 \\ & + (15bce - 5ade - 10bd^2)x^2y^4 + (9c^2e - ae^2 - 2bde - 6cd^2)xy^5 \\ & + (3cde - be^2 - 2d^3)y^6. \end{aligned}$$

The power of the modulus in the equality expressive of invariancy or covariancy of any one of these has for its index, it will be remembered, the weight of the invariant or of the seminvariant leader of the covariant. Thus we have the five equalities

$$u = X^4 + Y^4 + 6mX^2Y^2, \quad \dots (1)$$

$$I = M^4(1 + 3m^2), \quad \dots (2)$$

$$J = M^6(m - m^3), \quad \dots (3)$$

$$H = M^2 \{m(X^4 + Y^4) + (1 - 3m^2)X^2Y^2\}, \quad \dots (4)$$

$$G = M^3(1 - 9m^2)XY(X^4 - Y^4). \quad \dots (5)$$

The first observation made on an inspection of these equalities is that the two invariants  $I$  and  $J$  alone supply us with the equation for the determination of the  $m$ 's of the six canonical forms, and with the values of the modulus  $M'$ , going with each value of  $m$ , of the substitutions which express  $X$  and  $Y$  in terms of  $x$  and  $y$ .

Elimination of  $M'$  between (2) and (3) gives at once

$$I^3m^2(1 - m^2)^2 = J^2(1 + 3m^2)^3, \quad \dots (6)$$

the cubic whose roots are the three values of  $m^2$ . To each value of  $m$  there corresponds a value of  $M'^2$  given by

$$M'^2 = \frac{J}{I} \cdot \frac{1 + 3m^2}{m - m^3}, \quad \dots (7)$$

so that with each value of  $m$  go two of  $M'$ , equal but of

opposite signs. This is reasonable, for a canonical form is unaltered when we interchange  $X$  and  $Y$ , but the modulus is changed in sign. The equation for all values of  $M'$  should then be a cubic in  $M'^4$ , the two values  $\pm m$  of  $m$  giving two values  $\pm M'^2$  of  $M'^2$ . This cubic comes at once from elimination of  $m$  between (2) and (3), and is

$$J^2 = M'^{12} \frac{I - M'^4}{3M'^4} \left\{ 1 - \left( \frac{I - M'^4}{3M'^4} \right) \right\}^2,$$

i.e.  $(M'^4 - I)(4M'^4 - I)^2 + 27J^2 = 0. \quad \dots (8)$

The cubic for  $M'^2 m$  is simpler than either that for  $m^2$  or that for  $M'^4$ , and is given by taking

$$\frac{J}{M'^2 m} - I = -4M'^4 m^2,$$

and so is

$$4(M'^2 m)^3 - I(M'^2 m) + J = 0, \quad \dots (9)$$

which will be recognized as the ordinary reducing cubic of a quartic equation.

We shall consider this cubic further presently. Meanwhile let us pay a little close attention to the cubic (6) for  $m^2$ , the solution of which is the one which at once affords the canonical forms themselves. Written at length the cubic is

$$(I^3 - 27J^2)m^6 - (2I^3 + 27J^2)m^4 + (I^3 - 9J^2)m^2 - J^2 = 0. \dots (10)$$

The one parameter involved in it is the absolute invariant  $\frac{I^3}{J^2}$ .

We proceed to draw in the following article certain conclusions as to the reduction of special classes of quartics, which obey invariant conditions suggested by the coefficients in this cubic.

213.] **Quartics for which  $I = 0$ .** If a quartic belong to the special class for which

$$I \equiv ae - 4bd + 3c^2 = 0,$$

the cubic for  $m^2$  becomes

$$(1 + 3m^2)^3 = 0,$$

so that the three pairs of reductions to a canonical form

coalesce in form into a single pair, the alternative canonical forms being

$$X^4 + Y^4 \pm 2\sqrt{-3} X^2 Y^2,$$

and being thus of imaginary shape. From (3) we find, as corresponding to the values  $\pm \sqrt{-\frac{1}{3}}$  of  $m$  respectively,

$$M^6 = \mp \frac{3}{4} \sqrt{-3} J,$$

so that the values of the modulus as well as of  $m$  are all imaginary.

Since the relation  $1 + 3m^2 = 0$  may be written

$$\frac{1 - 3m^2}{6m} = -\frac{m}{1},$$

we see from (1) and (4) that the Hessian of

$$X^4 + Y^4 \pm 2\sqrt{-3} X^2 Y^2$$

is, but for the factor  $M'^2 m$ ,

$$X^4 + Y^4 \mp 2\sqrt{-3} X^2 Y^2.$$

Thus a quartic for which  $I = 0$  and its Hessian have reciprocal properties, each being, but for a constant factor, the Hessian of the other. Moreover they have, but for a constant factor, the same sextic covariant  $G$ .

Ex. 11. Prove that

$$\begin{aligned} X^4 + Y^4 \pm 2\sqrt{-3} X^2 Y^2 &\equiv \frac{1}{2 \mp 2\sqrt{-3}} \{ (X+Y)^4 + (X-Y)^4 \\ &\quad \mp 2\sqrt{-3} (X+Y)^2 (X-Y)^2 \} \\ &\equiv \frac{1}{2 \pm 2\sqrt{-3}} \{ (X+Y\sqrt{-1})^4 \\ &\quad + (X-Y\sqrt{-1})^4 \pm 2\sqrt{-3} (X+Y\sqrt{-1})^2 (X-Y\sqrt{-1})^2 \}, \end{aligned}$$

thus exhibiting the connexion between the three like pairs of canonical forms when  $I = 0$ .

Ex. 12. Three of the six anharmonic ratios of the range or pencil denoted by a binary quartic for which  $I = 0$  are equal to  $-\omega$ , and the other three to  $-\omega^2$ , where  $\omega$  and  $\omega^2$  are the imaginary cube roots of unity.

Ex. 13. When  $I = 0$ ,  $M^6 m^3 = -\frac{1}{4} J$ .

Ex. 14. When  $I = 0$ ,

$$u + \frac{4^{\frac{1}{3}}}{J^{\frac{1}{3}}} H, u + \omega \frac{4^{\frac{1}{3}}}{J^{\frac{1}{3}}} H, u + \omega^2 \frac{4^{\frac{1}{3}}}{J^{\frac{1}{3}}} H$$

are perfect squares; viz. numerical multiples of the squares of the products  $XY$  for canonical forms.

Ex. 15. When  $I = 0$ ,  $Jw^3 + 4H^3 = -G^2$ . Hence also prove Ex. 14.

214.] **Quartics for which  $J = 0$ .** When the catalecticant

$$J \equiv ace + 2bcd - ad^2 - b^2e - c^3 = 0,$$

so that (§ 208) one canonical form is a sum of two fourth powers, the cubic (6) or (10) of § 212 for  $m^2$  is

$$m^2(m^2 - 1)^2 = 0.$$

The second and third pairs of canonical forms coalesce then in the shape

$$X^4 + Y^4 \pm 6X^2Y^2.$$

The connexion of the different canonical forms for this case is exhibited in the identities

$$\begin{aligned} X^4 + Y^4 &\equiv X^4 + (Y\sqrt{-1})^4, \\ &\equiv \frac{1}{8} \{ (X+Y)^4 + (X-Y)^4 + 6(X+Y)^2(X-Y)^2 \}, \\ &\equiv \frac{1}{8} \{ (X+Y)^4 + (X\sqrt{-1} - Y\sqrt{-1})^4 \\ &\quad - 6(X+Y)^2(X\sqrt{-1} - Y\sqrt{-1})^2 \}, \\ &\equiv \frac{1}{8} \{ (X+Y\sqrt{-1})^4 + (X-Y\sqrt{-1})^4 \\ &\quad + 6(X+Y\sqrt{-1})^2(X-Y\sqrt{-1})^2 \}, \\ &\equiv \frac{1}{8} \{ (X+Y\sqrt{-1})^4 + (X\sqrt{-1} + Y)^4 \\ &\quad - 6(X+Y\sqrt{-1})^2(X\sqrt{-1} + Y)^2 \}. \end{aligned}$$

By § 212 (2),  $M'^4 = I$  goes with  $m = 0$ , and  $M'^4 = \frac{1}{4}I$  with  $m = \pm 1$ .

Ex. 16. When  $J = 0$ , the Hessian is, but for a constant factor, the square of the product of the  $X$  and  $Y$  of the canonical form  $X^4 + Y^4$ .

Ex. 17. In the same case, the two expressions  $u \pm 2I^{-\frac{1}{3}}H$  are eight times the squares of the products  $XY$  for the other two essentially distinct canonical forms, the third and fourth, and the fifth and sixth, of the above forms not being reckoned as essentially distinct.

Ex. 18. In the same case,  $(Iu^2 - 4H^2)H = G^2$ .

Ex. 19. The range or pencil denoted by a binary quartic for which  $J = 0$  is harmonic.

Ex. 20. So is the range or pencil composed of any two out of three pairs of elements which constitute what is denoted by the sextic covariant  $G$ .

215.] **Quartics for which  $I^3 = 27J^2$ .** When the coefficients in the quartic are such that  $I^3 - 27J^2 = 0$ , i. e. when the discriminant vanishes, so that the quartic has a square factor, one value of  $m^2$  given by § 212 (10) is infinite, and the quadratic for the other two values of  $m^2$  is

$$(9m^2 - 1)^2 = 0.$$

But we are here confronted with a case in which the reduction to the canonical form  $X^4 + Y^4 + 6mX^2Y^2$  is impossible unless a further condition is satisfied. The value  $m = \infty$  would make this canonical form an infinite multiple of  $X^2Y^2$ , i. e. of a perfect square, and the values  $m = \pm \frac{1}{3}$  would make it  $X^4 + Y^4 \pm 2X^2Y^2$ , again perfect squares. Now obviously a quartic with a square factor must have its conjugate quadratic factor also a perfect square for such a reduction to be possible.

For the explanation of this we must refer back to § 211. If the two conjugate quadratic factors there assumed have a common linear factor their eliminant

$$(a'c'' - a''c')^2 - 4(a'b'' - a''b')(b'c'' - b''c')$$

vanishes, and the quadratic in  $a$  has equal roots, so that  $a = \beta$ , and the method followed fails to find a distinct  $X$  and  $Y$ , and indeed fails to lead to any result which is not more obvious otherwise. And again, if one of the quadratic factors,  $a'x^2 + 2b'xy + c'y^2$  suppose, is a perfect square, so that  $a'c' = b'^2$ , it follows that  $p'q'(a - \beta)^2 = 0$ , so that either  $a = \beta$ , and there is failure as before, or else either  $p' = 0$  or  $q' = 0$ , which leads not to the form  $X^4 + Y^4 + 6mX^2Y^2$  but to the form  $X^4 + 6mX^2Y^2$ .

It is this form, or rather its further simplification

$$X^4 + 6X^2Y^2,$$

which is canonical for a quartic for which  $I^3 - 27J^2 = 0$ .

The more special quartic still which has not only one square factor but two square factors, i. e. which is a constant

multiple of the square of a quadratic, can however, it is clear, be given the form  $(X^2 + Y^2)^2$  or the form  $(X^2 - Y^2)^2$  as above. An even simpler form for such a quantic is  $6X^2Y^2$ .

Ex. 21. The Hessian of a binary quartic with a square factor has that same square factor. (This fact is easily proved for a binary quantic of any order with a square factor.)

Ex. 22. The sextic covariant  $G$  of a quartic with a square factor  $(lx + my)^2$  has the factor  $(lx + my)^5$ .

Ex. 23. If a binary quartic be the square of a quadratic it is the same but for a constant factor as its Hessian, so that

$$\frac{ac - b^2}{a} = \frac{ad - bc}{2b} = \frac{ae + 2bd - 3c^2}{6c} = \frac{be - cd}{2d} = \frac{ce - d^2}{e}.$$

Ans.  $2IH = 3Ju$ . (Cayley.)

Ex. 24. In the same case the sextic covariant  $G$  vanishes identically. Hence also determine the same conditions as in Ex. 23.

216.] **The general binary quartic.** We now proceed to apply the equalities of § 212 to the case of the general quartic. A pair of canonical forms  $X^4 + Y^4 \pm 6mX^2Y^2$  are not essentially distinct, the  $X$  and  $Y$  of one being merely the  $X$  and  $Y\sqrt{-1}$  of the other.

The sextic covariant  $G$  helps us to decide what are the  $X$  and  $Y$  of each of the two other essentially distinct canonical forms of the same shape as one  $X^4 + Y^4 + 6mX^2Y^2$ . This covariant  $G$  has, § 212 (5), the  $X$  and  $Y$  of  $X^4 + Y^4 + 6mX^2Y^2$  for factors. For the same reason it must have for factors the  $X$  and  $Y$  of each of the other canonical forms. It is in fact, therefore, but for a factor free from the variables, the product of the three  $X$ 's and the three  $Y$ 's of the essentially distinct canonical forms. We are thus led to expect that  $X^2 - Y^2$  and  $X^2 + Y^2$  are but for constant factors the products  $XY$  corresponding to the two other forms.

And it is in fact quite easy so to assign  $k', k'', m', m''$  as to satisfy the identities

$$\begin{aligned} X^4 + Y^4 + 6mX^2Y^2 &\equiv \{k'(X + Y)\}^4 + \{k'(X - Y)\}^4 \\ &\quad + 6m'\{k'(X + Y)\}^2\{k'(X - Y)\}^2 \\ &\equiv \{k''(X + Y\sqrt{-1})\}^4 + \{k''(X - Y\sqrt{-1})\}^4 \\ &\quad + 6m''\{k''(X + Y\sqrt{-1})\}^2\{k''(X - Y\sqrt{-1})\}^2. \end{aligned}$$

Thus for  $k', m'$  we have only to secure that

$$k'^4(2 + 6m') = 1, k'^4(12 - 12m') = 6m,$$

i. e. to take

$$m' = \frac{1-m}{3m+1}, k'^4 = \frac{1}{8}(3m+1),$$

and similarly for  $k'', m''$ .

217.] **Reduction of general quartic to canonical form.** Let us now take the simplest of the cubics of § 212, viz. (9)

$$4(M'^2m)^3 - I(M'^2m) + J = 0.$$

The solution of this affords a ready way of determining the  $X$  and  $Y$  of either of the canonical forms. The equations (1) and (4) of § 212 give us at once

$$u - \frac{H}{M'^2m} = \left(9m - \frac{1}{m}\right) X^2 Y^2.$$

Solve then the cubic above for  $M'^2m$ , and, taking either of its roots, form the corresponding  $M'^2mu - H$ . This, but for a multiplier free from the variables, is a perfect square, namely the square of  $XY$ . Break up the square root of  $M'^2mu - H$ , or any convenient multiple of it, into two factors  $gx + hy, g'x + h'y$ . The identity must hold

$$u \equiv (a, b, c, d, e) (x, y)^4 \equiv a'(gx + hy)^4 + 6c'(gx + hy)^2(g'x + h'y)^2 + e'(g'x + h'y)^4,$$

for some values of  $a', c', e'$ . These values can be found by identifying three of the coefficients on the left with those which correspond on the right. Having found them,

$$a'^{\frac{1}{4}}(gx + hy), e'^{\frac{1}{4}}(g'x + h'y) \text{ are } X, Y, \text{ and } c'a'^{-\frac{1}{2}}e'^{-\frac{1}{2}} \text{ is } m.$$

218.] **Syzygy among  $u, I, J, H, G$ .** That  $XY$  is, but for a factor free from  $x, y$ , the square root of  $M'^2mu - H$ , and consequently that the product of the three values of  $XY$  for the three essentially distinct canonical forms is, but for such a factor, the square root of the product of the three values of  $M'^2mu - H$  corresponding to the three roots of the cubic for  $M'^2m$ , tells us, when taken in connexion with § 216, that this product

$$(M_1'^2m_1u - H)(M_2'^2m_2u - H)(M_3'^2m_3u - H)$$

can only differ by a factor free from the variables from  $G^2$ .



Now by the theory of equations

$$4z^3 - Iz + J = 4(z - M_1'^2 m_1)(z - M_2'^2 m_2)(z - M_3'^2 m_3).$$

Consequently an identity must hold of the form

$$kG^2 = -4H^3 + Iu^2H - Ju^3.$$

That  $k$  here is a merely numerical constant, and not a function of the coefficients, is clear because  $G^2$  and the right-hand side are both of degree 6 in the coefficients. To find its value we may either substitute for  $u, I, J, H, G$  their values in terms of  $X, Y, M', m$  from § 212 and examine the identity of the coefficients of one term, say of  $X^{10}Y^2$ , on the two sides, or may notice that § 213, Ex. 15 gives us the particular form which the relation takes when  $I = 0$ . We thus find  $k = 1$ .

Accordingly the irreducible but not independent concomitants  $u, I, J, H, G$  of the binary quartic  $u$  are connected by the syzygy

$$Iu^2H - 4H^3 - Ju^3 = G^2,$$

the invariants and seminvariant leaders of the covariants being themselves connected by the syzygy

$$Ia^2(ac - b^2) - 4(ac - b^2)^3 - Ja^3 = (a^2d - 3abc + 2b^3)^2.$$

These syzygies have been otherwise obtained in previous chapters.

219.] **Canonical reduction with unit modulus.** There is often convenience in using, not the strictly canonical form of a quantic, i. e. the simplest form to which the quantic may be reduced by any linear substitution, but the simplest form to which it may be reduced by a substitution of unit modulus.

If the substitution which reduces the quartic  $(a, b, c, d, e)(x, y)^4$  to its canonical form  $X^4 + Y^4 + 6mX^2Y^2$  be

$$x = lX + mY, \quad y = l'X + m'Y,$$

so that  $lm' - l'm \equiv M'^{-1}$  in what precedes, the substitution

$$(lm' - l'm)^{\frac{1}{2}}x = lx' + my', \quad (lm' - l'm)^{\frac{1}{2}}y = l'x' + m'y',$$

whose modulus is unity, reduces the quartic to

$$M'^2(x'^4 + y'^4 + 6mx'^2y'^2).$$

Thus  $a'(x'^2 + y'^2) + 6c'x'^2y'^2 \equiv (a', 0, c', 0, a')(x', y')^2$

is a form to which the general binary quartic can be reduced by a substitution whose modulus is unity. We also see that  $c'$  is the  $M'^2m$  of §§ 212, 217, so that the reducing cubic of the quartic is the one whose roots are the three values of  $c'$ . We may find it very easily as follows. Since the modulus is 1,

$$I = a'^2 + 3c'^2,$$

$$J = a'^2c' - c'^3,$$

whence, eliminating  $a'^2$ ,

$$Ic' - J = 4c'^3,$$

the same cubic as that already found for  $M'^2m$ .

We might equally have found in this way the same cubic for  $c'$  if all that we had assumed were that a substitution of unit modulus reduces the quartic to the form

$$a'x'^4 + 6c'x'^2y'^2 + e'y'^4.$$

Adopting, however, the fact known as above that  $a'$  and  $e'$  may be made equal, the equalities (1) to (5) of § 212 are replaced by

$$u = a'(x'^4 + y'^4) + 6c'x'^2y'^2,$$

$$I = a'^2 + 3c'^2,$$

$$J = a'^2c' - c'^3,$$

$$H = a'c'(x'^4 + y'^4) + (a'^2 - 3c'^2)x'^2y'^2,$$

$$G = a'(a'^2 - 9c'^2)x'y'(x'^4 - y'^4),$$

whose right-hand sides are obviously connected by a syzygy, as they involve only four quantities  $a', c', x', y'$ . This is readily seen to be that of the preceding article.

Ex. 25. Prove that, if  $H'$  be the Hessian of the Hessian  $H$  of  $u$ ,  $H' = \frac{1}{4}Ju - \frac{1}{12}IH$ .

Ex. 26. The sextic covariant  $G'$  of  $H$  is  $-\frac{1}{4}JG$ .

Ex. 27. Find the Hessian and the sextic covariant of  $ku + k'H$ .

220.] The cubic for  $c'$  may be found in a different manner, which exhibits it in a form having its analogue in the case of higher binary quantics of even order.

It has been seen that the identity

$$\begin{aligned} ax^4 + 4bx^3y + 6cx^2y^2 + 4dxy^3 + ey^4 \\ \equiv a'(lx + my)^4 + e'(l'x + m'y)^4 + 6c'(lx + my)^2(l'x + m'y)^2 \end{aligned}$$

is one which can be satisfied simultaneously with

$$lm' - l'm = 1,$$

and in fact that we may take  $e' = a'$ .

Now operate on both sides of the identity with

$$\left(l \frac{d}{dy} - m \frac{d}{dx}\right) \left(l' \frac{d}{dy} - m' \frac{d}{dx}\right),$$

noticing that this annihilates both  $lx + my$  and  $l'x + m'y$ , and that

$$\begin{aligned} \left(l \frac{d}{dy} - m \frac{d}{dx}\right) \left(l' \frac{d}{dy} - m' \frac{d}{dx}\right) \{ (lx + my)^2 (l'x + m'y)^2 \} \\ = -4(lm' - l'm)^2 (lx + my) (l'x + m'y) \\ = -4(lx + my) (l'x + m'y). \end{aligned}$$

Equating coefficients of  $x^2$ ,  $xy$ ,  $y^2$  in the results of operating on the left and right-hand sides of the identity, we obtain

$$\begin{aligned} amm' - b(lm' + l'm) + cll' &= -2c'll', \\ bmm' - c(lm' + l'm) + dll' &= -c'(lm' + l'm), \\ cmm' - d(lm' + l'm) + ell' &= -2c'mm', \end{aligned}$$

equations linear in  $ll'$ ,  $lm' + l'm$ ,  $mm'$ . By elimination of these we at once obtain the cubic for  $c'$

$$\begin{vmatrix} a & , & b & , & c + 2c' \\ b & , & c - c' & , & d \\ c + 2c' & , & d & , & e \end{vmatrix} = 0,$$

$$\text{i. e. } ace + 2bcd - ad^2 - b^2e - c^3 - (ae - 4bd + 3c^2)c' + 4c'^3 = 0,$$

$$\text{i. e. } J - Ic'_e + 4c'^3 = 0,$$

the reducing cubic already obtained otherwise.

Taking either root  $c'$  of this cubic we can solve the linear equations in  $ll'$ ,  $lm' + l'm$ ,  $mm'$ , and so obtain their ratios, i. e. the product  $ll'x^2 + (lm' + l'm)xy + mm'y^2$  of  $lx + my$ ,  $l'x + m'y$  but for a constant factor.

221.] **Solution of a quartic equation.** When a quartic is reduced to its canonical form, or to the form  $a'x^4 + 6c'x^2y^2 + e'y^4$ , it is at once broken up into quadratic factors and solved. Two methods suggested by the articles which precede are here

exemplified. They are not so simple in their use as some of those given in works on the theory of equations.

Ex. 28. By use of § 220 solve the quartic equation

$$3x^4 - 4x^3 + 24x^2 - 16x + 48 = 0.$$

Here  $a, b, c, d, e$  have the values 3, -1, 4, -4, 48. Thus  $I = 176, J = 448$ , and the cubic for  $c'$  is  $c'^3 - 44c' + 112 = 0$ , of which 4 is a root. The corresponding ratios  $l'l' : lm' + l'm : mm'$  of § 220 are 1 : 0 : -4. The quartic has then the form

$$a''(x+2)^4 + e''(x-2)^4 + 6c''(x^2-4)^2 = 0,$$

and is in fact seen to be

$$(x+2)^4 + 2(x-2)^4 + 3(x^2-4)^2 = 0,$$

i.e.  $\{(x+2)^2 + (x-2)^2\} \{(x+2)^2 + 2(x-2)^2\} = 0,$

i.e.  $(x^2+4)(3x^2-4x+12) = 0,$

so that the roots are  $\pm 2\sqrt{-1}$  and  $\frac{1}{3}(2 \pm 4\sqrt{-2})$ .

Ex. 29. To the same quartic apply the method of § 217.

Here

$$\begin{aligned} H &\equiv (3x^2 - 2x + 4)(4x^2 - 8x + 48) - (-x^2 + 8x - 4)^2 \\ &\equiv 11x^4 - 16x^3 + 104x^2 - 64x + 176. \end{aligned}$$

Also a value of  $M'^2m$ , or  $c'$ , is, as above, 4. Thus

$$c'u - H \equiv x^4 - 8x^2 + 16 \equiv \{(x+2)(x-2)\}^2.$$

Hence the given equation has the form

$$a''(x+2)^4 + e''(x-2)^4 + 6c''(x^2-4)^2 = 0,$$

and the solution is completed as above.

222.] There is a symmetrical expression given by Cayley for a linear factor of the general binary quartic  $u$ .

If  $c_1, c_2, c_3$  are the roots of  $4c'^3 - Ic' + J = 0$ , it has been seen in § 217 that  $c_1u - H, c_2u - H, c_3u - H$  are multiples of squares of quadratics in the variables.

Thus

$$\lambda\sqrt{c_1u - H} + \mu\sqrt{c_2u - H} + \nu\sqrt{c_3u - H}$$

is a rational quadratic function of  $x$  and  $y$ . We seek  $\lambda, \mu, \nu$  that it may be the square of a factor of  $u$ .

A value of  $\frac{x}{y}$  which makes  $u$  vanish, i. e. a root of  $u$ , will make the quadratic function vanish if

$$(\lambda + \mu + \nu) \sqrt{-H} = 0,$$

i. e. if  $\lambda + \mu + \nu = 0$ .

The same value will make its differential coefficient with respect to  $x$ , i. e.

$$\frac{\lambda}{\sqrt{c_1 u - H}} \left\{ c_1 \frac{du}{dx} - \frac{dH}{dx} \right\} + \frac{\mu}{\sqrt{c_2 u - H}} \left\{ c_2 \frac{du}{dx} - \frac{dH}{dx} \right\} + \frac{\nu}{\sqrt{c_3 u - H}} \left\{ c_3 \frac{du}{dx} - \frac{dH}{dx} \right\},$$

vanish, if it make

$$\frac{1}{\sqrt{-H}} \left\{ (\lambda c_1 + \mu c_2 + \nu c_3) \frac{du}{dx} - (\lambda + \mu + \nu) \frac{dH}{dx} \right\} = 0,$$

as it will do if the further condition

$$\lambda c_1 + \mu c_2 + \nu c_3 = 0$$

is satisfied.

Now these two conditions are satisfied by taking

$$\frac{\lambda}{c_2 - c_3} = \frac{\mu}{c_3 - c_1} = \frac{\nu}{c_1 - c_2}.$$

We conclude then that

$$(c_2 - c_3) \sqrt{c_1 u - H} + (c_3 - c_1) \sqrt{c_2 u - H} + (c_1 - c_2) \sqrt{c_3 u - H}$$

is, but for a multiple not involving the variables, the square of a linear factor of the quartic  $u$ .

Ex. 30. Prove that

$(c_2 - c_3) \sqrt{c_1(c_1 u - H)} + (c_3 - c_1) \sqrt{c_2(c_2 u - H)} + (c_1 - c_2) \sqrt{c_3(c_3 u - H)}$   
is the square of a linear factor of the Hessian.

223.] **Geometry of concomitants of quartic.** The invariant and covariant geometry of a binary quartic is a geometry of anharmonic properties. The student of geometry will know that, if  $\rho$  is an anharmonic ratio of a pencil or range of four elements, the other five anharmonic ratios are

$$\frac{1}{\rho}, 1 - \rho, \frac{1}{1 - \rho}, 1 - \frac{1}{\rho}, \frac{\rho}{\rho - 1}.$$

Some parts of the geometry have already been obtained. Thus (§ 213, Ex. 12)  $I = 0$  is the condition that one anharmonic ratio of the pencil or range denoted by the quartic be  $-\omega$ , and consequently that two others be also  $-\omega$ , and the other three  $-\omega^2$ . In fact, if we take for  $\rho$  the anharmonic ratio  $\frac{\gamma-\beta}{\gamma-\delta} : \frac{\alpha-\beta}{\alpha-\delta}$ , where  $\alpha, \beta, \gamma, \delta$  are the roots, and notice that this is  $-\frac{u}{w}$  in the notation of § 80, we obtain

$$\begin{aligned} I &= \frac{1}{24} a^2 (u^2 + v^2 + w^2) \\ &= \frac{1}{24} a^2 (u^2 + v^2 - 2vw) \\ &= \frac{1}{12} a^2 \{u^2 + w(u+w)\} \\ &= \frac{1}{12} a^2 w^2 (\rho^2 - \rho + 1), \end{aligned}$$

$$\begin{aligned} u &= (\beta - \gamma) \\ w &= (\gamma - \alpha) \end{aligned}$$

so that  $I = 0$  means, unless two roots are equal which would imply a further invariant condition,

$$\rho = -\omega \text{ or } -\omega^2.$$

Again,  $J = 0$  is the condition (§ 214, Ex. 19) that the pencil or range be harmonic, i.e. that one anharmonic ratio be  $-1$ , and consequently the rest  $-1, 2, \frac{1}{2}, \frac{1}{2}, 2$ . In fact, referring again to § 80,

$$\begin{aligned} J &= -\frac{1}{432} a^3 (v-w)(w-u)(u-v) \\ &= -\frac{1}{432} a^3 (-u-2w)(w-u)(2u+w) \\ &= \frac{1}{432} a^3 w^3 (\rho-2)(\rho+1)(2\rho-1), \end{aligned}$$

so that  $J = 0$  necessitates that  $\rho$  be either  $-1$  or  $2$  or  $\frac{1}{2}$ .

We have at once, by elimination of  $aw$ , the equation whose roots are the six values of  $\rho$ , i.e. the six anharmonic ratios of the general quartic; namely,

$$\frac{I^3}{J^2} = 108 \frac{(\rho^2 - \rho + 1)^3}{(\rho + 1)^2 (\rho - 2)^2 (2\rho - 1)^2},$$

which may be given the simpler shape

$$\frac{(\rho^2 - \rho + 1)^3}{\rho^2(\rho - 1)^2} = \frac{27I^3}{4(I^3 - 27J^2)},$$

a cubic, in  $\rho(1-\rho)$ . The left-hand side of this may also be written

$$\frac{\left(\rho + \frac{1}{\rho} - 1\right)^3}{\rho + \frac{1}{\rho} - 2},$$

so that it is also a cubic in  $\rho + \frac{1}{\rho}$ .

To interpret the sextic covariant  $G$  we remember that it is, but for a constant factor, the product of the  $XY$ 's of the three essentially distinct canonical forms. Now, if the canonical form  $X^4 + 6mX^2Y^2 + Y^4$  be broken up into

$$(X^2 + \mu Y^2)(X^2 + \mu^{-1} Y^2),$$

we recognize that  $XY$  represents the common pair of harmonic conjugates of the pairs  $X^2 + \mu Y^2$ ,  $X^2 + \mu^{-1} Y^2$ . We thus conclude that  $G$  represents the three pairs of common harmonic conjugates of pairs into which the four factors of the quartic  $u$  can be separated, i.e. the double elements of the three involutions which are determined by taking the four linear factors of  $u$  in pairs.

It is clear, from the similarity of the canonical reduction (§ 212 (4)) of  $H$  the Hessian of  $u$  to that of  $u$  itself, that  $G$  has the same property with regard to  $H$  as it has with regard to  $u$ . It has also the same property with regard to  $ku + k'H$ , where  $k$  and  $k'$  have any values which do not make this a perfect square.

We notice the further property of  $G$ , gathering it from the canonical reduction, that its six linear factors occur in pairs  $XY$ ,  $X^2 - Y^2$ ,  $X^2 + Y^2$  such that either pair constitutes the double elements of the involution determined by the other pairs of elements.

The geometrical property of  $H$  is that of determining with  $u$  an infinite system of quartics  $ku + k'H$ , the factors of any one of which, taken in pairs in any way, have a pair chosen out of six elements constituting  $G$  for the double elements of the involution which they determine.

Ex. 31. Find a covariant which represents the four harmonic conjugates of the factors of  $u$ , each with regard to the Hessian of the other three factors.

*Ans.*  $(I^3 - 3J^2)u + 8IJH$ . To find it take the quartic in the form  $4(x^3 + y^3)(x + ay)$ , so that  $I = 12a$ ,  $J = -4(a^3 + 1)$ , and determine  $\theta$ , so that  $\theta u - H$ , where  $H$  is the Hessian of the quartic, may have  $x - ay$  for a factor.

Ex. 32. In terms of the roots of  $u$ , the quadratic factors of  $G$  which give the products  $XY$  for canonical forms are

$$(\delta + a - \beta - \gamma, \beta\gamma - a\delta, \delta a(\beta + \gamma) - \beta\gamma(\delta + a))(x, y)^2,$$

and two similar.

224.] **Higher binary quantics of even order.** We now pass to consider briefly the reduction to canonical forms of  $2n$ -ics, where  $n$  exceeds 2.

It has been seen (§ 208) that a binary  $2n$ -ic whose catalecticant vanishes can be expressed as a sum of  $n$   $2n$ th powers.

Now let  $u$  be the general binary  $2n$ -ic, whose catalecticant therefore does not vanish, and let  $v$  be any particular binary  $2n$ -ic with coefficients definitely chosen, either as constants or as functions of the coefficients in  $u$ , whose catalecticant does not vanish. Let  $\lambda$  be a constant free to have any value.

Write down the catalecticant of  $u - \lambda v$  and equate it to zero. The result is an equation in  $\lambda$ . This equation has a root or roots, i.e. there is a value, or values, of  $\lambda$  for which  $u - \lambda v$  is a sum of  $n$   $2n$ th powers.

A right form to assume for the general binary  $2n$ -ic  $u$  is then a sum of  $n$   $2n$ th powers together with a free multiple of any particular  $2n$ -ic  $v$  whose catalecticant does not vanish.

The most natural form to assume for the sextic would appear to be

$$X^6 + Y^6 + Z^6 + \lambda X^2 Y^2 Z^2,$$

but the reduction to this form has not, as a matter of fact, been effected.

The octavic however, as we shall presently see, has been brought by Sylvester to the corresponding form.

$$X^8 + Y^8 + Z^8 + W^8 + \lambda X^2 Y^2 Z^2 W^2.$$



225.] **The binary sextic.** The usual canonical form for the sextic is

$$x'^6 + y'^6 + z'^6 + \lambda' x' y' z' (y' - z') (z' - x') (x' - y'),$$

or, putting  $X + Y, \omega(X + \omega Y), \omega^2(X + \omega^2 Y)$  for  $x', y', z'$ ,

$$(X + Y)^6 + (X + \omega Y)^6 + (X + \omega^2 Y)^6 + \lambda(X^6 - Y^6).$$

It is convenient to take unity for the modulus of transformation. If  $x = lX + mY, y = l'X + m'Y$  be the substitution which reduces the sextic to the above form, the substitution

$$(lm' - l'm)^{\frac{1}{3}} x = lX + mY, (lm' - l'm)^{\frac{1}{3}} y = l'X + m'Y,$$

whose modulus is unity, reduces the sextic to the form

$$\nu\{(X + Y)^6 + (X + \omega Y)^6 + (X + \omega^2 Y)^6\} + \mu(X^6 - Y^6).$$

Let us assume only, with apparently smaller particularization, that

$$(a, b, c, d, e, f, g)(x, y)^6 \\ \equiv p(X + Y)^6 + q(X + \omega Y)^6 + r(X + \omega^2 Y)^6 + \mu(X^6 - Y^6).$$

We proceed to show how to find  $X^3 + Y^3$  and  $\mu$ . Suppose that

$$a'x^3 + 3b'x^2y + 3c'xy^2 + d'y^3 \equiv X^3 + Y^3.$$

Then, by § 46, the modulus being unity, we have also the equivalence of operations

$$a' \frac{d^3}{dy^3} - 3b' \frac{d^3}{dx dy^2} + 3c' \frac{d^3}{dx^2 dy} - d' \frac{d^3}{dx^3} = \frac{d^3}{dY^3} - \frac{d^3}{dX^3}.$$

Operate with the left-hand side here upon

$$(a, b, c, d, e, f, g)(x, y)^6,$$

and with the right on its equivalent in terms of  $X$  and  $Y$ . Remembering that

$$\left(\frac{d}{dY} - \frac{d}{dX}\right)(X + Y) = 0, \quad \left(\frac{d}{dY} - \omega \frac{d}{dX}\right)(X + \omega Y) = 0,$$

$$\left(\frac{d}{dY} - \omega^2 \frac{d}{dX}\right)(X + \omega^2 Y) = 0,$$

we see that the result on the right-hand side is

$$-120\mu(X^3 + Y^3),$$

which is the same as

$$-120\mu(a'x^3 + 3b'x^2y + 3c'xy^2 + d'y^3).$$

This is exhibited as the equivalent of another cubic in  $x$  and  $y$ . Equating the coefficients in the two, we have at once

$$a'd - 3b'c + 3c'b - d'a = -a'\mu,$$

$$a'e - 3b'd + 3c'c - d'b = -b'\mu,$$

$$a'f - 3b'e + 3c'd - d'c = -c'\mu,$$

$$a'g - 3b'f + 3c'e - d'd = -d'\mu,$$

which are made consistent for finding the mutual ratios of  $a', b', c', d'$  by choosing  $\mu$  so as to satisfy

$$\begin{vmatrix} d + \mu, c & , & b & , & a \\ e & , & d - \frac{1}{3}\mu, c & , & b \\ f & , & e & , & d + \frac{1}{3}\mu, c \\ g & , & f & , & e & , & d - \mu \end{vmatrix} = 0,$$

i. e. by solving a quartic equation; so that there are four such values of  $\mu$ .

Substitute one of these values of  $\mu$ . The ratios  $a' : b' : c' : d'$  are at once determined by any three of the linear equations now made consistent. It is now a matter of the solution of a cubic equation to split up

$$a'x^3 + 3b'x^2y + 3c'xy^2 + d'y^3$$

into its three factors, which must, but for constant multipliers, be the  $X + Y, X + \omega Y, X + \omega^2 Y$  of the canonical form.

The coefficients of the canonizing quartic equation in  $\mu$  must be invariants, as their property is quite independent of any linear transformation. The equation is in fact

$$\begin{vmatrix} d, c, b, a \\ e, d, c, b \\ f, e, d, c \\ g, f, e, d \end{vmatrix} + \frac{1}{9}(ag - 6bf + 15ce - 10d^2)\mu^2 + \frac{1}{9}\mu^4 = 0,$$

whose coefficients are the catalecticant and the quadric invariant.

It will be seen that the four values of  $\mu$  go in pairs  $\pm\mu_1, \pm\mu_2$ .

226.] Another, and perhaps a more useful, canonical form to which the sextic can be brought by linear transformation is that of Stephanos and Brill

$$x^6 + 15cx^4y^2 + 20dx^3y^3 + 15ex^2y^4 + y^6.$$

We may prove the possibility of the reduction as follows.

Taking the form

$$(X + Y)^6 + (X + \omega Y)^6 + (X + \omega^2 Y)^6 + 3\lambda(X^6 - Y^6),$$

to which the sextic has been brought by solution of a quadratic (in  $\mu^2$ ) and a cubic, put

$$X = plx + qmy, \quad Y = lx + my,$$

where  $p - q$  must not be zero. The equations found upon making the coefficients of  $x^5y$  and  $xy^5$  zero are

$$(p^5 + 10p^2)q + 10p^3 + 1 + \lambda(p^5q - 1) = 0,$$

$$(q^5 + 10q^2)p + 10q^3 + 1 + \lambda(pq^5 - 1) = 0.$$

Of these the difference has  $p + q$  as well as  $p - q$  for a factor. The former, which is allowable, makes either equation

$$-p^6 + 1 - \lambda(p^6 + 1) = 0,$$

giving  $p = -q = \left(\frac{1-\lambda}{1+\lambda}\right)^{\frac{1}{6}}$ . Thus we have a substitution which reduces a sextic to the form

$$a'x^6 + 15c'x^4y^2 + 20d'x^3y^3 + 15e'x^2y^4 + g'y^6,$$

or, altering the notation by putting  $x$  and  $y$  for  $a'^{\frac{1}{6}}x$  and  $g'^{\frac{1}{6}}y$ , the form required.

227.] **The binary octavic.** The canonical form of the octavic  $(a_0, a_1, \dots, a_8)(x, y)^8$  is

$$X^8 + Y^8 + Z^8 + W^8 + \lambda X^2 Y^2 Z^2 W^2,$$

where  $X, Y, Z, W$  are linear in  $x$  and  $y$ . We have to see that  $\lambda$  and the product

$$\begin{aligned} XYZW &\equiv (l_1x + m_1y)(l_2x + m_2y)(l_3x + m_3y)(l_4x + m_4y) \\ &\equiv (a', b', c', d', e')(x, y)^4, \text{ say,} \end{aligned}$$

can be found.

The operator  $(a', b', c', d', e') \left( \frac{d}{dy}, -\frac{d}{dx} \right)^4$  annihilates  $X^8$ ,  $Y^8$ ,  $Z^8$ ,  $W^8$ , as is clear, since  $(l \frac{d}{dy} - m \frac{d}{dx})(lx + my) = 0$ . We can further see that the same operator produces from  $X^2 Y^2 Z^2 W^2$  a constant multiple of  $XYZW$ . To do so affords a good example of the use of the concomitants of a quartic.

We have to see, in fact, that

$$(a', b', c', d', e') \left( \frac{d}{dy}, -\frac{d}{dx} \right)^4 \{ (a', b', c', d', e') (x, y)^4 \}^2$$

is a constant multiple of  $(a', b', c', d', e') (x, y)^4$ .

The expression is a covariant of the quartic by § 47. Its degree in the coefficients of the quartic is 3, and its order in the variables is 4. Now, referring to the complete list, § 212, of the irreducible concomitants of the quartic, we see that these,  $u, I, J, H, G$ , are of degree-orders (1, 4), (2, 0), (3, 0), (2, 4), (3, 6), and that the only covariant of degree-order (3, 4) which can be formed by combining them is the product  $Iu$  of the first two, i.e. is a constant multiple of  $u$  or

$$(a', b', c', d', e') (x, y)^4.$$

The result of operating with  $(a', b', c', d', e') \left( \frac{d}{dy}, -\frac{d}{dx} \right)^4$  on the identity of the octavic and its supposed canonical form is then the production of an identity

$$(a', b', c', d', e') \left( \frac{d}{dy}, -\frac{d}{dx} \right)^4 \cdot (a_0, a_1, \dots, a_8) (x, y)^8 \\ \equiv \mu' (a', b', c', d', e') (x, y)^4$$

of two quartics,  $\mu'$  being a constant qua  $x, y$ , namely a numerical multiple of the product of  $\lambda$  and the invariant  $I$  of the quartic on the right. Let us write  $8 \cdot 7 \cdot 6 \cdot 5 \cdot \mu$  for  $\mu'$ . We obtain, by equating coefficients of  $x^4, x^3y, x^2y^2, xy^3, y^4$  on the two sides, the equations

$$\begin{aligned} a'a_4 - 4b'a_3 + 6c'a_2 - 4d'a_1 + e'a_0 &= a'\mu, \\ a'a_5 - 4b'a_4 + 6c'a_3 - 4d'a_2 + e'a_1 &= b'\mu, \\ a'a_6 - 4b'a_5 + 6c'a_4 - 4d'a_3 + e'a_2 &= c'\mu, \\ a'a_7 - 4b'a_6 + 6c'a_5 - 4d'a_4 + e'a_3 &= d'\mu, \\ a'a_8 - 4b'a_7 + 6c'a_6 - 4d'a_5 + e'a_4 &= e'\mu, \end{aligned}$$

which are made consistent for determining  $a' : b' : c' : d' : e'$  by taking for  $\mu$  one of the roots of the canonizing quintic equation

$$\begin{vmatrix} a_4 - \mu, & a_3 & , & a_2 & , & a_1 & , & a_0 \\ a_5 & , & a_4 + \frac{1}{4}\mu, & a_3 & , & a_2 & , & a_1 \\ a_6 & , & a_5 & , & a_4 - \frac{1}{6}\mu, & a_3 & , & a_2 \\ a_7 & , & a_6 & , & a_5 & , & a_4 + \frac{1}{4}\mu, & a_3 \\ a_8 & , & a_7 & , & a_6 & , & a_5 & , & a_4 - \mu \end{vmatrix} = 0,$$

a quintic all whose non-vanishing coefficients are invariants of the octavic.

228.] **General binary  $2n$ -ic.** The success of the method adopted with variations in §§ 220, 225, 227 for the canonizing of a  $2n$ -ic depends on the knowledge, for the cases  $n = 2, 3, 4$ , of an auxiliary  $n$ -ic covariant  $V$  of an  $n$ -ic

$$(a'_0, a'_1, \dots, a'_n)(x, y)^n,$$

which is such that the derived  $n$ -ic covariant

$$(a'_0, a'_1, \dots, a'_n) \left( \frac{d}{dy}, -\frac{d}{dx} \right)^n \{ (a'_0, a'_1, \dots, a'_n)(x, y)^n \cdot V \}$$

is of the form

$$k(a'_0, a'_1, \dots, a'_n)(x, y)^n,$$

where  $k$  is a function of  $a'_0, a'_1, \dots, a'_n$  only.

If in the case of any higher value of  $n$  such a covariant  $V$  of the  $n$ -ic can be found, then the method of the preceding article will establish that the general binary  $2n$ -ic has the canonical form

$$X_1^{2n} + X_2^{2n} + \dots + X_n^{2n} + \lambda X_1 X_2 \dots X_n \cdot V,$$

where  $V$  is this covariant of  $X_1 X_2 \dots X_n$ , and  $\mu$ , a determinate constant multiple of  $\lambda$ , is any one of the roots of the canonizing  $(n+1)$ -ic equation, all whose coefficients are invariants,

and which for odd values of  $n$  is an  $\frac{1}{2}(n+1)$ -ic in  $\mu^2$  only,

$$\begin{vmatrix} a_n - \mu, & a_{n-1} & , & a_{n-2} & , & \dots, & a_0 \\ a_{n+1} & , & a_n + \frac{1}{n}\mu, & a_{n-1} & , & \dots, & a_1 \\ a_{n+2} & , & a_{n+1} & , & a_n - \frac{1 \cdot 2}{n(n-1)}\mu, & \dots, & a_2 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ a_{2n} & , & a_{2n-1} & , & a_{2n-2} & , & \dots, & a_n - (-1)^n \mu \end{vmatrix} = 0.$$

229.] **The ternary cubic.** For the reduction to canonical forms of ternary and quaternary quantics works (e.g. Salmon's) on geometry of two and three dimensions should be consulted. The reduction of the ternary cubic is alone given here.

The canonical form, due to Hesse, of the general ternary cubic is

$$X^3 + Y^3 + Z^3 + 6mXYZ.$$

The number of free constants ( $= 3 \times 3 + 1 = 10$ ) is the same as that of coefficients in the general ternary cubic. We have then an indication of the likelihood of the correctness of the form.

We shall prove the correctness by showing that, if we take any cubic which has the form, and give to all its ten coefficients quite arbitrary infinitesimal increments, the altered cubic also has the form. This will establish what is desired, for by continued repetition of a process of giving the coefficients independent infinitesimal increments one cubic may be made eventually to become any other.

Take a cubic which can be thrown into the form

$$X^3 + Y^3 + Z^3 + 6mXYZ.$$

Give to  $X, Y, Z$  and  $m$  the infinitesimal increments

$$\begin{aligned} \xi &= \epsilon_1 X + \epsilon_2 Y + \epsilon_3 Z, & \eta &= \eta_1 X + \eta_2 Y + \eta_3 Z, \\ & & \zeta &= \iota_1 X + \iota_2 Y + \iota_3 Z, & \mu, \end{aligned}$$

where  $\epsilon_1, \dots, \eta_1, \dots, \iota_1, \dots, \mu$  are arbitrary infinitesimal constants, ten in number. The consequent increment of the cubic, obtained by differentiation, is

$$\begin{aligned} &3(X^2 + 2mYZ)(\epsilon_1 X + \epsilon_2 Y + \epsilon_3 Z) \\ &+ 3(Y^2 + 2mZX)(\eta_1 X + \eta_2 Y + \eta_3 Z) \\ &+ 3(Z^2 + 2mXY)(\iota_1 X + \iota_2 Y + \iota_3 Z) + 6\mu XYZ. \end{aligned}$$

Now, if we identify this with the most general cubic in  $X, Y, Z$  with infinitesimal coefficients, we obtain ten equations of the first degree in  $\epsilon_1, \dots, \eta_1, \dots, \iota_1, \dots, \mu$  which are at once seen to determine these ten constants uniquely. And if  $X, Y, Z$  denote  $\lambda x + \mu y + \nu z, \lambda' x + \mu' y + \nu' z, \lambda'' x + \mu'' y + \nu'' z$ , whose coefficients are known constants, the most general cubic in  $X, Y, Z$  with infinitesimal coefficients is the same as the most general cubic in  $x, y, z$  with infinitesimal coefficients. We see then that, having a cubic which can be thrown into the form

$$X^3 + Y^3 + Z^3 + 6mXYZ,$$

the most general cubic obtained by giving its coefficients infinitesimal increments is also of the same form

$$(X + \xi)^3 + (Y + \eta)^3 + (Z + \zeta)^3 + 6(m + \mu)(X + \xi)(Y + \eta)(Z + \zeta).$$

Starting then from the cubic

$$x^3 + y^3 + z^3 + 6m'xyz,$$

which certainly has the form, we may pass by infinitesimal stages to any other cubic, and see that it must be expressible in the same form in the way indicated at the outset.

Ex. 33. Apply this method to show (§ 199) that  $X^3 + Y^3$  is a canonical form of the binary cubic.

Ex. 34. Also apply it to show (§ 211) that  $X^4 + Y^4 + 6mX^2Y^2$  is a canonical form of the binary quartic.

230.] **Catalecticant of ternary quartic.** We conclude this chapter with a theorem due to Sylvester as to the impossibility in general of a reduction which a mere counting of the constants might lead us hastily to assume possible. This will illustrate the necessity of a care which in much that has preceded may have seemed superfluous.

The general ternary *quartic*

$$(a, b, c, \dots)(x, y, z)^4$$

contains fifteen ( $= 1 + 2 + 3 + 4 + 5$ ) coefficients; and the sum of five fourth powers

$$\sum_{r=1}^{r=5} (l_r x + m_r y + n_r z)^4$$

contains exactly the same number of constants, which may be chosen at will.

It would then be expected that a canonical form of the general ternary quartic would be a sum of five fourth powers. But this is not the case. The fifteen coefficients in the sum of powers are, as we shall see, connected by a relation, which must consequently also connect the coefficients in the ternary quartic for the reduction to be possible.

The fact is akin to that of §§ 208, 210, in which, however, there is nothing in the same way paradoxical, as in the sum of two binary fourth powers the number of constants is obviously one less than in the general binary quartic.

The function of the coefficients in a ternary quartic which must vanish that the quartic may be a sum of five fourth powers is an invariant, called in analogy with the catalecticant of a binary quartic its catalecticant, i. e. is the eliminant of its six second partial differential coefficients, which are linear functions of  $x^2$ ,  $y^2$ ,  $z^2$ ,  $yz$ ,  $zx$ ,  $xy$ . Let us use a triple suffix notation according to which  $a_{rst}$ , where  $r+s+t=4$ , is the coefficient in the ternary quartic of the term  $k_{rst}x^r y^s z^t$  which occurs in the expansion of  $(x+y+z)^4$  by the multinomial theorem. The catalecticant is

$$\begin{vmatrix} a_{400} & a_{310} & a_{301} & a_{220} & a_{211} & a_{202} \\ a_{310} & a_{220} & a_{211} & a_{130} & a_{121} & a_{112} \\ a_{301} & a_{211} & a_{202} & a_{121} & a_{112} & a_{103} \\ a_{220} & a_{130} & a_{121} & a_{040} & a_{031} & a_{022} \\ a_{211} & a_{121} & a_{112} & a_{031} & a_{022} & a_{013} \\ a_{202} & a_{112} & a_{103} & a_{022} & a_{013} & a_{004} \end{vmatrix}.$$

The same function of the coefficients in  $(lx+my+nz)^4$  is

$$\begin{vmatrix} l^4 & l^3m & l^3n & l^2m^2 & l^2mn & l^2n^2 \\ l^3m & l^2m^2 & l^2mn & lm^3 & lm^2n & lmn^2 \\ l^3n & l^2mn & l^2n^2 & lm^2n & lmn^2 & ln^3 \\ l^2m^2 & lm^3 & lm^2n & m^4 & m^3n & m^2n^2 \\ l^2mn & lm^2n & lmn^2 & m^3n & m^2n^2 & mn^3 \\ l^2n^2 & lmn^2 & ln^3 & m^2n^2 & mn^3 & n^4 \end{vmatrix},$$

in which it will be noticed that the columns are respectively



$l^2, lm, ln, m^2, mn, n^2$  times the one column  $l^2, lm, ln, m^2, mn, n^2$ . The columns are then identical, but for different multipliers applied to them severally.

For the sum of five fourth powers the catalecticant is a determinant, obtained from that last written by writing for each constituent in it a sum of five like ones obtained by giving to  $l, m, n$ , or such of them as occur in it, the suffixes 1, 2, 3, 4, 5 in succession.

Now the determinant thus obtained is a sum of  $5^6$  determinants like the last written, except that the constituents have suffixes, which in any one of the determinants are the same in any column, but not, except in the case of five of the determinants, the same in all columns. All possibilities of applying the suffixes 1, 2, 3, 4, 5 to columns, one suffix to each, in fact occur in different determinants of the whole set of  $5^6$ .

But there are six columns and only five suffixes. In every one of the  $5^6$  determinants there must therefore be at least two columns with the same suffix. By the above, then, every one of the determinants contains two columns which, upon removal of factors such as  $l^2, lm, ln, m^2, mn, n^2$  for some suffix or other, are identical. Every one then vanishes. Consequently their sum, the catalecticant of the sum of five fourth powers, vanishes.

231.] The student will easily convince himself in like manner of the following facts.

The quaternary quartic  $(a, b, \dots)(x, y, t, u)^4$  contains thirty-five coefficients, and the sum of nine fourth powers of linear forms contains thirty-six constants, apparently one more than is necessary. Yet a quaternary quartic cannot be expressed as a sum of nine fourth powers unless its catalecticant, i.e. the eliminant of its second partial derivatives, vanishes. For ten, the number of these derivatives, exceeds nine, the number of squares.

The quinary quartic  $(a, b, \dots)(x, y, z, u, v)^4$  contains seventy coefficients, and the sum of fourteen fourth powers of linear functions of  $x, y, z, u, v$  contains seventy free constants. Yet a quinary quartic cannot be written as a sum of fourteen fourth powers unless its catalecticant vanishes, since  $15 > 14$ .

## CHAPTER XIII.

### INVARIANTS AND COVARIANTS OF THE BINARY QUINTIC AND SEXTIC.

232.] THE study of the binary quintic and its concomitants has been carried to a high degree of completeness by investigators, among whom Hermite, Cayley, Sylvester, Salmon, Clebsch, Gordan, and Faa de Bruno should be named. The present chapter contents itself with calling attention to the main facts, and some of the simpler applications thereof. It is beyond the scope of an introductory treatise to give a full synopsis of the mass of results at which the theory has arrived, or to endeavour to reproduce in outline more than the most elementary of the investigations which have produced those results.

The three absolutely independent invariants of lowest degrees have been encountered in previous chapters, and are of degrees 4, 8, 12 respectively. Any other invariant is a function of these by § 30: but there is a fourth of degree 18 (§ 114, Ex. 22), discovered by Hermite, which is irreducible in that it is not a rational integral function of them, but is connected with them by a syzygy which will be exhibited later. A method by which the existence of the syzygy is proved has been noticed in § 143.

The whole number of irreducible covariants and invariants of the quintic, the quintic itself being counted as one, is twenty-three, a number which the arithmetical method by analysis of a generating function, whose beginnings have been sketched in chapter viii, has been successful in indicating. The honour, not only of pointing out the number, but of exhibiting symbolically the concomitants themselves, is Gordan's. Their explicit forms have been investigated in

Cayley's second, third, fifth, eighth and ninth memoirs on quantics.

233.] **Canonical and semi-canonical forms.** For the detailed study of the quintic much use has been made of the form to which it may be reduced

$$aX^5 + bY^5 + c'(X + Y)^5,$$

or, say,

$$aX^5 + bY^5 + cZ^5,$$

where

$$X + Y + Z \equiv 0.$$

One of the three coefficients  $a, b, c$  may, in accordance with § 205, be taken as unity; or, if we allow  $a, b, c$  to be all arbitrary, we may suppose that the modulus of the substitution which reduces the general quintic to the form is unity. Chapter xviii of Salmon's *Higher Algebra* gives the forms, symmetrical of course in  $a, b, c$  and in  $X, Y, Z$ , of the concomitants for this symmetrical shape of canonical form.

A very convenient canonical form is Hammond's

$$x^5 + 5b'x^4y' + 5e'x'y'^4 + y'^5,$$

i.e.

$$(1, b', 0, 0, e', 1) (x', y')^5,$$

in which the two end coefficients are units and the two middle ones zero. He uses more the form

$$(a, b, 0, 0, e, f) (x, y)^5,$$

which contains too many free coefficients to be properly called canonical, but to which a substitution of unit modulus reduces any quintic, and which has the advantage of not excluding some special classes of quintics, whose coefficients obey invariant conditions, to which the more restricted canonical form does not apply.

We must see that the reduction to this form is possible.

Take the quintic in the form

$$aX^5 + bY^5 + c(X + Y)^5,$$

from which the most general linear substitution produces

$$a(lx + my)^5 + b(l'x + m'y)^5 + c\{(l + l')x + (m + m')y\}^5.$$

In this the coefficients of  $10x^3y^2$  and  $10x^2y^3$  are

$$al^3m^2 + bl'^3m'^2 + c(l + l')^3(m + m')^2$$

and

$$al^2m^3 + bl'^2m'^3 + c(l + l')^2(m + m')^3,$$

which both vanish if

$$al^2m^2 = bl'^2m'^2 = -c(l+l')^2(m+m')^2,$$

i. e. if  $\pm a^{\frac{1}{2}}lm = \pm b^{\frac{1}{2}}l'm' = c^{\frac{1}{2}}\sqrt{-1}(l+l')(m+m')$ ,

which suffice to determine  $\frac{l}{l'}$  and  $\frac{m}{m'}$ , whence  $lm' - l'm = 1$  derives  $lm$  and  $l'm'$ , leaving still one of  $l, m, l', m'$  which may be assigned arbitrarily.

In case (cf. § 204) the assumed reduction to the form

$$aX^5 + bY^5 + c(X + Y)^5$$

is impossible, there is as a rule no exception to the reducibility to Hammond's form. In fact it has been seen in § 204 that in the ordinary special case, when  $\beta = \gamma$  only, the special form

$$ax^5 + 5exy^4 + fy^5$$

is assumed, which is the case of Hammond's when  $b$  as well as  $c$  and  $d$  is zero. This is the case when  $I_{12}$ , the irreducible invariant of degree 12, vanishes.

In the more special case, when  $a = \beta = \gamma$ , we saw in the article referred to that the form

$$x^3(a'x^2 + 5b'xy + 10c'y^2)$$

is taken, and this, if we take for a new  $y$  one of the factors of  $a'x^2 + 5b'xy + 10c'y^2$ , becomes

$$x^3(5bxy + 10cy^2).$$

This is the one case of exception to the general applicability of Hammond's form.

It might be thought from mere counting of the constants that it would be possible in general to make the coefficient of  $x^4y$  as well as those of  $x^3y^2$  and  $x^2y^3$  vanish. It will be seen later, however, that this can only be done when an invariant condition  $I_{12} = 0$  is satisfied. That it can be done when  $I_{12} = 0$  has been seen above.

234.] **List of concomitants of binary quintic.** The twenty-three concomitants of a quintic, arranged in the order of Cayley's ninth memoir on quantics, are as follows. Many of them have been already met with. It will be seen that all are invariants or covariants from their methods of

formation. That they are irreducible, and form the complete irreducible system, it is beyond our scope to establish here. The best proof reposes on the method of transvectants (cf. § 61). We use the notation  $C_{i, \varpi}$  to denote a covariant of degree  $i$  in the coefficients and order  $\varpi$  in the variables, and  $I_i$  to denote an invariant of degree  $i$ . The arrangement is according to degree, and for the same degree according to order.

(1)  $u$  or  $C_{1, 5}$  is the quintic  $(a, b, c, d, e, f) (x, y)^5$  itself.

(2)  $C_{2, 2}$  is the quadratic covariant whose leading coefficient is the seminvariant  $ae - 4bd + 3c^2$ . It is the fourth transvectant of  $u$  and itself, or the quadratic invariant  $a'e' - 4b'd' + 3c'^2$  of the fourth emanant of  $u$ .

(3)  $C_{2, 6}$  is the Hessian of  $u$ . Its leading coefficient is  $ac - b^2$ .

(4)  $C_{3, 3}$  has for its leading coefficient

$$ace + 2bcd - ad^2 - b^2e - c^3.$$

It is obtained as this invariant of the fourth emanant of  $u$ , or as the result of putting  $\frac{d}{dy}$ ,  $-\frac{d}{dx}$  for  $x$  and  $y$  in (2) and operating on  $u$ . Let us express this shortly by saying that it is the result of operating with (2) on  $u$ . It is the canonizant (§ 207) of the quintic.

(5)  $C_{3, 5}$  is the Jacobian of  $u$  and (2). Its leading coefficient is  $a^2f - 5abe + 2acd + 8b^2d - 6bc^2$ .

(6)  $C_{3, 9}$  is the covariant whose leading coefficient is

$$a^2d - 3abc + 2b^3.$$

It is the Jacobian of  $u$  and its Hessian (3).

(7)  $I_4$  is the invariant of lowest degree. It is the discriminant of (2), viz.

$$(ae - 4bd + 3c^2)(bf - 4ce + 3d^2) - (af - 3be + 2cd)^2.$$

(8)  $C_{4, 4}$  is formed by adding nine times the square of (2) to the result of operating with (2) on (3), in the manner described in connexion with (4), and dividing by fifteen. Its leading coefficient is

$$a^2(e^2 - df) + a(3bcf - 3bde - 4c^2e + 4cd^2) \\ + 5b^2ce + 2b^2d^2 - 2b^3f - 8bc^2d + 3c^4.$$

It will be noticed that we have given it a different sign from that of our general convention in § 71.

(9)  $C_{4,6}$ , a second sextic covariant, is the Jacobian of  $u$  and (4). Its leading coefficient is

$$a^2(cf - de) - \alpha(b^2f + 2bce - 4bd^2 + c^2d) + 3b^3e - 6b^2cd + 3bc^3.$$

(10)  $C_{5,1}$ , the linear covariant of lowest degree in the coefficients, is the result of operating with (2) on (4), with sign changed.

(11)  $C_{5,3}$ , a second cubic covariant, is the Jacobian of (4) and (2).

(12)  $C_{5,7}$  is the Jacobian of (3) and (4).

(13)  $C_{6,2}$ , a second quadratic covariant, is given by operation with (2) on (8).

(14)  $C_{6,4}$ , a second quartic covariant, is given by operation with (10) on  $u$ .

(15)  $C_{7,1}$ , a second linear covariant, is the Jacobian of (10) and (2).

(16)  $C_{7,5}$ , a third quintic covariant, reckoning  $u$  itself as one, is the Jacobian of (13) and  $u$ .—(N.B. The quintic covariant (16) of Salmon's *Higher Algebra*, § 232, or Faa de Bruno's *Formes Binaires*, No. 12, Table v, is the result of subtracting from this  $C_{7,5}$  the product  $C_{2,2} C_{5,3}$  of (2) and (11).)

(17)  $I_8$ , the second invariant, is found as the invariant  $ac' + a'e - 2bb'$  of (2) and (13).

(18)  $C_{8,2}$ , a third quadratic covariant, is found as the Jacobian of (4) and (10).

(19)  $C_{9,3}$ , a third cubic covariant, is the Jacobian of (13) and (4).—(N.B. The covariant of degree 9 and order 3 in Faa de Bruno's *Formes Binaires*, No. 15, Table v, is  $96C_{9,3} - 16C_{2,2} C_{7,1} + 7I_4 C_{5,3}$ .)

(20)  $C_{11,1}$ , a third linear covariant, is given by operation with (2) on (19).

(21)  $I_{12}$ , the third invariant, is the discriminant, but for a numerical factor, of (13) or of (4).

(22)  $C_{13,1}$ , the fourth linear covariant, is the result of operating with (19) on (8). (Faa de Bruno's, No. 17, Table v, is  $-6C_{13,1} - 2I_8 C_{5,1}$ .)

(23)  $I_{18}$ , the fourth irreducible invariant, is the eliminant of the two linear covariants (10) and (22). It is also the catalecticant of (14).

Thus, to sum up, the irreducible concomitants of a binary quintic are

4 invariants	$I_4, I_8, I_{12}, I_{18},$
4 linear covariants	$C_{5,1}, C_{7,1}, C_{11,1}, C_{13,1},$
3 quadratic „	$C_{2,2}, C_{6,2}, C_{8,2},$
3 cubic „	$C_{3,3}, C_{5,3}, C_{9,3},$
2 quartic „	$C_{4,4}, C_{6,4},$
3 quintic „	$u, C_{3,5}, C_{7,5},$
2 sextic „	$C_{2,6}, C_{4,6},$
1 septic „	$C_{5,7},$
1 nonic „	$C_{3,9}.$

235.] **Forms of the concomitants when  $c = 0, d = 0$ .** The kindness of Mr. Hammond has supplied me with the forms taken by the twenty-three concomitants when the quintic is given his form  $(a, b, 0, 0, e, f)(x, y)^5$ , in which the two middle terms are wanting, this form being, as we have seen, in effect general. None of the expressions are of great complexity. They are

$$(1) C_{1,5} \equiv u = ax^5 + 5bx^4y + 5exy^4 + fy^5,$$

$$(2) C_{2,2} = aex^2 + (af - 3be)xy + bfy^2,$$

$$(3) C_{2,6} = -(b^2x^6 + e^2y^6) + 3(aex^2 + bfy^2)x^2y^2 + (af + 7be)x^3y^3,$$

$$(4) C_{3,3} = -\{b^2ex^3 + b^2fx^2y + ae^2xy^2 + be^2y^3\},$$

$$(5) C_{3,5} = (af - 5be)(ax^5 - fy^5) + (5af - 9be)(bx^3 - ey^3)xy + 8(b^2fx - ae^2y)x^2y^2,$$

$$(6) C_{3,9} = 2(b^3x^9 - e^3y^9) + 2(a^2ex^7 - bf^2y^7)xy + (af + 11be)(ax^5 - fy^5)x^2y^2 + (7af + 29be)(bx^3 - ey^3)x^3y^3 + 16(b^2fx - ae^2y)x^4y^4,$$

$$(7) I_4 = a^2f^2 - 10abef + 9b^2e^2,$$

$$(8) C_{4,4} = (a^2e^2 - 2b^3f)x^4 + (b^2f^2 - 2ae^3)y^4 + 4be(aex^2 + bfy^2)xy + 18b^2e^2x^2y^2,$$

- (9)  $C_{4,6} = (3be - af)(b^2x^6 - e^2y^6) - 2(a^2e^2 + b^3f)x^5y$   
 $+ 2(b^2f^2 + ae^3)xy^5 - 10be(aex^2 - bfy^2)x^2y^2,$
- (10)  $C_{5,1} = (a^2e^3 - ab^2f^2 + 6b^3ef)x + (b^3f^2 - a^2e^2f + 6abe^3)y,$
- (11)  $C_{5,3} = be(9be - af)(bx^3 - ey^3) + (4a^2e^3 - ab^2f^2 - 3b^3ef)x^2y$   
 $- (4b^3f^2 - a^2e^2f - 3abe^3)xy^2,$
- (12)  $C_{5,7} = 2(b^4fx^7 - ae^4y^7) + 10b^2e^2(ax^5 - fy^5)xy$   
 $+ 3be(af + 9be)(bx^3 - ey^3)x^2y^2$   
 $+ (ab^2f^2 - 6a^2e^3 + 19b^3ef)x^4y^3$   
 $- (a^2e^2f - 6b^3f^2 + 19abe^3)x^3y^4,$
- (13)  $C_{6,2} = (3ae^3 - b^2f^2)b^2x^2 - (af - 9be)b^2e^2xy$   
 $+ (3b^3f - a^2e^2)e^2y^2,$
- (14)  $C_{6,4} = (a^3e^2f - 5a^2be^3 - 2ab^3f^2 + 6b^4ef)x^4$   
 $- (ab^2f^3 - 5b^3ef^2 - 2a^2e^3f + 6abe^4)y^4$   
 $+ 4(a^2e^2f - 6abe^3 - b^3f^2)bx^3y$   
 $- 4(ab^2f^2 - 6b^3ef - a^2e^3)exy^3,$
- (15)  $C_{7,1} = (3a^3e^3f - a^2b^2f^3 - 15a^2be^4 + 7ab^3ef^2 - 18b^4e^2f)x$   
 $- (3ab^3f^3 - a^3e^2f^2 - 15b^4ef^2 + 7a^2be^3f - 18ab^2e^4)y,$
- (16)  $C_{7,5} = (a^2e^2f - 3abe^3 - 2b^3f^2)b^2x^5$   
 $+ (2a^3e^2 - 3ab^3f - 27b^4e)e^2x^4y$   
 $+ 8(a^2e^2 - 3b^3f)be^2x^3y^2$   
 $- 8(b^2f^2 - 3ae^3)b^2ex^2y^3$   
 $- (2b^2f^3 - 3ae^3f - 27be^4)b^2xy^4$   
 $- (ab^2f^2 - 3b^3ef - 2a^2e^3)e^2y^5,$
- (17)  $I_8 = a^2b^2e^2f^2 - 2a^3e^5 - 2b^5f^3 + 27b^4e^4,$
- (18)  $C_{8,2} = (4a^2e^3f - ab^2f^3 - 18abe^4 + 3b^3ef^2)b^2x^2$   
 $+ 2(a^3e^5 - b^5f^3)xy$   
 $- (4ab^3f^2 - a^3e^2f - 18b^4ef + 3a^2be^3)e^2y^2,$
- (19)  $C_{9,3} = (2b^2f^3 - 9ae^3f + 27be^4)b^4x^3$   
 $+ 3(ab^2f^2 + 9b^3ef - 6a^2e^3)b^2e^2x^2y$   
 $- 3(a^2e^2f + 9abe^3 - 6b^3f^2)b^2e^2xy^2$   
 $- (2a^3e^2 - 9ab^3f + 27b^4e)e^4y^3,$
- (20)  $C_{11,1} = (5a^3e^5f - a^2b^2e^2f^3 - 27a^2be^6 - 9ab^3e^3f^2 + 2b^5f^4$   
 $+ 54b^4e^4f)b^2x - (5ab^5f^3 - a^3b^2e^2f^2 - 27b^6ef^2$   
 $- 9a^2b^3e^3f + 2a^4e^5 + 54ab^4e^4)e^2y,$
- (21)  $I_{12} = b^2e^2(a^2b^2e^2f^2 - 4a^3e^5 - 4b^5f^3 + 18ab^3e^3f - 27b^4e^4),$



$$(22) C_{13,1} = (a^5 e^7 + b^8 f^4 - 8a^3 b^3 e^5 f + 9ab^6 e^3 f^2 + 27a^2 b^4 e^6 - 54b^7 e^4 f) ex + (b^7 f^5 + a^4 e^8 - 8ab^5 e^3 f^3 + 9a^2 b^3 e^6 f + 27b^6 e^1 f^2 - 54ab^4 e^7) by,$$

$$(23) I_{18} = (a^3 e^5 - b^5 f^3) \{ (af - 5be) (a^3 e^5 + b^5 f^3) - 10a^2 b^3 e^3 f^2 + 90ab^4 e^4 f - 216b^5 e^5 \}.$$

236.] **Discriminant of quintic.** It will be noticed that the discriminant of the quintic does not occur among the irreducible invariants  $I_4, I_8, I_{12}, I_{18}$ . Its degree is  $2(5-1) = 8$ . It might have been taken instead of  $I_8$ , being, as will be seen, the difference of a multiple of  $I_8$  and  $I_4^2$ ; but, as  $I_8$  itself is the simpler of the two, we prefer to speak of that and not of the discriminant as the irreducible invariant.

For the quintic in its form  $(a, b, 0, 0, e, f)(x, y)^5$  the discriminant is easily formed by elimination, following Bézout's method, between the two first deriveds

$$ax^4 + 4bx^3y + ey^4, \quad bx^4 + 4exy^3 + fy^4,$$

and is found to be

$$a^4 f^4 - 20a^3 b e f^3 + 256(a^3 e^5 + b^5 f^3) - 10a^2 b^2 e^2 f^2 - 180ab^3 e^3 f - 3375b^4 e^4,$$

which may be written

$$(a^2 f^2 - 10abef + 9b^2 e^2)^2 - 128(a^2 b^2 e^2 f^2 - 2a^3 e^5 - 2b^5 f^3 + 27b^4 e^4),$$

so that, by § 235 (7) and (17), the expression for the discriminant in terms of  $I_4$  and  $I_8$  is

$$\Delta = I_4^2 - 128I_8.$$

237.] **Syzygy among the invariants.** The four invariants  $I_4, I_8, I_{12}, I_{18}$ , though irreducible, must, as we have often seen, be connected by a syzygy. This may be expected to give the square of  $I_{18}$  in terms of the others. It is here sought.

As the quintic can be brought to the form

$$(a, b, 0, 0, e, f)(x, y)^5$$

by a substitution of modulus unity, it can in general be further brought to the canonical form

$$(a', 1, 0, 0, 1, f')(x', y')^5$$

by a further linear substitution which replaces  $bx^4y$  and  $exy^4$  by  $x'^4y'$  and  $x'y'^4$ . Let the modulus of the resultant

substitution which brings the quintic from its general to this last form be  $M$ . Then, from the expressions in § 235,

$$\begin{aligned} M^{10} I_4 &= a'^2 f'^2 - 10 a' f' + 9 \\ &= (a' f' - 1)(a' f' - 9), \\ M^{20} I_8 &= a'^2 f'^2 - 2 a'^3 - 2 f'^3 + 27, \\ M^{30} I_{12} &= a'^2 f'^2 - 4 a'^3 - 4 f'^3 + 18 a' f' - 27, \\ M^{45} I_{18} &= (a'^3 - f'^3) \{ (a' f' - 5)(a'^3 + f'^3) \\ &\quad - 10 a'^2 f'^2 + 90 a' f' - 216 \}. \end{aligned}$$

It is possible to eliminate  $a'$ ,  $f'$  and  $M$  between these four equations and obtain the syzygy required.

As a guidance see what happens when  $a' = f'$  so that  $I_{18} = 0$ . Writing  $J_4$ ,  $J_8$ ,  $J_{12}$  for the values taken by  $I_4$ ,  $I_8$ ,  $I_{12}$ , we have

$$\begin{aligned} M^{10} J_4 &= (a'^2 - 1)(a'^2 - 9) \\ M^{20} J_8 &= a'^4 - 4 a'^3 + 27 = (a' - 3)^2 (a'^2 + 2 a' + 3) \\ M^{30} J_{12} &= a'^4 - 8 a'^3 + 18 a'^2 - 27 = (a' - 3)^3 (a' + 1), \end{aligned}$$

or, writing  $\mu$  for  $\frac{M^{10}}{a' - 3}$ ,

$$\begin{aligned} \mu J_4 &= (a'^2 - 1)(a' + 3) \\ \mu^2 J_8 &= a'^2 + 2 a' + 3, \\ \mu^3 J_{12} &= a' + 1. \end{aligned}$$

These give, by substitution for  $a'$  from the last in the others,

$$\begin{aligned} \mu J_4 &= \mu^3 J_{12} (\mu^6 J_{12}^2 - 4), \\ \mu^2 J_8 &= \mu^6 J_{12}^2 + 2, \end{aligned}$$

by combination of which

$$J_4 = \mu^2 J_{12} (\mu^2 J_8 - 6).$$

We thus have a simple quadratic and cubic from which to eliminate  $\mu^2$ . The result is

$$\begin{aligned} J_4 J_8^4 + 8 J_8^3 J_{12} - 2 J_4^2 J_8^2 J_{12} - 72 J_4 J_8 J_{12}^2 \\ - 432 J_{12}^3 + J_4^3 J_{12}^2 = 0. \end{aligned}$$

It is suggested then to try whether the same function of  $I_4$ ,  $I_8$ ,  $I_{12}$  as this on the left is of  $J_4$ ,  $J_8$ ,  $J_{12}$ , a function whose

degree is 36, is of the form  $\lambda I_{18}^2$ , where  $\lambda$  is a constant. This proves to be the case with  $\lambda = 16$ . Thus

$$16 I_{18}^2 = I_4 I_8^4 + 8 I_8^3 I_{12} - 2 I_4^2 I_8^2 I_{12} - 72 I_4 I_8 I_{12}^2 - 432 I_{12}^3 + I_4^3 I_{12}^2$$

is the syzygy required.

A usual and elegant way of obtaining this syzygy is to show that, formed by the methods of § 234, the values of the invariants for the canonical form of unit modulus

$$lX^5 + mY^5 - n(X + Y)^5$$

$$\text{are } I_4 = (mn + nl + lm)^2 - 4lmn(l + m + n),$$

$$I_8 = l^2 m^2 n^2 (mn + nl + lm),$$

$$I_{12} = l^4 m^4 n^4,$$

$$I_{18} = l^5 m^5 n^5 (m - n)(n - l)(l - m),$$

so that  $l, m, n$  are the roots of the cubic

$$l^3 + \frac{I_4 I_{12} - I_8^2}{4 I_{12}^{\frac{5}{4}}} l^2 + \frac{I_8}{I_{12}^{\frac{1}{2}}} l - I_{12}^{\frac{1}{4}} = 0,$$

and  $I_{18}^2 I_{12}^{-\frac{5}{2}}$  is the product of the squares of differences between roots of this cubic.

238.] The quintic in a form with invariant coefficients. Hermite's Formes-types. It is an interesting proposition that if a quintic be so transformed that its variables are any two of its linear covariants, the coefficients are all invariants; and the same is true for any binary quantic whatever which has two linear covariants.

$$\text{Let } X = Px + Qy, \quad Y = P'x + Q'y$$

be any two of the linear covariants  $C_{5,1}, C_{7,1}, C_{11,1}, C_{13,1}$  of the quintic  $(a, b, c, d, e, f)(x, y)^5$ . We have

$$x = \frac{Q'X - QY}{PQ' - P'Q}, \quad y = \frac{-P'X + PY}{PQ' - P'Q},$$

in which the denominator is the modulus of the  $(X, Y)$  to  $(x, y)$  substitution, and is also an invariant, being the eliminant of two covariants.

We have now to show that in

$$(a, b, c, d, e, f)(Q'X - QY, -P'X + PY)^5$$

all the coefficients are invariants. This will be proved if we can show that they are annihilated by  $\Omega$  and by  $O$ , of which the first is

$$a\delta_b + 2b\delta_c + 3c\delta_d + 4d\delta_e + 5e\delta_f.$$

Now, if  $u, v$  stand for  $Q'X - QY, -P'X + PY$ ,

$$\begin{aligned} \Omega(a, b, c, d, e, f)(u, v)^5 &= 5v(a, b, c, d, e)(u, v)^4 \\ &+ 5(a, b, c, d, e)(u, v)^4 \Omega u + 5(b, c, d, e, f)(u, v)^4 \Omega v. \end{aligned}$$

We may here mean that the operation is not on  $X$  and  $Y$ , but only on coefficients of powers and products of powers of  $X$  and  $Y$  when the quintic is expressed in terms of  $a, b, c, d, e, f; P, Q, P', Q'; X, Y$ . Now, since (§ 109)  $\Omega P = 0, \Omega Q = P, \Omega P' = 0, \Omega Q' = P'$ , the operation being in this sense,

$$\Omega u = \Omega(Q'X - QY) = P'X - PY = -v,$$

and  $\Omega v = \Omega(-P'X + PY) = 0$ .

Consequently

$$\Omega(a, b, c, d, e, f)(u, v)^5 = 0,$$

i. e.  $\Omega(a, b, c, d, e, f)(Q'X - QY, -P'X + PY)^5 = 0$ ,

the operation not being on  $X, Y$ , but only on coefficients of  $X^5, X^4Y, \dots, Y^5$ .

All these coefficients are then annihilated by  $\Omega$ . Similarly all are annihilated by  $O$ . Accordingly all are invariants.

239.] **Quintics for which  $I_{12} = 0$ .** In § 233 it was stated that a quintic can only be linearly transformed to the form  $(a, 0, 0, 0, e, f)(x, y)^5$ , wanting its second as well as its third and fourth terms, when an invariant condition is satisfied. And it was seen that the said reduction can be effected when  $I_{12}$ , which is the discriminant of the canonizant, vanishes. To prove the necessity of this condition take Hammond's forms of the invariants (§ 235) of  $(a, b, 0, 0, e, f)(x, y)^5$ , and put  $b = 0$  in them. We get

$$I_4 = a^2 f^2,$$

$$I_8 = -2a^3 e^5,$$

$$I_{12} = 0,$$

$$I_{18} = a^7 e^{10} f,$$

of which the third proves the necessity stated.

From the values here of  $I_4, I_8, I_{12}$  it follows that for a quintic which can be reduced to the form now contemplated

$$16 I_{18}^2 = I_4 I_8^4,$$

and this is correctly what the syzygy of § 237 becomes when  $I_{12} = 0$ .

It is not hard to prove from the expressions for the invariants  $I_4, I_8, I_{12}$  of  $(a, b, 0, 0, e, f)(x, y)^5$ , which involve  $af, be$  and  $a^3e^5 + b^5f^3$  only, that  $be$ , which call  $\beta$ , is given by the equation

$$\left(I_4 - \frac{2I_8}{\beta^2} + \frac{I_{12}}{\beta^4}\right)^2 = 64 \left(2I_8 - \frac{I_{12}}{\beta^2}\right),$$

so that the product of all the values which  $be$  can have for reductions of the form  $(a, b, 0, 0, e, f)(x, y)^5$  is

$$I_{12}^2 (I_4^2 - 128 I_8)^{-1},$$

unless the discriminant  $I_4^2 - 128 I_8$  vanishes, when the product is still a multiple of  $I_{12}$ . We thus have quite clearly exhibited that when  $I_{12}$  vanishes some one at least of these values of  $be$  is zero, so that a reduction to the form  $(a, 0, 0, 0, e, f)(x, y)^5$  or the in fact equivalent form  $(a, b, 0, 0, 0, f)(x, y)^5$  is possible. The conclusion converse to that proved above, which was in effect arrived at before, as stated already, is thus confirmed.

A quintic for which  $I_{12} = 0$  cannot be expressed as a sum of three fifth powers, as was seen in § 204. In fact, the cano-  
nizant of  $ax^5 + 5exy^4 + y^5$ , to which form it can be reduced, is

$$C_{3,3} \equiv -ae^2xy^2.$$

Thus, if the reduction were possible, one of the  $X, Y, Z$  would be a multiple of  $x$  and the other two of  $y$ . Now

$$ax^5 + 5exy^4 + fy^5 \equiv lx^5 + my^5 + ny^5$$

is an impossibility unless  $e = 0$ , i.e. unless  $I_8 = 0$  and  $I_{18} = 0$  as well as  $I_{12} = 0$ .

We are thus guarded against an erroneous conclusion which might hastily be drawn from the last forms of the invariants in § 237. It might appear from those expressions that, whenever  $I_{12} = 0$ , either  $l = 0$  or  $m = 0$  or  $n = 0$ , and therefore  $I_8 = 0$  and  $I_{18} = 0$ . But this is not the case, the forms not being applicable to the case when  $I_{12} = 0$ .

240.] **Formes-types when  $I_{12} = 0$ .** Another interesting fact as to quintics for which  $I_{12} = 0$  may be derived from observing what the four linear covariants of the quintic become when  $b = 0$ ,  $c = 0$ ,  $d = 0$ .

Putting  $b = 0$  in (10) (15) (20) and (22) of § 235 we obtain

$$\begin{aligned} L_5 &\equiv C_{5,1} = \alpha^2 e^2 (ex - fy), \\ L_7 &\equiv C_{7,1} = \alpha^3 e^2 f (3ex + fy), \\ L_{11} &\equiv C_{11,1} = -2\alpha^4 e^7 y, \\ L_{13} &\equiv C_{13,1} = \alpha^5 e^8 x. \end{aligned}$$

Thus the  $y$  and  $x$  of the form  $(a, 0, 0, 0, e, f)(x, y)^5$ , to which a quintic for which  $I_{12} = 0$  may be reduced by a linear substitution of modulus unity, are multiples of the linear covariants of the 11th and 13th degrees, and are easily expressed also as sums of multiples of any two of the four linear covariants. We have, in fact, using the values of  $I_4$ ,  $I_8$ ,  $I_{18}$  in § 239,

$$\begin{aligned} \alpha x^5 &= a^{-24} e^{-40} L_{13}^5 = 256 I_8^{-8} L_{13}^5, \\ exy^4 &= \frac{1}{16} a^{-21} e^{-35} L_{13} L_{11}^4 = -8 I_8^{-7} L_{13} L_{11}^4, \\ fy^5 &= -\frac{1}{32} a^{-20} e^{-35} f L_{11}^5 = 16 I_8^{-9} I_{18} L_{11}^5. \end{aligned}$$

Thus the quintic reduced to the form  $(a, 0, 0, 0, e, f)(x, y)^5$ , with modulus unity, is

$$8 I_8^{-9} \{ 32 I_8 L_{13}^5 - 5 I_8^2 L_{13} L_{11}^4 + 2 I_{18} L_{11}^5 \}.$$

Consequently when  $I_{12} = 0$  one of the six ways (§ 238) of expressing the quintic with invariant coefficients expresses it in the reduced form.

Ex. 1. When  $I_{12} = 0$  prove that the other five expressions of the quintic with invariant coefficients are

- (2)  $I_8^{-18} (-8, 0, 0, 0, 4 I_8^9, 16 I_8^9 I_{18}) (4 I_{18} L_{11} + I_8^3 L_5, L_{11})^5$ ,
- (3)  $3^{-5} I_4^{-2} I_8^{-6} I_{18}^{-1} (2, 0, 0, 0, -81 I_4^2 I_8, 243 I_4^3 I_8)$   
 $(I_8 L_7 + I_4 L_{11}, L_{11})^5$ ,
- (4)  $2^{-6} I_{18}^{-4} (64 I_4^2, 0, 0, 0, -2 I_8, -I_8) (L_{13}, 2 L_{13} + I_8 L_5)^5$ ,
- (5)  $2^7 I_4^{-7} I_8^{-12} (2 I_4^7 I_8^4, 0, 0, 0, -I_4 I_8, 1)$   
 $(L_{13}, 3 I_4 I_8 L_{13} - 2 I_{18} L_7)^5$ ,
- (6)  $2^{-10} I_4^{-4} I_8^{-11} I_{18}^{-1} (2 I_4^2, 0, 0, 0, -I_8, -I_8)$   
 $(I_8^2 L_7 - 4 I_{18} L_5, I_8^2 L_7 + 12 I_{18} L_5)^5$ . (*Hammond*.)

241.] The classes of quintics for which respectively  $I_4 = 0$  and  $I_8 = 0$  will not long occupy us. We content ourselves with noticing that when  $I_4 = 0$  for the quintic

$$(a, b, 0, 0, e, f)(x, y)^5$$

the condition is, by § 235 (7),

$$(af - be)(af - 9be) = 0.$$

Thus when  $I_4 = 0$  a quintic may be reduced by a substitution of unit modulus with one degree of arbitrariness in its coefficients to one or other of the forms

$$ax^5 + 5bx^4y + 5exy^4 + \frac{be}{a}y^5,$$

$$ax^5 + 5bx^4y + 5exy^4 + \frac{9be}{a}y^5.$$

By a linear substitution of non-unit modulus it can then be given one of the forms

$$ax^5 + \frac{1}{a}y^5 + 5xy(x^3 + y^3),$$

$$ax^5 + \frac{9}{a}y^5 + 5xy(x^3 + y^3),$$

or, equally, one or other of the forms

$$x^5 + y^5 + 5xy\left(bx^3 + \frac{1}{b}y^3\right),$$

$$x^5 + y^5 + 5xy\left(bx^3 + \frac{1}{9b}y^3\right).$$

When  $I_8 = 0$ , provided  $I_{12}$  does not also vanish, by the last expressions in § 237, the quintic can by substitution of unit modulus be given the form

$$lX^5 + mY^5 - n(X + Y)^5,$$

where  $\frac{1}{l} + \frac{1}{m} + \frac{1}{n} = 0$ , i.e. the form

$$lX^5 + mY^5 + \frac{lm}{l+m}(X + Y)^5.$$

By a substitution of non-unit modulus it can be given the form

$$\frac{X^5}{\lambda} + \frac{Y^5}{1-\lambda} + (X + Y)^5.$$

When both  $I_8 = 0$  and  $I_{12} = 0$ , or both  $I_4 = 0$  and  $I_{12} = 0$ , we gather from § 237 or § 239 that also  $I_{18} = 0$ , and the cases are such as will occur below.

242.] Quintics for which  $I_{18} = 0$ . Quintics for which the skew invariant  $I_{18}$  vanishes have a special simplicity in that they are soluble.  $I_{18}$  is skew, for its weight is  $\frac{1}{2} 18 \cdot 5 = 45$ , i.e. is odd.

Consider the canonical form of unit modulus

$$lX^5 + mY^5 + nZ^5, \quad \text{where } X + Y + Z = 0.$$

Referring to § 237, we see that if  $I_{18} = 0$ , and if  $I_{12}$  does not vanish, so that the reduction to this form is possible,

$$(m - n)(n - l)(l - m) = 0,$$

which requires that two of  $l, m, n$  be equal. Thus a quintic for which  $I_{18} = 0$  and  $I_{12} \neq 0$  can be given by linear transformation of modulus unity the form

$$l(X^5 + Y^5) - m(X + Y)^5,$$

i.e. the form

$$(X + Y) \{ (l - m)(X^4 + Y^4) - (l + 4m)XY(X^2 + Y^2) + (l - 6m)X^2Y^2 \},$$

$$\text{i.e. } (X + Y) \{ (l - m)(X^2 + Y^2)^2 - (l + 4m)XY(X^2 + Y^2) - (l + 4m)X^2Y^2 \},$$

which can be broken up into  $X + Y$  and two quadratic, and then into five linear, factors.

Thus, so far, a quintic for which  $I_{18} = 0$  and  $I_{12} \neq 0$  can be solved.

Moreover the factors are of the forms

$$\begin{aligned} X + Y, \\ X^2 + Y^2 + pXY, \\ X^2 + Y^2 + qXY, \end{aligned}$$

and of these the single one which comes first is, speaking geometrically, one of the common harmonic conjugates  $X^2 - Y^2$  of the other two pairs.

Thus, if for a binary quintic  $I_{18} = 0$ , and  $I_{12} \neq 0$ , some one of its five linear factors is a double element of one of the three involutions determined by the other four factors taken in pairs.



Putting  $x$  for  $X + Y$  and  $y$  for  $X - Y$  the quintic above may be written in the form

$$kx(x^2 - \lambda y^2)(x^2 - \mu y^2),$$

i. e.  $ax^5 + 10cx^3y^2 + 5exy^4,$

a form in which all the coefficients of odd weight are wanting.

Conversely, if one of the five factors of a quintic be one of the common harmonic conjugates of two pairs which together constitute the other four factors, then  $I_{18} = 0$ .

For such a quintic can be given the form

$$ax(x^2 - \lambda y^2)(x^2 - \mu y^2),$$

i. e.  $ax^5 + 10cx^3y^2 + 5exy^4,$

in which no non-vanishing coefficient, and therefore no non-vanishing rational integral function of the coefficients, and consequently, in particular, no non-vanishing invariant, can be of odd weight.

Now all skew invariants (§ 95), and  $I_{18}$  in particular, are of odd weight. For such a quintic as contemplated then,  $I_{18}$  and all other skew invariants vanish.

Granting then, as we shall see in the next article, that the temporarily reserved case when  $I_{12} = 0$  as well as  $I_{18} = 0$  is only special and not exceptional, we have arrived at the fact that the condition  $I_{18} = 0$  is the necessary and sufficient one that the quintic may have the special property which has been expressed geometrically above.

We can also conclude that  $I_{18}$  is the only irreducible skew invariant which a quintic possesses. If  $I_{18} = 0$  the quintic has the above property. If it have that property all skew invariants vanish. Thus every skew invariant vanishes if  $I_{18}$  vanishes.  $I_{18}$  is then a factor of every skew invariant, and the invariant obtained by removing that factor is no longer skew. If its expression in terms of irreducible invariants involve another skew invariant, this may be analyzed in like manner; and so on.

Ex. 2. Solve the quintic equation

$$ax^5 + 5bx^4 + 5ex + \sqrt[3]{\frac{a^3e^5}{b^5}} = 0.$$

Ex. 3. By actual substitution, as in § 233, prove that

$$a(X^5 + Y^5) - c(X + Y)^5$$

can be transformed with modulus unity into

$$a'x^5 + 5b'x^4y + 6e'xy^4 + f'y^5,$$

where  $a'e^5 = b'^5f'^3$ .

*Ans.* As in § 233 a way to make  $c' = 0, d' = 0$  is given by taking  $a^{\frac{1}{2}}lm = a^{\frac{1}{2}}l'm' = c^{\frac{1}{2}}(l+l')(m+m')$ , whence, if

$$t = \frac{l}{m'} = \frac{l'}{m} = \frac{l+l'}{m+m'}, \text{ we get } \frac{a'}{f'} = t^5 \text{ and } \frac{b'}{e'} = t^3.$$

Ex. 4. Hence the form of Ex. 2 is a general one for quintics for which  $I_{18} = 0$ , but  $I_{12} \neq 0$ .

Ex. 5. Prove that

$$\begin{vmatrix} 1, & 2a, & a^2 \\ 1, & \beta + \gamma, & \beta\gamma \\ 1, & \delta + \epsilon, & \delta\epsilon \end{vmatrix} = 0$$

is the condition that  $x - ay$  be a common harmonic conjugate of the pairs

$$(x - \beta y)(x - \gamma y), (x - \delta y)(x - \epsilon y),$$

and that the determinant is a function of differences between pairs of  $a, \beta, \gamma, \delta, \epsilon$ .

*Ans.* It is annihilated by

$$\frac{d}{da} + \frac{d}{d\beta} + \frac{d}{d\gamma} + \frac{d}{d\delta} + \frac{d}{d\epsilon}.$$

It is  $(a - \delta)(a - \beta)(\epsilon - \gamma) + (a - \gamma)(a - \epsilon)(\delta - \beta)$ .

Ex. 6. One may be taken out of  $a, \beta, \gamma, \delta, \epsilon$  in five ways, and the rest go in pairs in three ways. Prove that  $a^{18}$  times the product of the  $5 \times 3 = 15$  determinants

$$\begin{vmatrix} 1, & 2a, & a^2 \\ 1, & \beta + \gamma, & \beta\gamma \\ 1, & \delta + \epsilon, & \delta\epsilon \end{vmatrix}$$

is an invariant of degree 18 and weight 45, and is consequently a numerical multiple of  $I_{18}$ . (*Cayley*.)

Ex. 7. Prove that the product of

$$\begin{aligned} & (a - \delta)(a - \beta)(\epsilon - \gamma) + (a - \epsilon)(a - \gamma)(\delta - \beta), \\ & (a - \beta)(a - \gamma)(\epsilon - \delta) + (a - \epsilon)(a - \delta)(\beta - \gamma), \\ & (a - \gamma)(a - \delta)(\epsilon - \beta) + (a - \epsilon)(a - \beta)(\gamma - \delta) \end{aligned}$$

is symmetrical in  $\beta, \gamma, \delta, \epsilon$ ; and hence also that  $a^{18}$  times the product of the five products of three such terms, with  $a, \beta, \gamma, \delta, \epsilon$  in succession taken for  $a$ , is a numerical multiple of an invariant  $I_{18}$ .  
(*Hermite.*)

243.] We temporarily reserved in § 242 the case of quintics for which  $I_{12} = 0$  as well as  $I_{18} = 0$ .

When  $I_{12} = 0$  the quintic can (§ 239) be given by substitution of unit modulus the form

$$ax^5 + 5exy^4 + fy^5,$$

and the condition  $I_{18} = 0$  is then

$$a^7 e^{10} f = 0,$$

so that either  $f = 0$  or  $e = 0$  or  $a = 0$ .

We also see from the syzygy  $16I_{18}^2 = I_4 I_8^4$ , which holds when  $I_{12} = 0$ , that the conditions  $I_{12} = 0, I_{18} = 0$  necessitate also that either  $I_4 = 0$  or  $I_8 = 0$ .

When  $f = 0$  the form taken is

$$ax^5 + 5exy^4,$$

i. e.  $X(X^4 + Y^4)$ .

This is the case when  $I_4 = 0$  as well as  $I_{12} = 0, I_{18} = 0$ . There is no exception here to the geometrical property in § 242.

When  $e = 0$  the form taken is

$$ax^5 + fy^5,$$

i. e.  $X^5 + Y^5$ ,

and again there is no exception. The case is that of quintics for which  $I_8 = 0$  as well as  $I_{12} = 0$  and  $I_{18} = 0$ .

When  $a = 0$  the form taken is

$$5exy^4 + fy^5,$$

i. e.  $Y^4(X + Y)$ ,

for which the property holds in a limiting form, for  $X + Y$  or any other linear form in  $X$  and  $Y$  is one of a pair of harmonic conjugates, the other being  $Y$ , of the coincident factors of  $Y^2$ .

This last class of cases is included in the class for which all the invariants vanish, but is not coextensive with that class. As we have seen in § 28, Ex. 3, all invariants vanish for a quintic

$$X^3(X + Y)(pX + qY)$$

with a perfect cube for factor, for such a quintic can be given the form

$$a_0 x^5 + 5 a_1 x^4 y + 10 a_2 x^3 y^2,$$

and no product of  $i$  factors chosen from among  $a_0, a_1, a_2$  can have a weight so great as to satisfy  $5i = 2w$ .

It will be remembered from § 233 that this case of a quintic with a cube factor is the one of irreducibility to Hammond's form. It is not one of exception to the geometrically expressed theorem as to quintics for which  $I_{18} = 0$ , for  $X$  is one of the common harmonic conjugates of the pairs

$$X^2, (X + Y)(pX + qY).$$

244.] **The binary sextic.** We will only give a list of what prove to be the complete system of irreducible concomitants of the sextic.

As indicated in § 143 the sextic has five irreducible invariants. Of these four  $I_2, I_4, I_6, I_{10}$  are absolutely independent. The fifth,  $I_{15}$ , is skew, and its square is given in terms of the rest by a syzygy of degree 30.

Clebsch and Gordan have found that the whole number of irreducible covariants and invariants, including the sextic itself, is 26, and the method of Cayley and Sylvester by means of generating functions, which has been referred to in chapter viii, confirms the result. The complete system has been exhibited as follows, the arrangement being, as in the case of the quintic, according to degrees in the coefficients, and for the same degree according to orders in the variables.

(1)  $u$ , or  $C_{1,6}$ , is the sextic  $(a, b, c, d, e, f, g)(x, y)^6$  itself.

(2)  $I_2$ , the invariant of degree 2, is  $ag - 6bf + 15ce - 10d^2$ . Cf. § 48.

(3)  $C_{2,4}$ , the first quartic covariant, is the covariant whose leading coefficient is the seminvariant

$$ae - 4bd + 3c^2.$$

It is the fourth transvectant of  $u$  and itself.

(4)  $C_{2,8}$ , an octavic covariant, is the Hessian, whose leading coefficient is  $ac - b^2$ .

(5)  $C_{3,2}$  is a quadratic covariant obtained by operation with  $u$ , having replaced in it  $x$  and  $y$  by  $\frac{d}{dy}$  and  $-\frac{d}{dx}$ , on the

Hessian  $C_{2,8}$ . The seminvariant which leads it is

$$(ac - b^2)g - 3(ad - bc)f + 2ae^2 - bde - 3c^2e + 2cd^2.$$

It can also be obtained by operation with  $C_{2,4}$  on  $u$ .

(6)  $C_{3,6}$  is a sextic covariant whose leader is

$$ace + 2bcd - ad^2 - b^2e - c^3$$

(§ 114, Ex. 13), the catalecticant of the fourth emanant.

(7)  $C_{3,12}$  is a duodecimic with the seminvariant

$$a^2d - 3abc + 2b^3$$

(§ 114, Ex. 15) for leader.

(8)  $C_{3,8}$ , a second octavic covariant, has for its leader

$$a^2f - 5abe + 2acd - 6bc^2 + 8b^2d$$

(§ 165).

(9)  $I_4$ , the irreducible invariant of degree 4, is the result of operating with  $u$  on  $C_{3,6}$ . It is the catalecticant (§ 208)

$$\begin{vmatrix} a, & b, & c, & d \\ b, & c, & d, & e \\ c, & d, & e, & f \\ d, & e, & f, & g \end{vmatrix}.$$

(10)  $C_{4,4}$  is a second quartic covariant, the Hessian but for a numerical factor of the first quartic covariant  $C_{2,4}$ . Its seminvariant leader is

$$2(ae - 4bd + 3c^2)(ag - 9ce + 8d^2) - 3(af - 3be + 2cd)^2,$$

where the coefficient 2 is, contrary to the usual convention of § 71, given to the alphabetically leading term  $a^2eg$  in order to avoid fractional coefficients.

(11)  $C_{4,6}$ , a third sextic covariant, is the Jacobian of  $u$  and  $C_{3,2}$ .

(12)  $C_{4,10}$ , a decimic, is the Jacobian of  $C_{2,8}$  and  $C_{2,4}$ .

(13)  $C_{5,2}$ , a second quadratic, is the result of operating on  $C_{2,4}$  with  $C_{3,2}$ .

(14)  $C_{5,4}$ , a third quartic, is the Jacobian of  $C_{2,4}$  and  $C_{3,2}$ .

(15)  $C_{5,8}$ , a third octavic, is the Jacobian of  $C_{2,8}$  and  $C_{3,2}$ .

(16)  $I_6$ , the irreducible invariant of degree 6, is the discriminant of  $C_{3,2}$ .

(17)  $C_{6,6}$ , a fourth sextic covariant, is the Jacobian of  $u$  and  $C_{5,2}$ .

(18)  $K_{6,6}$ , another covariant of the same degree and order 6, 6 as the last, is the Jacobian of  $C_{3,6}$  and  $C_{3,2}$ .

(19)  $C_{7,2}$ , a third quadratic, is the result of operating on  $C_{2,4}$  with  $C_{5,2}$ .

(20)  $C_{7,4}$ , a fourth quartic, is the Jacobian of  $C_{4,4}$  and  $C_{3,2}$ .

(21)  $C_{8,2}$ , a fourth quadratic, is the Jacobian  $C_{3,2}$  and  $C_{5,2}$ .

(22)  $C_{9,4}$ , a fifth quartic, is the Jacobian of  $C_{4,4}$  and  $C_{5,2}$ .

(23)  $I_{10}$ , the invariant of degree 10, is the discriminant of  $C_{5,2}$ .

(24)  $C_{10,2}$ , a fifth quadratic, is the Jacobian of  $C_{3,2}$  and  $C_{7,2}$ .

(25)  $C_{12,2}$ , a sixth quadratic, is the Jacobian of  $C_{5,2}$  and  $C_{7,2}$ .

(26)  $I_{15}$ , the skew invariant of degree 15 and weight  $\frac{1}{2} 6 \cdot 15 = 45$ , is the determinant (§ 17, Ex. 25) of the quadratics  $C_{3,2}$ ,  $C_{5,2}$ ,  $C_{7,2}$ . It is the criterion for those three quadratic covariants forming an involution.

245.] There are then altogether for the sextic the following irreducible concomitants:—

5 invariants  $I_2, I_4, I_6, I_{10}, I_{15}$ , of which the last is skew,

6 quadratic covariants  $C_{3,2}, C_{5,2}, C_{7,2}, C_{8,2}, C_{10,2}, C_{12,2}$ ,

5 quartic                    "      $C_{2,4}, C_{4,4}, C_{5,4}, C_{7,4}, C_{9,4}$ ,

5 sextic                     "      $u, C_{3,6}, C_{4,6}, C_{6,6}, K_{6,6}$ ,

3 octavic                   "      $C_{2,8}, C_{3,8}, C_{5,8}$ ,

1 decimic                   "      $C_{4,10}$ ,

1 duodecimic               "      $C_{3,12}$ .

None of the covariants are of odd order. Indeed, we have seen (§ 39) that no binary quantic of even order can have a covariant of odd order. In particular a sextic, or other binary quantic of even order, has no linear covariant.

We notice the occurrence of two irreducible covariants  $C_{6,6}$ ,  $K_{6,6}$  of the sixth order and the sixth degree, i.e. of two covariants of that order and degree which are linearly independent of one another, and of the covariants of the same order and degree which can be formed as products of lower covariants and invariants. This is the first instance of a state

of things which often occurs in connexion with quantics above the sixth order, but only in this one instance up to the sextic inclusively.

In forming covariants and invariants by operations which involve only differentiations with respect to variables, as for instance in the ordinary methods of finding them as hyper-determinants or transvectants, or, in particular, as Hessians, Jacobians, or results of operating with one covariant or quantic on another, we may, it is clear, with safety use canonical forms. Only operations of this class occur in the determinations of the more complicated of the above concomitants from the simpler ones. We may apply them to the canonical form of unit modulus

$$a(x^6 + y^6) + 15cx^4y^2 + 20dx^3y^3 + 15ex^2y^4,$$

as to which see § 226.

It should be mentioned, however, as of general applicability, that methods which use differentiations with regard to coefficients, such as that of evectants (§§ 67, 68), cannot as a rule be used in connexion with canonical forms. Such methods are not contemplated above.

246.] Complete systems of concomitants of the binary septic and octavic have been symbolically exhibited by Von Gall, and a good deal has been done with regard to quantics of a few orders higher. For no higher quantic, however, have explicit results been arrived at with completeness, except for that of infinite order, whose theory has been touched upon in chapter xi.

#### ADDITIONAL EXAMPLES.

Ex. 8. Prove that a quintic, deprived of its second term by writing  $x = X - bY$ ,  $y = aY$ , may be written

$$a(1, 0, C, D, a^2E - 3C^2, a^2F - 2CD)(X, Y)^5,$$

where

$$C = ac - b^2, \quad D = a^2d - 3abc + 2b^3, \quad E = ae - 4bd + 3c^2, \\ F = a^2f - 5abe + 8b^2d + 2acd - 6bc^2;$$

and that, if

$$J = ace + 2bcd - ad^2 - b^2e - c^3,$$

the relation  $D^2 = -a^3J + a^2CE - 4C^3$  can be used to reduce any expression to the first degree in  $D$ . (*Cayley*.)

Ex. 9. If  $\alpha, \beta, \gamma, \delta, \epsilon$  be the roots of a quintic, prove that

$$\alpha^2 \Sigma (\alpha - \beta)^2 (x - \gamma y)^2 (x - \delta y)^2 (x - \epsilon y)^2 = -100 C_{2,6}.$$

Ex. 10. If  $\alpha, \beta, \gamma, \delta, \epsilon, \zeta$  be the roots of a sextic, prove that

$$\alpha^2 \Sigma (\alpha - \beta)^2 (x - \gamma y)^2 (x - \delta y)^2 (x - \epsilon y)^2 (x - \zeta y)^2 = -180 C_{2,8}.$$

Ex. 11. For the quintic  $\alpha^4 \Sigma (\alpha - \beta)^2 (\beta - \gamma)^2 (\gamma - \alpha)^2 (\delta - \epsilon)^4$  is an invariant, and must be a numerical multiple of  $I_4$ .

Ex. 12. For the quintic  $\alpha^2 \Sigma (\alpha - \beta)^2 (\gamma - \delta)^2 (x - \epsilon y)^2$  is a covariant, and a numerical multiple of  $C_{2,2}$ .

Ex. 13. For the sextic  $\alpha^4 \Sigma (\alpha - \beta)^2 (\gamma - \delta)^2 (x - \epsilon y)^2 (x - \zeta y)^2$  is a numerical multiple of  $C_{2,4}$ .

Ex. 14. Prove that

$$\alpha^4 \Sigma (\alpha - \beta)^2 (\beta - \gamma)^2 (\gamma - \alpha)^2 (\delta - \epsilon)^2 (x - \delta y)^2 (x - \epsilon y)^2$$

is a covariant of a quintic which vanishes identically when the quintic has three equal roots. It must be a linear function of  $C_{4,4}$  and  $C_{2,2}^2$ . Show by considering the quintic  $x^5 + 10cx^3y^2$ , which has three equal roots, that it is a multiple of  $C_{2,2}^2 - 3C_{4,4}$ . (Cayley.)

Ex. 15. Prove that

$$\alpha^5 \Sigma (\alpha - \delta) (\alpha - \epsilon) (\beta - \delta) (\beta - \epsilon) (\gamma - \delta) (\gamma - \epsilon) (\delta - \epsilon)^2 \\ (x - \alpha y)^3 (x - \beta y)^3 (x - \gamma y)^3$$

is a covariant of degree 5 and order 9 of a quintic, which vanishes for a quintic having two pairs of equal roots. It must be a linear function of  $u C_{2,2}^2, u C_{4,4}$  and  $C_{2,6} C_{3,3}$ . Prove that it is a numerical multiple of  $50 C_{2,6} C_{3,3} - u (C_{4,4} + 3 C_{2,2}^2)$ . (Cayley.)

Ex. 16. Prove that

$$\alpha^4 \Sigma (\alpha - \beta) (\alpha - \gamma) (\alpha - \delta) (\alpha - \epsilon) (x - \beta y)^3 (x - \gamma y)^3 (x - \delta y)^3 (x - \epsilon y)^3$$

is a covariant of degree 4 and order 12, which vanishes for a quintic having three roots equal and the other two roots equal, and express it as  $k(3u^2 C_{2,2} - 25 C_{2,6}^2)$ . (Cayley.)

Ex. 17. For the sextic

$$I_2 = -\frac{1}{2} \frac{1}{10} \alpha^2 \Sigma (\alpha - \beta)^2 (\gamma - \delta)^2 (\epsilon - \zeta)^2. \quad (\text{Sylvester.})$$

Ex. 18. For the sextic

$$120(71 I_2 + 900 I_4) = \alpha^4 \Sigma (\alpha - \beta)^4 (\gamma - \delta)^4 (\epsilon - \zeta)^4. \quad (\text{Sylvester.})$$

Ex. 19. Show that

$$\left| \begin{array}{ccc} \alpha\beta, & \alpha + \beta, & 1 \\ \gamma\delta, & \gamma + \delta, & 1 \\ \epsilon\zeta, & \epsilon + \zeta, & 1 \end{array} \right|,$$

i. e.  $\alpha\beta(\gamma + \delta - \epsilon - \zeta) + \gamma\delta(\epsilon + \zeta - \alpha - \beta) + \epsilon\zeta(\alpha + \beta - \gamma - \delta),$



whose vanishing expresses that the quadratics

$$(x - \alpha y)(x - \beta y), (x - \gamma y)(x - \delta y), (x - \epsilon y)(x - \zeta y)$$

form an involution, is a function of the differences between pairs of  $\alpha, \beta, \gamma, \delta, \epsilon, \zeta$ .

Ex. 20. The six letters  $\alpha, \beta, \gamma, \delta, \epsilon, \zeta$  can be divided into pairs  $\alpha, \beta; \gamma, \delta; \epsilon, \zeta$ , in fifteen ways. Take each triad of pairs in a definite order, and write down the fifteen values of the function of differences in Ex. 19. Show that the product is symmetric in the roots, and must be a numerical multiple of  $a^{-15}I_{15}$ , where  $I_{15}$  is the skew invariant of the sextic of which  $\alpha, \beta, \gamma, \delta, \epsilon, \zeta$  are the roots.

(Joubert.)

Ex. 21. The vanishing of the skew invariant  $I_{15}$  of a sextic is the necessary and sufficient condition that the sextic be the product of three quadratics which form an involution, and consequently that, except in a very special case, it can be written as a cubic in  $X^2, Y^2$ , where  $X$  and  $Y$  are linear in the original variables.

Ex. 22. All skew invariants vanish for a sextic which can be thrown into the form  $aX^6 + 15cX^4Y^2 + 15eX^2Y^4 + gY^6$ .

Ex. 23. From the last two examples a sextic has no irreducible skew invariant but  $I_{15}$ .

## CHAPTER XIV.

### SEVERAL BINARY QUANTICS.

247.] It will be remembered (§§ 103, 115) that an invariant of several binary quantics

$$a_0 x^p + p a_1 x^{p-1} y + \frac{p(p-1)}{1 \cdot 2} a_2 x^{p-2} y^2 + \dots + a_p y^p,$$

$$a'_0 x^{p'} + p' a'_1 x^{p'-1} y + \frac{p'(p'-1)}{1 \cdot 2} a'_2 x^{p'-2} y^2 + \dots + a'_{p'} y^{p'},$$

&c., &c.,

has the two annihilators

$$\begin{aligned} \Sigma \Omega \equiv & \left( a_0 \frac{d}{da_1} + 2 a_1 \frac{d}{da_2} + \dots + p a_{p-1} \frac{d}{da_p} \right) \\ & + \left( a'_0 \frac{d}{da'_1} + 2 a'_1 \frac{d}{da'_2} + \dots + p' a'_{p'-1} \frac{d}{da'_{p'}} \right) + \dots, \end{aligned}$$

$$\begin{aligned} \Sigma O \equiv & \left( p a_1 \frac{d}{da_0} + \overline{p-1} a_2 \frac{d}{da_1} + \dots + a_p \frac{d}{da_{p-1}} \right) \\ & + \left( p' a'_1 \frac{d}{da'_0} + \overline{p'-1} a'_2 \frac{d}{da'_1} + \dots + a'_{p'} \frac{d}{da'_{p'-1}} \right) + \dots; \end{aligned}$$

that any covariant has the two annihilators

$$\Sigma \Omega - y \frac{d}{dx},$$

$$\Sigma O - x \frac{d}{dy};$$

and that any seminvariant, the leading coefficient in a covariant, has the one annihilator  $\Sigma \Omega$ .

It will also be remembered (§§ 103, 115) that for any invariant

$$ip + i'p' + \dots = 2w,$$

and that for any seminvariant which is not an invariant  $ip + i'p' + \dots$  exceeds  $2w$ , the excess

$$ip + i'p' + \dots - 2w$$

being the order in the variables of the covariant which the seminvariant leads. We here and throughout this chapter mean by seminvariant *rational integral* seminvariant.

We proceed to illustrate the theory of concomitants of several binary quantities by consideration of a few early cases.

248.] **Linear form and  $p$ -ic.** Let the linear form and the  $p$ -ic be

$$u \equiv \xi x + \eta y,$$

$$v \equiv (a_0, a_1, a_2, \dots a_p)(x, y)^p.$$

The linear form alone has no invariant, and no covariant distinct from itself.

The  $p$ -ic alone has a system of invariants and covariants which, in the preceding pages, have been investigated for values of  $p$  up to 4, and given for the values 5 and 6 of  $p$ .

The other *invariants* of the system are (§ 69) given by substituting  $\xi$  for  $y$  and  $-\eta$  for  $x$  in the covariants of the  $p$ -ic, including the  $p$ -ic itself. They are the eliminants of  $u$  and the covariants of  $v$ . They are also spoken of (§ 68) as the contravariants of  $v$  alone, if we regard  $u$  as the universal concomitant (§ 66) of two contragredient systems  $x, y; \xi, \eta$ . We shall not dwell on this aspect of them, but the notation is chosen so as to accord with it.

We seek information as to the other *covariants* of the system, or as to the *mixed concomitants* (§ 66) of  $v$ . The quest for these covariants is that for the *seminvariants* of the system which lead them.

These seminvariants are rational integral functions of  $\xi$  and  $\eta$  and  $a_0, a_1, \dots a_p$ , which are homogeneous, of different degrees  $i, i'$  it may be, both in  $\xi$  and  $\eta$  and in  $a_0, a_1, \dots a_p$ , and are isobaric in the two sets of coefficients taken together,  $\xi$  and  $\eta$  being of weights 0, 1 respectively. They have the one annihilator

$$\xi \frac{d}{d\eta} + \Omega.$$

Suppose that

$$\xi^i P_w + i \xi^{i-1} \eta P_{w-1} + \frac{i(i-1)}{1 \cdot 2} \xi^{i-2} \eta^2 P_{w-2} + \dots + \eta^i P_{w-i}$$



e. by the results of replacing  $x$  and  $y$  by  $-\eta$  and  $\xi$  in

$$\frac{1}{p!} \frac{d^p v}{dx^p}, \quad \frac{1!}{p!} \frac{d^{p-1} v}{dx^{p-1}}, \quad \frac{2!}{p!} \frac{d^{p-2} v}{dx^{p-2}}, \dots, \frac{1}{p} \frac{dv}{dx}, \quad v,$$

and after substitution dividing through by  $\xi^{w-i}$ .

Any seminvariant which is not a mere power of  $\xi$ , or, in particular, any invariant, is then a gradient in  $a_0, a_1, \dots, a_p$ , or such a gradient multiplied or divided by a power of  $\xi$ . Moreover, any gradient in them is a seminvariant or, in particular, invariant. For  $a_0, a_1, \dots, a_p$  are seminvariants themselves—the last of them, in fact, an invariant—being all annihilated by

$$\xi \frac{d}{d\eta} + \Omega.$$

249.] All seminvariants and invariants, including those of  $u$  and  $v$  singly, being thus rational integral functions of some or all of  $\xi$  and  $a_0, a_1, a_2, \dots, a_p$ , or such rational integral functions divided by powers of  $\xi$ , the search for the complete system of *irreducible* concomitants of  $u$  and  $v$  is the search for homogeneous isobaric functions of  $\xi$  and  $a_0, a_1, a_2, \dots, a_p$ , from which, when they are expressed in terms of  $\xi, \eta$  and  $a_0, a_1, a_2, \dots, a_p$ , powers of  $\xi$  may be removed by division, leaving the result integral. Such new forms have again to be combined with  $\xi, a_0, a_1, a_2, \dots, a_p$  and with one another, or with such of them as in the process are not excluded as themselves composite, and new forms derived by removal of  $\xi$  factors, till the process can be continued no longer. The method for thus arriving at all the irreducible concomitants is that illustrated in §§ 169, 170. Two early cases follow.

250.] **Case of two linear forms.** Let the two forms be

$$\xi x + \eta y, \quad ax + by.$$

The seminvariants  $\xi, a_0, a_1$  are

$$\xi, \quad a, \quad b\xi - a\eta;$$

and  $\xi$  is not a factor of any combination of them. These then are the only seminvariants and invariant; so that the complete system of concomitants of two linear forms consists of the two forms themselves and one invariant, their eliminant.

251.] Case of linear form and quadratic. For the system

$$\begin{aligned} u &\equiv \xi x + \eta y, \\ v &\equiv ax^2 + 2bxy + cy^2, \end{aligned}$$

the independent seminvariants  $\xi, a_0, a_1, a_2$  are

$$\xi, a, b\xi - a\eta, c\xi^2 - 2b\xi\eta + a\eta^2.$$

Here, if  $\xi$  were zero, we should have  $a_0 a_2 = a_1^2$ . Thus  $a_0 a_2 - a_1^2$  is divisible by  $\xi$ . In fact,

$$a_0 a_2 - a_1^2 = (ac - b^2) \xi^2.$$

Thus  $ac - b^2$  is a seminvariant (invariant) newly given. It is irreducible, for it is no function of the other one  $a$  which does not involve  $\xi$ .

Further examining the results of putting  $\xi = 0$  in the seminvariants now before us, i.e.

$$0, a, -\eta a, \eta^2 a, ac - b^2,$$

we see that no new relation connects them. We have before us then the complete system of irreducible concomitants, viz.

(1) the linear form itself, led by  $\xi$ ;

(2) the quadratic itself, led by  $a$ ;

(3) a covariant led by  $b\xi - a\eta$ . It is a linear covariant, the Jacobian

$$(b\xi - a\eta)x + (c\xi - b\eta)y.$$

Geometrically it is the harmonic conjugate of the linear form with regard to the quadratic;

(4) an invariant  $c\xi^2 - 2b\xi\eta + a\eta^2$ , the eliminant;

(5) an invariant  $ac - b^2$ , the discriminant of the quadratic.

252.] Another method, which might be pursued in examining the system of a linear form and  $p$ -ic, consists in using instead of

$$\xi, a_0, a_1, a_2, \dots, a_p$$

the system of  $p + 2$  protomorphs of § 168,

$$\begin{aligned} \xi, b\xi - a\eta, a_0, a_0 a_2 - a_1^2, a_0^2 a_3 - 3a_0 a_1 a_2 + 2a_1^3, \\ a_0 a_4 - 4a_1 a_3 + 3a_2^2, \dots \end{aligned}$$

A third method, which will be adopted, depends on the fact that if  $S$  be a seminvariant, or invariant, of  $\xi x + \eta y$  and  $v$ , which involves  $\xi$  and  $\eta$ , i.e. which is not either a power of  $\xi$  or a seminvariant of  $v$  alone, then  $\frac{dS}{d\eta}$  is another.

The fact is an immediate consequence of the identity of operators

$$\left(\xi \frac{d}{d\eta} + \Omega\right) \frac{d}{d\eta} = \frac{d}{d\eta} \left(\xi \frac{d}{d\eta} + \Omega\right),$$

which tells us that when  $\xi \frac{d}{d\eta} + \Omega$  annihilates  $S$ , so that

$$\frac{d}{d\eta} \left(\xi \frac{d}{d\eta} + \Omega\right) S = 0,$$

then  $\xi \frac{d}{d\eta} + \Omega$  also annihilates  $\frac{d}{d\eta} S$ .

Thus from an *invariant*

$$(Q_{w-i}, Q_{w-i+1}, \dots, Q_w)(\eta, -\xi)^i$$

of  $\xi x + \eta y$  and  $v$ , formed as in § 248 from  $v$  or a covariant of  $v$ , are derived the series of seminvariants

$$(Q_{w-i}, Q_{w-i+1}, \dots, Q_{w-1})(\eta, -\xi)^{i-1},$$

$$(Q_{w-i}, Q_{w-i+1}, \dots, Q_{w-2})(\eta, -\xi)^{i-2},$$

. . . . .

$$Q_{w-i}\eta - Q_{w-i+1}\xi,$$

$$Q_{w-i},$$

of which the last is the corresponding seminvariant of  $v$  only.

The way in which, in the following two cases, this is utilized for the determination of complete systems of  $\xi x + \eta y$  and  $v$ , when the complete system of  $v$  is known, is general. We shall see that  $\xi$ , the irreducible invariants of  $v$ , the invariants of  $\xi x + \eta y$  and  $v$  obtained by putting  $\eta, -\xi$  for  $x, y$  in the irreducible covariants of  $v$ , and the successive derivatives of these with regard to  $\eta$ , constitute together the complete system of seminvariants and invariants.

253.] **Case of linear form and cubic.** Take the linear form

$$\xi x + \eta y$$

and the cubic

$$ax^3 + 3bx^2y + 3cxy^2 + dy^3.$$

The complete system for the linear form alone is itself.

The complete system for the cubic alone consists (§ 169) of three covariants

$$(a, b, c, d)(x, y)^3, \text{ i. e. the cubic itself,}$$

$$(ac - b^2)x^2 + (ad - bc)xy + (bd - c^2)y^2,$$

$$(a^2d - 3abc + 2b^3, abd - 2ac^2 + b^2c, -acd + 2b^2d - bc^2, \\ -ad^2 + 3bcd - 2c^3)(x, y)^3,$$

and one invariant

$$(ad - bc)^2 - 4(ac - b^2)(bd - c^2).$$

Thus the system has

(1) the seminvariant  $\xi$ ;

(2) the invariant  $(a, b, c, d)(\eta, -\xi)^3$ ,

by § 248, from which flow, by § 252, the seminvariants

(3)  $(a, b, c)(\eta, -\xi)^2$ ,

(4)  $a\eta - b\xi$ ,

(5)  $a$ ;

(6) the invariant  $(ac - b^2)\eta^2 - (ad - bc)\eta\xi + (bd - c^2)\xi^2$ ,

from which flow the seminvariants

(7)  $2(ac - b^2)\eta - (ad - bc)\xi$ ,

(8)  $ac - b^2$ ;

(9) the invariant  $(a^2d - 3abc + 2b^3, -abd + 2ac^2 - b^2c, \\ -acd + 2b^2d - bc^2, ad^2 - 3bcd + 2c^3)(\eta, \xi)^3$ ,

from which flow the seminvariants

(10)  $(a^2d - 3abc + 2b^3, -abd + 2ac^2 - b^2c, \\ -acd + 2b^2d - bc^2)(\eta, \xi)^2$ ,

(11)  $(a^2d - 3abc + 2b^3)\eta - (abd - 2ac^2 + b^2c)\xi$ ,

(12)  $a^2d - 3abc + 2b^3$ ;

and lastly the invariant

(13)  $(ad - bc)^2 - 4(ac - b^2)(bd - c^2)$ .

None of these thirteen is a rational integral function of any of the others. That (5), (8), (12) and (13) are irreducible is the theory of the single cubic (§ 169). That (1) is irreducible is obvious. We have still to see that none of the rest of the



thirteen, i. e. none of those which involve  $\xi$  and  $\eta$ , is a rational integral function of others of the thirteen. Suppose, if possible, that

$$P\eta^m + Q\eta^{m-1}\xi + \dots + Z\xi^m,$$

where  $m$  is 1 or 2 or 3, and  $P, Q, \dots Z$  do not involve  $\xi$  and  $\eta$ , is the expansion of a rational integral function of some of the thirteen, not merely one of them, and proves to be the same as one of them. The coefficient  $P$  of  $\xi^m$  in a rational integral function of (1) to (13) must be either a rational integral function of

$$a, ac - b^2, a^2d - 3abc + 2b^3, (ad - bc)^2 - 4(ac - b^2)(bd - c^2)$$

or must vanish. Now no rational integral function of this complete system of irreducible seminvariants of a single cubic is equal to one of them except that one itself, this being the fact of their irreducibility; and  $P$  cannot vanish and yet be the first coefficient in one of (1) to (13), for none of these is divisible by  $\eta$ .

We have still to show that this system of thirteen irreducible seminvariants and invariants is complete.

254.] **The system is complete.** We have to see that any seminvariant or invariant whatever of the linear form and cubic can be rationally and integrally expressed in terms of the system (1) to (13) of the preceding article.

Firstly, any seminvariant or invariant in which only  $a, b, c, d$  occur is a rational integral function of (5), (8), (12), (13), by the theory of the single cubic.

Secondly, any seminvariant in which  $a, b, c, d$  do not occur is a mere power of  $\xi$ .

It remains to consider seminvariants and invariants in which both sets  $a, b, c, d$  and  $\xi, \eta$  are represented. Let

$$S \equiv P\eta^i + Q\eta^{i-1}\xi + R\eta^{i-2}\xi^2 + \dots + Z\xi^i$$

be one, from which factors which are powers of  $\xi$  have been removed.  $P, Q, \dots Z$  are rational integral functions of degree  $i'$  in  $a, b, c, d$  or some of them.

By § 248,  $P$  is a seminvariant in  $a, b, c, d$ . It can therefore be rationally and integrally expressed in terms of

$$a, ac - b^2, a^2d - 3abc + 2b^3, (ad - bc)^2 - 4(ac - b^2)(bd - c^2),$$

which call  $a, C, D, \Delta$ .

If  $w$  be the whole weight of  $S$ , we have (§ 247)

$$i + 3i' \leq 2w.$$

Now  $P$  consists of a sum of positive and negative numerical multiples of such products as  $\alpha^p C^q D^r \Delta^s$ , where

$$p + 2q + 3r + 4s = i'.$$

The term  $P\eta^i$  in  $S$  is then a sum of numerical multiples of such terms as

$$\alpha^p C^q D^r \Delta^s \cdot \eta^i,$$

for which, expressing that  $i + 3i' \leq 2w$ , we have

$$i + 3(p + 2q + 3r + 4s) \leq 2(i + 2q + 3r + 6s),$$

i. e.

$$3p + 2q + 3r \leq i$$

$$= i + t, \text{ say,}$$

where  $t$  is a positive integer or zero. It is important to see that this implies, among other things, that  $p, q, r$  cannot all vanish, since  $i$  does not.

Now in the product

$$(2)^p (6)^q (9)^r (13)^s$$

the highest term in  $\eta$  is

$$\alpha^p C^q D^r \Delta^s \cdot \eta^{3p+2q+3r},$$

and the product

$$(5)^p (8)^q (12)^r (13)^s$$

is

$$\alpha^p C^q D^r \Delta^s;$$

and, if in this last we replace one of the  $p$  factors (5) by (4), or one of the  $q$  factors (8) by half of (7), or one of the  $r$  factors (12) by (11), we produce a first term

$$\alpha^p C^q D^r \Delta^s \cdot \eta,$$

and again, by a like process of retrogression, we produce

$$\alpha^p C^q D^r \Delta^s \cdot \eta^2,$$

and so on. Continuing the process of unit retrogression we must arrive, before or upon reaching the final first term

$$\alpha^p C^q D^r \Delta^s \cdot \eta^{3p+2q+3r},$$

at the desired first term

$$\alpha^p C^q D^r \Delta^s \cdot \eta^{3p+2q+3r-t},$$

i. e.

$$\alpha^p C^q D^r \Delta^s \cdot \eta^i,$$

as a rule in a number of different ways.

Similarly for any other term of which  $P\eta^i$  consists.

Thus we can, and as a rule in a number of ways, obtain by composition of (2) to (13) a seminvariant

$$P\eta^i + Q'\eta^{i-1}\xi + R'\eta^{i-2}\xi^2 + \dots + Z'\xi^i,$$

whose  $\eta^i$  term is the same as that of  $S$ .

Subtracting it from  $S$ , and removing the seminvariant factor  $\xi$ , we obtain a seminvariant

$$(Q - Q')\eta^{i-1} + (R - R')\eta^{i-2}\xi + \dots + (Z - Z')\xi^{i-1}.$$

Let the same process be repeated. We can form a combination of (2) to (13) whose highest term in  $\eta$  is  $(Q - Q')\eta^{i-1}$ . Subtracting this, and dividing by  $\xi$ , we have a seminvariant

$$(R - R' - R'')\eta^{i-2} + \dots + (Z - Z' - Z'')\xi^{i-2}.$$

Repeat the process again; and continually as long as necessary. We get, lastly, unless at some stage or other the result of subtraction has vanished, in which case the desired expression of  $S$  is obtained, a residual

$$Z - Z' - Z'' - \dots - Z_i,$$

which is a seminvariant free from  $\xi$ ,  $\eta$ , and so a rational integral function of (5), (8), (12), (13).

Thus, finally, we have  $S$  expressed as

$$S_1 + \xi S_2 + \xi^2 S_3 + \dots + \xi^i S_i,$$

where  $S_1, S_2, S_3, \dots, S_i$  are rational integral functions of (2) to (13) or some of them.

We see then that every seminvariant or invariant of the linear form and cubic is a rational integral function of (1) to (13) or some of them. These were seen to be irreducible. The proof is now complete that they form the complete system of irreducible seminvariants and invariants.

The complete system of irreducible covariants and invariants follows at once from the complete system of irreducible seminvariants and invariants. The covariant corresponding to any one of the seminvariants  $S$  is, by § 115,

$$x^\omega e^{\frac{y}{x}(\eta \frac{d}{d\xi} + 0)} S,$$

where

$$\omega = i + 3i' - 2w.$$

255.] **Case of linear form and quartic.** The irreducible covariants and invariants of the quartic

$$(a, b, c, d, e)(x, y)^4,$$

are, § 170, the quartic itself, the covariants

$$(ac - b^2)x^4 + 2(ad - bc)x^3y + (ae + 2bd - 3c^2)x^2y^2 \\ + 2(be - cd)xy^3 + (ce - d^2)y^4,$$

and

$$x^6 e^{\frac{y}{x} 0} (a^2 d - 3abc + 2b^3),$$

and the invariants

$$I \equiv ae - 4bd + 3c^2,$$

$$J \equiv ace + 2bcd - ad^2 - b^2e - c^3.$$

The invariants of the system consisting of  $\xi x + \eta y$  and the quartic are then

$$(1) I,$$

$$(2) J,$$

$$(3) (a, b, c, d)(\eta, -\xi)^4,$$

$$(4) (ac - b^2)\eta^4 - 2(ad - bc)\eta^3\xi + (ae + 2bd - 3c^2)\eta^2\xi^2 \\ - 2(be - cd)\eta\xi^3 + (ce - d^2)\xi^4,$$

$$(5) \eta^6 e^{-\frac{\xi}{\eta} 0} (a^2 d - 3abc + 2b^3),$$

and the other seminvariants are, as in the last two articles,  $\xi$ , the four successive differential coefficients of (3), the four successive differential coefficients of (4), and the six successive differential coefficients of (5), all with respect to  $\eta$ .

Altogether we have twenty seminvariants and invariants.

The twenty are all irreducible. This is established exactly as in § 253.

Moreover, as in § 254, any seminvariant or invariant whatever can be rationally and integrally expressed in terms of the twenty or some of them. They form, then, the complete irreducible system of seminvariants and invariants.

From every seminvariant of the system the corresponding irreducible covariant is formed by the operation and multiplication

$$x^{i+4i'-2w} e^{\frac{y}{x} \left( \eta \frac{d}{d\xi} + 0 \right)}.$$

256.] **Case of  $n$  linear forms.** For the case of  $n$  linear forms

$$a_1x + b_1y, a_2x + b_2y, \dots, a_nx + b_ny,$$

an *algebraically* complete system of concomitants, i.e. a system of which all other covariants and invariants are functions, though not necessarily rational integral functions, consists of the  $n$  linear forms themselves and the  $n-1$  invariants

$$a_1b_2 - a_2b_1, a_1b_3 - a_3b_1, \dots, a_1b_n - a_nb_1,$$

which are the eliminants of a chosen one of the forms and the other  $n-1$ .

For

$$a_1, a_2, \dots, a_n, a_1b_2 - a_2b_1, a_1b_3 - a_3b_1, \dots, a_1b_n - a_nb_1$$

are all independent, each involving a letter which does not occur in any previous one, and their whole number  $2n-1$  is less by 4, the number of  $l, m, l', m'$ , than  $2n+2+1$ , i.e. than the number of equations which express the equalities of coefficients of  $X$  and  $Y$  in the given forms and their linear transformations together with the equations of substitution

$$x = lX + mY, \quad y = l'X + m'Y$$

and the one equation

$$M = lm' - l'm. \quad (\text{Cf. } \S 42.)$$

The complete *irreducible* system consists of the above and the other eliminants

$$a_2b_3 - a_3b_2, a_2b_4 - a_4b_2, \dots, a_3b_4 - a_4b_3, \dots, \dots, a_{n-1}b_n - a_nb_{n-1}$$

of pairs of the  $n$  forms. The whole number of the system is the sum of  $n$ , the number of linear forms, and  $\frac{1}{2}n(n-1)$ , the number of eliminants, i.e. is

$$\frac{1}{2}n(n+1).$$

We must see that they are all irreducible, and that there is no other covariant or invariant which is not a rational integral function of them.

They are all irreducible. For the leaders

$$a_1, a_2, \dots, a_n$$

of the forms themselves, being different and of the first degree,

are irreducible, and the eliminants  $a_1b_2 - a_2b_1, \dots$  are all of degree 2, so that any one of them if not irreducible would have to be a linear function of the rest and of the squares and products of  $a_1, a_2, \dots, a_n$ . These last cannot enter with the others in any linear relation, for they are all of weight zero, and the rest are of weight 1: and no linear relation can connect the eliminants alone, for they are of different partial degrees in the coefficients of the  $n$  forms separately.

To prove that every other seminvariant is reducible in terms of  $a_1, a_2, \dots, a_n$  and the eliminants, we may proceed by mathematical induction. Assume it true for the above  $n$  forms, and take an  $(n+1)$ th  $\xi x + \eta y$ . Exactly as in § 254, any seminvariant of the system of  $n+1$  forms is a rational integral function of

$$a_1b_2 - a_2b_1, a_1b_3 - a_3b_1, \dots, a_2b_3 - a_3b_2, \dots, \dots, a_{n-1}b_n - a_nb_{n-1}, \\ a_1\eta - b_1\xi, a_2\eta - b_2\xi, \dots, a_n\eta - b_n\xi,$$

and the  $\eta$ -derivatives of these

$$a_1, a_2, \dots, a_n,$$

together with

$$\xi.$$

This proves the theorem for  $n+1$  forms when we know it for  $n$ . But (§ 250) we know it for  $n=2$ . Consequently it is true for  $n=3, 4, 5, \dots$ , i.e. universally.

In proofs by the method of § 254 the critical fact is that, in the inequalities like the

$$3p + 2q + 3r \leq i$$

of that article, the non-vanishing coefficients on the left are all positive. This is a universal fact, for every coefficient is the  $\Sigma(ip) - 2w$  of a seminvariant, and this is never negative (§ 247).

257.] **System of two quadratics.** This is the only system of two quantities neither of which is linear which we shall discuss at length.

Let the two quadratics be

$$u \equiv ax^2 + 2bxy + cy^2,$$

$$v \equiv a'a^2 + 2b'xy + c'y^2.$$

Six seminvariants and invariants we have at once the means of writing down, viz.

- (1)  $a$ , the leading coefficient of  $u$ ,
- (2)  $a'$ , the leading coefficient of  $v$ ,
- (3)  $ac - b^2$ , the one invariant of  $u$  alone, its discriminant,
- (4)  $a'c' - b'^2$ , the one invariant of  $v$  alone, its discriminant,
- (5)  $ac' + a'c - 2bb'$ , the invariant of  $u$  and  $v$  intermediate to (3) and (4), found in § 18.
- (6)  $ab' - a'b$ , the leading coefficient of the Jacobian of  $u$  and  $v$ .

These six form the complete irreducible system. This may be seen as follows.

Firstly none of them is a rational integral function of the rest. This is clearly the case with regard to the two  $a, a'$  of the first degree. As to the rest all are of the second degree. If any one of them is a rational integral function of the rest it must be a linear function of  $a^2, aa', a'^2$  and of the other three of (3) to (6). Now it is clear that every one of (3) to (6) consists of terms which do not occur in the rest or in  $a^2, aa', a'^2$ .

The six, however, are not all independent, but are connected by one syzygy. For, combining (1) to (5) so as to get an expression free from  $c$  and  $c'$ , we find

$$aa'(ac' + a'c - 2bb') - a^2(a'c' - b'^2) - a'^2(ac - b^2) = (ab' - a'b)^2. \quad \dots (7)$$

We have still to see that any seminvariant or invariant can be expressed rationally and integrally in terms of (1) to (6). Writing  $C, C', I$  for (3), (4) and (5), we have

$$c = \frac{C + b^2}{a}, \quad c' = \frac{C' + b'^2}{a'},$$

so that any rational integral function of  $a, b, c, a', b', c'$  may be written as a sum of such terms as

$$\lambda a^m a'^n b^p b'^q (C + b^2)^r (C' + b'^2)^s,$$

where the indices are integers, zero allowed, and, with the exception of  $m$  and  $n$ , certainly positive. This again is a sum of such terms as

$$\lambda' a^m a'^n b^p b'^q C^r C'^s.$$

If the sum is a seminvariant or invariant it is annihilated by

$$\left(a \frac{d}{db} + 2b \frac{d}{dc}\right) + \left(a' \frac{d}{db'} + 2b' \frac{d}{dc'}\right),$$

which also annihilates  $a, a', C, C'$  separately, and so in its effect on the sum is the same as

$$a \frac{d}{db} + a' \frac{d}{db'}$$

on it as a function of  $b$  and  $b'$  as they occur explicitly. Now the annihilation by this implies, as the theory of partial differential equations tells us, that  $b$  and  $b'$  only occur in the connexion

$$ab' - a'b.$$

Any seminvariant or invariant is then a sum of such terms as

$$\mu \alpha^{m'} a'^{n'} (ab' - a'b)^r C'^r C'^{\sigma'},$$

and consequently, by use of the syzygy (7), can be written as a sum of terms each belonging to one of the types

$$\mu' a^\alpha a'^\beta C^\rho C'^\sigma \Gamma^\tau,$$

$$\mu' a^\alpha a'^\beta (ab' - a'b) C^\rho C'^\sigma \Gamma^\tau.$$

Terms of both types cannot occur, for the whole weights of the two types are one even and the other odd.

Thus a seminvariant or invariant is, either a sum of terms like

$$\mu' a^\alpha a'^\beta C^\rho C'^\sigma \Gamma^\tau,$$

or such a sum multiplied by  $ab' - a'b$ . Here  $\rho, \sigma, \tau$  are positive integers, zero not excluded, and  $\alpha, \beta$  are integral or zero, but not yet proved positive. The factors  $a, a', C, C', \Gamma$  of any term are absolutely independent, and the last three are invariants.

We have to see that for no term as above can  $\alpha$  or  $\beta$  be negative. Suppose if possible that there are terms in which  $\alpha$ , for instance, is negative, and let  $a'$  be the greatest positive value of  $-a$  which occurs in any term. We remember that the seminvariant must lead a covariant, and that the last coefficient in the covariant is, but for a numerical factor, obtained by operating on the seminvariant with  $(O + O')^\varpi$ , where

$$O + O' \equiv \left(2b \frac{d}{da} + c \frac{d}{db}\right) + \left(2b' \frac{d}{da'} + c' \frac{d}{db'}\right),$$



and  $\varpi$  is the order of the covariant, i.e.  $\Sigma(ip) - 2w$ , in this case  $2(\alpha + \beta)$ , which is accordingly non-negative and constant throughout. This last coefficient is annihilated by  $O + O'$ , so that the seminvariant is annihilated by  $(O + O')^{\varpi+1}$ . Now the result of operating with this upon it, if as supposed it contains a sum of terms

$$a^{-\alpha'} \Sigma(a'^{\beta} C^{\rho} C'^{\sigma} \Gamma^{\tau})$$

and none with the factor  $a^{-\alpha'-1}$ , contains the terms

$$(-1)^{\varpi+1} a' (a' + 1) \dots (a' + \varpi) a^{-\alpha'-\varpi-1} \Sigma(a'^{\beta} C^{\rho} C'^{\sigma} \Gamma^{\tau}) 2^{\varpi+1} b^{\varpi+1},$$

and no other terms involving  $a^{-\alpha'-\varpi-1}$  against which they can cancel. The result then cannot vanish; and the supposition was unsound.

It is then completely established that any seminvariant or invariant is either a rational integral function of (1), (2), (3), (4), (5), or such a function multiplied by (6).

The system (1) to (6) is then the complete system of irreducible seminvariants and invariants.

Consequently also the complete system of irreducible covariants and invariants consists of the two quadratics themselves, the three invariants (3), (4), (5), and a third covariant, the Jacobian

$$(ab' - a'b)x^2 + (ac' - a'c)xy + (bc' - b'c)y^2.$$

258.] **Canonical forms of two quadratics.** It is easily seen that by a linear substitution of modulus unity the two quadratics can be given the simultaneous forms

$$aX^2 + cY^2, \quad a'X^2 + c'Y^2,$$

with new values of  $a, c, a', c'$ .

For to make simultaneously

$$all' + b(lm' + l'm) + cmm' = 0,$$

$$a'll' + b'(lm' + l'm) + c'mm' = 0,$$

we have only so to choose  $l : m$  as to make

$$\frac{al + bm}{a'l + b'm} = \frac{bl + cm}{b'l + c'm},$$

i.e. to solve a quadratic, and then to take for  $l' : m'$

$$-\frac{bl + cm}{al + bm}.$$

The absolute values of  $l$ ,  $m$  and  $l'$ ,  $m'$ , whose ratios in pairs are thus determined, may then be taken, and still with one degree of freedom, so as to make  $lm' - l'm = 1$ .

For these canonical forms of unit modulus the six concomitants are

$$\begin{aligned} aX^2 + cY^2 & , \\ a'X^2 + c'Y^2 & , \\ ac & , \\ a'c' & , \\ ac' + a'c & , \\ (ac' - a'c)XY. & \end{aligned}$$

The  $X$  and  $Y$  of the canonical forms are then factors of the Jacobian.

The failing case of two quadratics which are special in that their Jacobian has equal roots is left as an exercise to the student.

Ex. 1. Interpret geometrically the vanishing of the invariant  $ac' + a'c - 2bb'$ .

Ex. 2. The Jacobian of two quadratics represents the double elements of the involution which they determine.

Ex. 3. Express the seminvariant  $ab^2 - 2ba'b' + ca'^2$  in terms of (1) to (6).

$$\text{Ans.} \quad a'(ac' + a'c - 2bb') - a(a'c' - b'^2).$$

Ex. 4. By means of the canonical forms prove that the covariant led by the seminvariant of Ex. 3, represents the harmonic conjugates of the factors of  $ax^2 + 2bxy + cy^2$  with regard to  $a'x^2 + 2b'xy + c'y^2$ .

Ex. 5. Prove the same by means of § 251 (3).

Ex. 6. Find the covariant which consists of the harmonic conjugates of the factors of  $v$  with regard to  $u$ .

Ex. 7. Prove that the covariants of Ex. 4 and Ex. 6 are quadratics which belong to the involution of Ex. 2.

Ex. 8. Express the eliminant  $(ac' - a'c)^2 - 4(ab' - a'b)(bc' - b'c)$  of  $u$  and  $v$  in terms of the invariants (3), (4), (5) of § 257.

Ans.  $(ac' + a'c - 2bb')^2 - 4(ac - b^2)(a'c' - b'^2)$ . Use the canonical forms.

259.] **Linear form and two quadratics.** Let the quadratics be as before

$$u \equiv ax^2 + 2bxy + cy^2,$$

$$v \equiv a'x^2 + 2b'xy + c'y^2,$$

and the linear form  $\xi x + \eta y$ .

Of the linear form alone the one seminvariant is

$$\xi,$$

and of the quadratics the irreducible concomitants are

$$ax^2 + 2bxy + cy^2,$$

$$a'x^2 + 2b'xy + c'y^2,$$

$$(ab' - a'b)x^2 + (ac' - a'c)xy + (bc' - b'c)y^2,$$

$$ac - b^2,$$

$$a'c' - b'^2,$$

$$ac' + a'c - 2bb'.$$

As in § 248 the invariants of the system are these last three, and the eliminants of  $\xi x + \eta y$ , and the preceding three, i. e.

$$a\eta^2 - 2b\eta\xi + c\xi^2,$$

$$a'\eta^2 - 2b'\eta\xi + c'\xi^2,$$

$$(ab' - a'b)\eta^2 - (ac' - a'c)\eta\xi + (bc' - b'c)\xi^2;$$

and as in §§ 253, 254 the remaining irreducible seminvariants of the system are the successive  $\eta$ -derivatives of these three last, i. e.

$$a\eta - b\xi,$$

$$a,$$

$$a'\eta - b'\xi,$$

$$a',$$

$$2(ab' - a'b)\eta - (ac' - a'c)\xi,$$

$$ab' - a'b.$$

The covariants, which these lead are readily written down. The last, last but two, and last but four, occur above.

Ex. 9. The vanishing of the invariant

$$(ab' - a'b)\eta^2 - (ac' - a'c)\eta\xi + (bc' - b'c)\xi^2$$

expresses that the linear form is a double element of the involution determined by the quadratics.

Ex. 10. The covariant  $(a\eta - b\xi)x + (b\eta - c\xi)y$  represents the harmonic conjugate of  $\xi x + \eta y$  with regard to  $u$ .

Ex. 11. Interpret the linear covariants led by

$$a'\eta - b'\xi, \quad 2(ab' - a'b)\eta - (ac' - a'c)\xi.$$

260.] System of quadratic and cubic. Take the forms

$$u \equiv ax^2 + 2bxy + cy^2,$$

$$v \equiv a'x^3 + 3b'x^2y + 3c'xy^2 + d'y^3.$$

For the quadratic only the irreducible concomitants are

$$(1) \quad u,$$

and the invariant

$$(2) \quad ac - b^2.$$

For the cubic only they are

$$(3) \quad v,$$

the covariants

$$(4) \quad (a'c' - b'^2)x^2 + (a'd' - b'c')xy + (b'd' - c'^2)y^2,$$

$$(5) \quad (a'^2d' - 3a'b'c' + 2b'^3)x^3 + \dots,$$

and the invariant

$$(6) \quad (a'd' - b'c')^2 - 4(a'c' - b'^2)(b'd' - c'^2).$$

The remaining irreducible concomitants of the system prove to be nine in number. They may be taken to be the following

(7) the Jacobian of  $u$  and  $v$

$$(ab' - a'b)x^3 + \dots,$$

(8) the Jacobian of  $u$  and (4)

$$\{a(a'd' - b'c') - 2b(a'c' - b'^2)\}x^2 + \dots,$$

(9) the result of operating with  $u$  on  $v$ , after substituting

$\frac{d}{dy}, -\frac{d}{dx}$  for  $x, y$  in  $u$ ,

$$(ac' + a'c - 2bb')x + (ad' + cb' - 2bc')y,$$

(10) the result of operating with (9) on  $u$

$$- \{a^2d' - 3abc' + (ac + 2b^2)b' - bca'\}x \\ + \{c^2a' - 3ccb' + (ac + 2b^2)c' - abd'\}y,$$

(11) the result of operating with  $u$  on (5)

$$\{a(-a'c'd' + 2b'^2d' - b'c'^2) + 2b(-a'b'd' + 2a'c'^2 - b'^2c') \\ + c(a'^2d' - 3a'b'c' + 2b'^3)\}x - \{c(-a'b'd' + 2a'c'^2 - b'^2c') \\ + 2b(-a'c'd' + 2b'^2d' - b'c'^2) + a(a'd'^2 - 3b'c'd' + 2c'^3)\}y,$$

(12) the result of operating with (11) on  $u$ , a linear covariant (the fourth) of degree 2 in the coefficients of  $u$  and 3 in those of  $v$ ,

(13) the intermediate invariant  $AC' + A'C - 2BB'$  of  $u$  and (4), i. e.

$$a(b'd' - c'^2) - b(a'd' - b'c') + c(a'c' - b'^2),$$

(14) an invariant of partial degrees 3, 2, the eliminant of  $u$  and  $v$ ,

(15) an invariant of partial degrees 3, 4, the eliminant of the two linear covariants (9) and (12), or (10) and (11).

Of the system five are invariants, (2), (6), (13), (14), (15), four are linear covariants (9), (10), (11), (12), three are quadratics,  $u$ , (4), (8), and three cubics  $v$ , (5) and (7).

The above is Salmon's list (*Higher Algebra*, § 198). They are, though all irreducible, connected by many syzygies, which have been fully exhibited by Hammond (*Am. J.* vol. viii). There is of course a considerable freedom allowed in choosing the complete list of fifteen, it being allowable to take, in place of any one of the more complicated ones above, any linear function of that one and compounds of the right order and partial degrees of those that are simpler. In fact Hammond finds it convenient to modify the last linear covariant (12) and the last invariant but one (14) by addition of products of others of the set in a way suggested in the next article.

261.] We may, in accordance with the chapter on canonical forms, reduce the cubic by linear transformation of modulus unity to the form

$$a'x^3 + d'y^3,$$

with different  $a'$ ,  $d'$ ,  $x$ ,  $y$ . The same substitution does not affect the form of

$$ax^2 + 2bxy + cy^2,$$

but only makes the  $a$ ,  $b$ ,  $c$  different.

For purposes, then, of the study of the combinations of the concomitants, it suffices to consider  $u$ ,  $v$  in the forms

$$u \equiv ax^2 + 2bxy + cy^2,$$

$$v \equiv a'x^3 + d'y^3.$$

With this simplification it is easy to form all of (1) to (15) by the methods described. In the following notation  $I, L, Q, C$  denote respectively invariants and linear, quadratic, and cubic covariants. The first suffix denotes degree in the coefficients of  $u$ , and the second degree in those of  $v$ . The list is, in the same order as before,

- (1)  $Q_{10} \equiv u \equiv ax^2 + 2bxy + cy^2$ ,
- (2)  $I_{20} \equiv ac - b^2$ ,
- (3)  $C_{01} \equiv a'x^3 + d'y^3$ ,
- (4)  $Q_{02} \equiv a'd'xy$ ,
- (5)  $C_{03} \equiv a'd'(a'x^3 - d'y^3)$ ,
- (6)  $I_{04} \equiv a'^2d'^2$ ,
- (7)  $C_{11} \equiv -b(a'x^3 - d'y^3) - (ca'x - ad'y)xy$ ,
- (8)  $Q_{12} \equiv a'd'(ax^2 - cy^2)$ ,
- (9)  $L_{11} \equiv ca'x + ad'y$ ,
- (10)  $L_{21} \equiv (bca' - a^2d')x - (abd' - c^2a')y$   
 $\equiv b(ca'x - ad'y) - (a^2d'x - c^2a'y)$ ,
- (11)  $L_{13} \equiv a'd'(ca'x - ad'y)$ ,
- (12)  $L_{23} \equiv a'd' \{(a^2d' + bca')x + (abd' + c^2a')y\}$   
 $\equiv a'd'(a^2d'x + c^2a'y) + ba'd'(ca'x + ad'y)$   
 $\equiv a'd'(a^2d'x + c^2a'y) - I_{12}L_{11}$ ,
- (13)  $I_{12} \equiv -ba'd'$ ,
- (14)  $I_{32} \equiv a^3d'^2 + (6abc - 8b^3)a'd' + c^3a'^2$   
 $\equiv a^3d'^2 - 2abca'd' + c^3a'^2 - 8I_{12}I_{20}$ ,
- (15)  $I_{34} \equiv a'd'(c^3a'^2 - a^3d'^2)$ .

Mr. Hammond's modification is to take instead of  $L_{23}$  and  $I_{32}$  the concomitants of like type, but simpler canonical shape,

$$L'_{23} \equiv L_{23} + I_{12}L_{11} \equiv a'd'(a^2d'x + c^2a'y),$$

$$I'_{32} \equiv I_{32} + 8I_{12}I_{20} \equiv a^3d'^2 - 2abca'd' + c^3a'^2.$$

This last is the eliminant of (9) and (10). Its full value in the notation of § 260 is

$$a^3d'^2 + c^3a'^2 - 6a^2bc'd' - 6bc^2a'b' + 2(ac + 2b^2)(ab'd' + ca'c')$$

$$+ (ac + 8b^2)(ac'^2 + cb'^2) - 2abca'd' - 2b(5ac + 4b^2)b'c'.$$

Many good exercises in geometrical interpretation are afforded by the above canonical expressions.

262.] **Linear form quadratic and cubic.** From the system for the quadratic and cubic, the system for them and a linear form  $\xi x + \eta y$ , is derived exactly as in § 259.

The invariants of the system are the invariants  $I_{20}, I_{04}, I_{12}, I_{32}, I_{34}$  of the quadratic and cubic, and the results of replacing  $x$  by  $\eta$  and  $y$  by  $-\xi$  in the quadratic and cubic and their covariants. The other seminvariants are  $\xi$  and the successive derivatives of the invariants which contain  $\xi, \eta$  with regard to  $\eta$ .

Ex. 12. Any seminvariant of  $ax^3 + 3bx^2y + 3cxy^2 + dy^3$  is a rational integral function of  $a$  and invariants of the system

$$ax + by, ax^2 + 2bxy + cy^2, ax^3 + 3bx^2y + 3cxy^2 + dy^3. \quad (\text{Kempe.})$$

Ans. It is a rational integral function of

$$a, ac - b^2, a^2d - 3abc + 2b^3, \text{ and } (ad - bc)^2 - 4(ac - b^2)(bd - c^2),$$

of which the second and fourth are invariants of the quadratic and cubic respectively, and the third is the invariant of the linear form quadratic and cubic given by § 260 (9).

Ex. 13. Any seminvariant of  $(a, b, c, d, e)(x, y)^4$  is a rational integral function of  $a$  and invariants of the system consisting of the quartic and its successive derivatives with regard to  $x$ . (Kempe.)

Ans. Since it is a rational integral function of

$$a, ac - b^2, a^2d - 3abc + 2b^3, ae - 4bd + 3c^2, ace + 2bcd - ad^2 - b^2e - c^3.$$

Ex. 14. Express the seminvariant  $a^2f - 5abe + 2acd + 8b^2d - 6bc^2$  of extent 5 as an invariant of the quintic and its successive derivatives with regard to  $x$ .

Ans. Form a covariant of the system by operating on the quintic with its first  $x$ -derivative, and substitute  $-b$  for  $x$  and  $a$  for  $y$ .

Ex. 15. All invariants of a linear form a quadratic and a cubic are functions, not necessarily rational integral functions, of the discriminant of the quadratic and the eliminants of the linear form with the quadratic, the cubic, the Hessian and cubicovariant of the latter, and the Jacobian of the quadratic and cubic. (Forsyth.)

263.] **Case of  $p+1$  binary quantics of orders  $p, p-1, p-2, \dots, 1, 0$ .** An algebraically complete system of semin-





252, of sets of  $p, p-1, p-2, \dots, 2, 1, 1$  protomorphs of the  $p+1$  quantics, and the Jacobians of the linear one and the  $p-1$  of higher orders.

264.] **Systems of quantics of one order. Combinants.** We will conclude this chapter by allusion, without developement, to an important class of invariants of several binary  $p$ -ics, to which their first discoverer, Sylvester, has given the name *Combinants*. There are also combinants of systems of  $q$ -ary  $p$ -ics, for any the same  $q$  as well as the same  $p$ .

A combinant of a number of  $p$ -ics  $u, v, w, \dots$ , in the same variables, is an invariant which differs only by a function of  $\lambda, \mu, \nu, \dots, \lambda', \mu', \nu', \dots, \dots$  as factor, by a power of

$$\begin{vmatrix} \lambda, \mu, \nu, \dots \\ \lambda', \mu', \nu', \dots \\ \lambda'', \mu'', \nu'', \dots \\ \dots \end{vmatrix}$$

in fact, from the same invariant of

$$\lambda u + \mu v + \nu w + \dots, \quad \lambda' u + \mu' v + \nu' w + \dots, \\ \lambda'' u + \mu'' v + \nu'' w + \dots, \dots$$

It is, in fact, an invariant quâ linear transformation of the  $p$ -ics as well as quâ linear transformations of the variables.

If  $a, b, c, \dots; a', b', c', \dots; a'', b'', c'', \dots; \dots$  be corresponding coefficients in  $u, v, w, \dots$  it is readily seen that the conditions for an invariant to be a combinant are that it have the pairs of annihilators

$$a' \frac{d}{da} + b' \frac{d}{db} + c' \frac{d}{dc} + \dots, \\ a \frac{d}{da'} + b \frac{d}{db'} + c \frac{d}{dc'} + \dots,$$

corresponding to pairs of the  $p$ -ics  $u, v, w, \dots$ .

There are also combinant covariants.

265.] A few of the more obvious facts with regard to combinants are the following.

The eliminant or resultant of two binary  $p$ -ics  $u, v$  is a combinant. For if  $u, v$  have a common factor so have  $\lambda u + \mu v, \lambda' u + \mu' v$ , so that the eliminant of  $u, v$  is a factor

of that of  $\lambda u + \mu v$ ,  $\lambda' u + \mu' v$ . The remaining factor involves  $\lambda, \mu, \lambda', \mu'$  only, as we are told by consideration of dimensions in the coefficients of  $u, v$ .

The eliminant or resultant of three ternary  $p$ -ics is also a combinant; and so on.

A combinant of  $u, v, w, \dots$  is of equal partial degrees in the coefficients of  $u, v, w, \dots$  separately. For, if we denote

$$\begin{aligned} a' \frac{d}{da} + b' \frac{d}{db} + c' \frac{d}{dc} + \dots, \\ a \frac{d}{da'} + b \frac{d}{db'} + c \frac{d}{dc'} + \dots \end{aligned}$$

by  $\phi', \phi$  respectively,

$$\begin{aligned} \phi' \phi - \phi \phi' = & \left( a' \frac{d}{da'} + b' \frac{d}{db'} + c' \frac{d}{dc'} + \dots \right) \\ & - \left( a \frac{d}{da} + b \frac{d}{db} + c \frac{d}{dc} + \dots \right), \end{aligned}$$

whose effect is, by Euler's theorem, the same as that of the multiplier  $i' - i$ . Thus if  $\phi C = 0$  and  $\phi' C = 0$  the first and second partial degrees  $i, i'$  of  $C$  are equal. In like manner the first and third, the first and fourth, &c., partial degrees are equal.

An intermediate invariant (§§ 18, 19) is *not* a combinant. Consider two  $p$ -ics  $u, v$  only. The operation  $\phi$  repeated a number of times on an intermediate invariant produces the invariant of  $u$  only, between which and the same invariant of  $v$  the supposed invariant is intermediate. This intermediate invariant is not then annihilated by  $\phi$ . In like manner as to intermediate invariants of more  $p$ -ics than two.

Let  $I$  be any invariant of  $u$ , and form the same invariant of  $\lambda u + \mu v$ , as in §§ 18, 19. This is

$$\lambda^i I + \frac{1}{1} \lambda^{i-1} \mu \phi' I + \frac{1}{1 \cdot 2} \lambda^{i-2} \mu^2 \phi'^2 I + \dots + \frac{1}{i!} \mu^i \phi'^i I,$$

or, as it may be also written,

$$\frac{1}{i!} \lambda^i \phi^i I' + \frac{1}{(i-1)!} \lambda^{i-1} \mu \phi^{i-1} I' + \dots + \frac{1}{1} \lambda \mu^{i-1} \phi I' + \mu^i I'.$$

Call it

$$A \lambda^i + i B \lambda^{i-1} \mu + \frac{i(i-1)}{1 \cdot 2} C \lambda^{i-2} \mu^2 + \dots + i J \lambda \mu^{i-1} + K \mu^i,$$

so that

$$\begin{aligned}\phi' A &= iB, & \phi' B &= (i-1)C, \\ & & \phi' C &= (i-2)D, \dots \phi' J = K, \phi' K = 0.\end{aligned}$$

We thus have

$$\begin{aligned}\phi' F(A, B, C, \dots J, K) &= \frac{dF}{dA} \phi' A + \frac{dF}{dB} \phi' B + \dots + \frac{dF}{dK} \phi' K \\ &= \left( iB \frac{d}{dA} + (i-1)C \frac{d}{dB} + \dots + K \frac{d}{dJ} \right) F(A, B, C, \dots J, K),\end{aligned}$$

where the operator on the right is of the form of  $O$ . In like manner

$$\begin{aligned}\phi F(A, B, C, \dots J, K) \\ = \left( A \frac{d}{dB} + 2B \frac{d}{dC} + \dots + iJ \frac{d}{dK} \right) F(A, B, C, \dots J, K),\end{aligned}$$

where the operator on the right is of the form of  $\Omega$ .

We thus see that any invariant of an invariant of  $\lambda u + \mu v$  regarded as a quantic in  $\lambda, \mu$  is a combinant of  $u$  and  $v$ . For it is an invariant of  $u$  and  $v$ , being a rational integral homogeneous isobaric function of invariants  $A, B, \dots K$ , and it is annihilated by  $\phi$  and  $\phi'$ .

In like manner any invariant of an invariant of  $\lambda u + \mu v + \nu w$  regarded as a quantic in  $\lambda, \mu, \nu$  is a combinant of  $u, v, w$ , by the principles of the next chapter but one; and so in general.

Ex. 16. If  $u, v$  be binary quadratics, the combinant which is the discriminant of the quadratic in  $\lambda, \mu$  which is the discriminant of  $\lambda u + \mu v$  is the eliminant of  $u, v$ . (*Boole.*)

Ex. 17. The lineo-linear invariant of two binary  $p$ -ics is a combinant if  $p$  is odd, but not if  $p$  is even. (*Cayley.*)

Ex. 18. The criterion of an involution

$$\begin{vmatrix} a_1, & b_1, & c_1 \\ a_2, & b_2, & c_2 \\ a_3, & b_3, & c_3 \end{vmatrix}$$

of three binary quadratics is a combinant.

Ex. 19. A combinant of the fewer than  $p+1$   $p$ -ics

$(a_1, b_1, c_1, \dots)(x, y, \dots)^p, (a_2, b_2, c_2, \dots)(x, y, \dots)^p, (a_3, b_3, c_3, \dots)(x, y, \dots)^p, \dots$   
is a function of determinants obtained by erasing columns from

$$\begin{vmatrix} a_1, & b_1, & c_1, & d_1, & e_1, & \dots \\ a_2, & b_2, & c_2, & d_2, & e_2, & \dots \\ a_3, & b_3, & c_3, & d_3, & e_3, & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \end{vmatrix}. \quad (\text{Sylvester.})$$

## CHAPTER XV.

### RESTRICTED SUBSTITUTIONS. METRICAL GEOMETRY OF PENCILS.

266.] BESIDES actual invariants and covariants, which have the invariantic or covariantic connexion with a quantic or quantics whatever be the linear substitution for the variables, there exist functions which possess the property of invariancy or covariancy for particular classes of substitutions.

Thus, for instance, seminvariants of binary quantics are not invariants for all linear substitutions, but are invariantic for the particular class of substitutions  $x = lX + mY$ ,  $y = m'Y$ .

A very important class of quasi-invariants and covariants is that of functions which are invariants and covariants as far as all substitutions are concerned which in Cartesian geometry express change of reference from one set of axes to another, the old and new variables being both sets of coordinates in the ordinary sense.

Confining attention to plane geometry, the most general equations of substitution, those which express change from old axes at an angle  $\omega$  to new axes at an angle  $\omega' = \beta - \alpha$ , through a point  $(h, k)$ , and inclined at angles  $\alpha, \beta$  respectively to the old axis of  $x$ , are

$$x = X \frac{\sin(\omega - \alpha)}{\sin \omega} + Y \frac{\sin(\omega - \beta)}{\sin \omega} + [Z] h,$$

$$y = X \frac{\sin \alpha}{\sin \omega} + Y \frac{\sin \beta}{\sin \omega} + [Z] k,$$

$$[z] = \qquad \qquad \qquad [Z] = 1.$$

Of these, whether  $h, k$  be present or absent, i. e. whether the

substitution be taken as ternary or binary, the modulus is

$$\begin{aligned} \frac{\sin(\omega - a) \sin \beta - \sin(\omega - \beta) \sin a}{\sin^2 \omega} &= \frac{\sin \omega \sin(\beta - a)}{\sin^2 \omega} \\ &= \frac{\sin(\beta - a)}{\sin \omega} = \frac{\sin \omega'}{\sin \omega}. \end{aligned}$$

267.] **Boolean and orthogonal invariants, &c.** The study of certain invariants and covariants for Cartesian transformations preceded and led to the investigation of invariants and covariants generally. The chief early contribution to the study is in a paper by Boole (*Cambridge Math. Journal*, Vol. III), to which reference has already been made in § 18. It is proposed here to give the name *Boolean* invariants and covariants to functions which have the restricted invariantic and covariantic properties contemplated, the name being given without in the least implying that Boole confined his attention to such restricted invariants and covariants. His work not only led to the development of the more general invariant algebra, but began that development.

The original theorem was that a binary quadratic

$$ax^2 + 2bxy + cy^2$$

has the two Boolean invariants

$$ac - b^2, \quad a + c - 2b \cos \omega;$$

in fact that, if  $a'X^2 + 2b'XY + c'Y^2$  is the quadratic transformed so as to be referred to new axes at an angle  $\omega'$ ,

$$a'c' - b'^2 = \left(\frac{\sin \omega'}{\sin \omega}\right)^2 (ac - b^2),$$

$$a' + c' - 2b' \cos \omega' = \left(\frac{\sin \omega'}{\sin \omega}\right)^2 (a + c - 2b \cos \omega).$$

The first of these Boolean invariants is known to be an invariant for all linear substitutions. The second is not so.

Boole's well-known method depends on the fact that, the transformation being a binary one, i. e. one with no change of origin,

$$x^2 + 2xy \cos \omega + y^2 = X^2 + 2XY \cos \omega' + Y^2.$$

He in fact determines, by his method § 18, the invariants in

the ordinary sense of the system

$$ax^2 + 2bxy + cy^2,$$

$$x^2 + 2xy \cos \omega + y^2.$$

Quite generally, the Boolean invariants and covariants of a binary quantic, among which are included the full invariants and covariants of that quantic, are the full invariants and covariants of the system consisting of that quantic and the quadratic

$$x^2 + 2xy \cos \omega + y^2.$$

If we take  $\cos \omega' = \cos \omega = 0$ , i. e. if we consider only transformations from one pair to another of *rectangular* axes, Boolean invariants and covariants take particular forms which may be called *orthogonal* invariants and covariants.

268.] As a first instance of a Boolean covariant we may take the Jacobian of the quadratic

$$ax^2 + 2bxy + cy^2,$$

and

$$x^2 + 2xy \cos \omega + y^2,$$

which proves to be

$$(a \cos \omega - b)x^2 + (a - c)xy + (b - c \cos \omega)y^2.$$

This, then, is a pair of lines having an invariable geometrical relation to the pair of lines denoted by the given quadratic and the pair of lines  $x^2 + 2xy \cos \omega + y^2$  to the circular points at infinity.

To see what the relation is take  $2xy$  for the given quadratic. The Boolean covariant is at once

$$y^2 - x^2,$$

and so represents the bisectors of the angles between the lines forming the quadratic  $2xy$ . These bisectors are presented as the common harmonic conjugates of the lines  $xy$  and

$$x^2 + 2xy \cos \omega + y^2.$$

Quite generally, a Boolean covariant represents a pencil of lines having an invariable geometrical relation to the pencil represented by a binary quantic and the pencil

$$x^2 + 2xy \cos \omega + y^2$$

from the vertex to the circular points at infinity, i. e. represents

a pencil having an invariable relation to the pencil represented by the quantic, into the expression of which relation magnitudes of angles enter or may enter as well as descriptive and projective connexions. Conversely, any such pencil is represented by a Boolean covariant.

The vanishing of a Boolean *invariant* expresses a geometrical relation between the lines of a pencil denoted by the binary quantic to which the invariant belongs, into the expression of which relation magnitudes of angles may enter as well as descriptive and projective connexions.

For instance,  $a + c - 2b \cos \omega = 0$  expresses that a quadratic denotes lines harmonically conjugate with regard to the lines to the circular points, i.e. denotes lines at right angles.

269.] The method of emanants (§§ 52, &c.) applies to Boolean invariants and covariants. It was proved that any invariant of an emanant of  $u$  is a covariant, or, in particular, invariant, of  $u$ . Now, just as this was seen, it follows also that any function which is for a restricted class of substitutions an invariant of the emanant is for the same class of substitutions a covariant, or invariant, of  $u$ . In particular, this is the case for the substitutions of Cartesian geometry.

For instance, the second emanant of  $u$

$$x'^2 \frac{d^2 u}{dx^2} + 2x'y' \frac{d^2 u}{dxdy} + y'^2 \frac{d^2 u}{dy^2}$$

has the Boolean invariant

$$\frac{d^2 u}{dx^2} + \frac{d^2 u}{dy^2} - 2 \frac{d^2 u}{dxdy} \cos \omega.$$

This, then, is a Boolean covariant of the binary quantic  $u$ , if the order of  $u$  exceeds 2. For the order 2 of  $u$  it is the Boolean invariant used in its production.

The reasoning as to the power of the modulus in § 56 still applies. Thus the expression of the fact of covariancy of the function before us is

$$\begin{aligned} \frac{d^2 u}{dX^2} + \frac{d^2 u}{dY^2} - 2 \frac{d^2 u}{dXdY} \cos \omega' \\ = \left( \frac{\sin \omega'}{\sin \omega} \right)^2 \left\{ \frac{d^2 u}{dx^2} + \frac{d^2 u}{dy^2} - 2 \frac{d^2 u}{dxdy} \cos \omega \right\}. \end{aligned}$$

In particular,  $\frac{d^2u}{dx^2} + \frac{d^2u}{dy^2}$  is an *orthogonal* covariant, i.e., since  $\left(\frac{\sin \omega'}{\sin \omega}\right)^2 = 1$  for all orthogonal substitutions, is unaltered by any change of rectangular axes without change of origin.

270.] **Cogrediency identical with contragrediency for orthogonal substitutions.** This last fact as to an orthogonal covariant is a case of an interesting theorem due to Boole.

It should be noticed that there are two discrete classes of orthogonal substitutions, which may be called *direct* and *skew* respectively. In the direct class the sense of rotation from the axis of  $X$  to that of  $Y$  is the same as that from the axis of  $x$  to that of  $y$ , while in the skew class the senses of rotation are opposite. The modulus for the direct class is  $+1$ , whereas that for the skew class is  $-1$ .

The formulae for orthogonal substitution are

$$x = X \cos \theta \mp Y \sin \theta,$$

$$y = X \sin \theta \pm Y \cos \theta,$$

where the upper signs refer to direct and the lower to skew substitutions. From these there follow

$$\frac{d}{dX} = \cos \theta \frac{d}{dx} + \sin \theta \frac{d}{dy},$$

$$\frac{d}{dY} = \mp \sin \theta \frac{d}{dx} \pm \cos \theta \frac{d}{dy},$$

so that

$$\frac{d}{dx} = \cos \theta \frac{d}{dX} \mp \sin \theta \frac{d}{dY},$$

$$\frac{d}{dy} = \sin \theta \frac{d}{dX} \pm \cos \theta \frac{d}{dY}.$$

Thus for all orthogonal substitutions  $\frac{d}{dx}$  and  $\frac{d}{dy}$  are cogredient with  $x$  and  $y$  (§ 51). For orthogonal substitutions then contragrediency is identical with cogrediency (cf. §§ 46, 68). This is readily seen to be the case if we take  $\xi, \eta$ , any quantities or symbols contragredient with  $x$  and  $y$ , instead of  $\frac{d}{dx}$  and  $\frac{d}{dy}$  in particular.



The application of the cogrediency now before us is as follows. If the result of transforming a binary quantic orthogonally is

$$u \equiv (a_0, a_1, a_2, \dots a_p)(x, y)^p = (A_0, A_1, A_2, \dots A_p)(X, Y)^p, \dots (1)$$

$$\text{while of course} \quad x^2 + y^2 = X^2 + Y^2, \quad \dots (2)$$

and the modulus of the substitution is in magnitude  $\cos^2 \theta + \sin^2 \theta$ , i.e. unity, we have also

$$(a_0, a_1, a_2, \dots a_p) \left( \frac{d}{dx}, \frac{d}{dy} \right)^p \\ = (A_0, A_1, A_2, \dots A_p) \left( \frac{d}{dX}, \frac{d}{dY} \right)^p \quad \dots (3)$$

$$\text{and} \quad \left( \frac{d}{dx} \right)^2 + \left( \frac{d}{dy} \right)^2 = \left( \frac{d}{dX} \right)^2 + \left( \frac{d}{dY} \right)^2. \quad \dots (4)$$

Moreover, if  $K(a_0, a_1, a_2, \dots a_p)(x, y)^w$  be any covariant or orthogonal covariant,

$$K(a_0, a_1, a_2, \dots a_p) \left( \frac{d}{dx}, \frac{d}{dy} \right)^w \\ = \pm K(A_0, A_1, A_2, \dots A_p) \left( \frac{d}{dX}, \frac{d}{dY} \right)^w, \quad \dots (5)$$

the upper sign being correct except when both the covariant and the substitution are skew.

Thus by operation with any combination of the left-hand sides of (3) and (4), or with any covariant operator such as the left of (5), on any combination of the left-hand sides of (1) and (2) or on any covariant or orthogonal covariant, we obtain an orthogonal covariant or invariant.

Consider in particular the quadratic

$$ax^2 + 2bxy + cy^2.$$

We have by (3) on (1) an orthogonal invariant

$$a^2 + 2b^2 + c^2,$$

and by (4) on (1) or (3) on (2) another

$$a + c.$$

The known one  $ac - b^2$  is, of course,

$$\frac{1}{2} \{ (a + c)^2 - (a^2 + 2b^2 + c^2) \}.$$

Again, from the cubic

$$ax^3 + 3bx^2y + 3cxy^2 + dy^3$$

we have by (3) on (1) an orthogonal invariant

$$a^2 + 3b^2 + 3c^2 + d^2,$$

by (4) on (1) an orthogonal covariant

$$(a+c)x + (b+d)y,$$

and again, by operating on this with  $(a+c)\frac{d}{dx} + (b+d)\frac{d}{dy}$ , another orthogonal invariant

$$(a+c)^2 + (b+d)^2.$$

For another example take the quintic  $(a, b, c, d, e, f)(x, y)^5$ . We obtain at once the orthogonal invariant

$$a^2 + 5b^2 + 10c^2 + 10d^2 + 5e^2 + f^2,$$

a cubic orthogonal covariant which leads to the orthogonal invariant  $(a+c)^2 + 3(b+d)^2 + 3(c+e)^2 + (d+f)^2$ ,

and from it again a linear orthogonal covariant and the orthogonal invariant

$$(a+2c+e)^2 + (b+2d+f)^2,$$

besides two other orthogonal invariants obtained by operations with orthogonal covariants found above on others.

The number of independent orthogonal invariants including full invariants of the binary  $p$ -ic is  $p$ . The difficulty of discovering the independence or interdependence of orthogonal invariants determined as above, and the investigation of complete irreducible systems, would have to be attacked separately in the case of every order  $p$ .

Ex. 1. In orthogonal transformations in three dimensions prove that  $\frac{d}{dx}, \frac{d}{dy}, \frac{d}{dz}$  are cogredient with  $x, y, z$ . (*Boole.*)

Ex. 2. For orthogonal transformations covariants and contravariants coincide. (*Sylvester.*)

Ex. 3. The ternary cubic

$$ax^3 + by^3 + cz^3 + 3dx^2y + 3exy^2 + 3fx^2z + 3gxyz + 3hy^2z + 3iyz^2 + 6kxyz$$

has the orthogonal invariants

$$a^2 + b^2 + c^2 + 3(d^2 + e^2 + f^2 + g^2 + h^2 + i^2) + 6k^2,$$

$$(a+e+g)^2 + (b+d+i)^2 + (c+f+h)^2. \quad (\textit{Boole.})$$

271.] **Annihilator of orthogonal covariants and invariants.** Sylvester has expressed by a linear differential equation the condition that a function be an orthogonal covariant or invariant of a binary quantic.

If we express that the function is unaltered, to the first order of infinitesimals, when the substitution is made which effects the turning of the rectangular axes through an infinitesimal angle, we express equally that it is unaltered when we turn them through a finite angle, for, if they be turned through this angle  $a$  by an infinite succession  $n$  of infinitesimal turnings through  $\frac{1}{n}a$ , the whole increment of the function is at most comparable with  $n\left(\frac{1}{n}a\right)^2$ , i.e. is an infinitesimal, and vanishes in the limit.

Now the formulae of substitution for turning through an infinitesimal angle  $\theta$  are

$$\begin{aligned}x &= X - \theta Y, \\y &= \theta X + Y,\end{aligned}$$

omitting infinitesimals of the second order. The modulus of this, to the first order of infinitesimals, is 1. Indeed 1 is its absolute value, as in all cases of direct orthogonal substitution, when infinitesimals of higher orders are expressed in the formulae of substitution. Now these are, to the same order of infinitesimals,

$$\begin{aligned}X &= x + \theta y, \\Y &= y - \theta x.\end{aligned}$$

Thus the substitution amounts to giving  $x$  and  $y$  the increments  $\theta y$  and  $-\theta x$ .

Again, if the quantic under consideration be

$$(a_0, a_1, a_2, \dots a_p)(x, y)^p,$$

the substitution may be effected, correctly to the first order in  $\theta$ , by first writing it

$$(a_0, a_1, a_2, \dots a_p)(X - \theta y, y)^p,$$

which makes it  $(a'_0, a'_1, a'_2, \dots a'_p)(X, y)^p,$

where (§ 91)

$$a'_0 = a_0, a'_1 = a_1 - a_0\theta, a'_2 = a_2 - 2a_1\theta, \dots, a'_p = a_p - pa_{p-1}\theta,$$

and then writing it

$$(a_0', a_1', a_2', \dots a_p')(X, \theta X + Y)^p,$$

i. e.  $(A_0, A_1, A_2, \dots A_p)(X, Y)^p,$

where (§ 94)

$$A_0 = a_0' + pa_1'\theta = a_0 + pa_1\theta, \quad \text{to the first order in } \theta,$$

$$A_1 = a_1' + (p-1)a_2'\theta = a_1 + (p-1)a_2\theta - a_0\theta, \quad \text{,, ,, ,}$$

$$A_2 = a_2' + (p-2)a_3'\theta = a_2 + (p-2)a_3\theta - 2a_1\theta, \quad \text{,, ,, ,}$$

. . . . .

$$A_{p-1} = a_{p-1}' + a_p'\theta = a_{p-1} + a_p\theta - (p-1)a_{p-2}\theta, \quad \text{,, ,}$$

$$A_p = a_p' = a_p - pa_{p-1}\theta, \quad \text{,, ,, .}$$

Thus  $X, Y, A_0, A_1, A_2, \dots A_{p-1}, A_p$  differ from

$$x, y, a_0, a_1, a_2, \dots a_{p-1}, a_p$$

by the increments

$$\delta x = \theta y, \quad \delta y = -\theta x, \quad \delta a_0 = pa_1\theta, \quad \delta a_1 = \{(p-1)a_2 - a_0\}\theta, \\ \delta a_2 = \{(p-2)a_3 - 2a_1\}\theta, \dots, \delta a_p = -pa_{p-1}\theta,$$

whence it follows that the increment of

$$F(x, y, a_0, a_1, a_2, \dots a_p)$$

is  $\theta \left\{ y \frac{d}{dx} - x \frac{d}{dy} + \left( pa_1 \frac{d}{da_0} + (p-1)a_2 \frac{d}{da_1} + \dots + a_p \frac{d}{da_{p-1}} \right) \right. \\ \left. - \left( a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + \dots + pa_{p-1} \frac{d}{da_p} \right) \right\} F.$

The necessary and sufficient condition that  $F$  be a covariant for *direct* orthogonal substitutions is, then, if as usual we adopt the notation of chapter vi, that  $F$  have the annihilator

$$y \frac{d}{dx} - x \frac{d}{dy} + O - \Omega.$$

That it be a covariant also for *skew* orthogonal substitutions a further condition is necessary and sufficient. It must be either unaltered or only changed in sign when  $y$  and the alternate coefficients  $a_1, a_3, a_5, \dots$  have their signs altered. For the most general skew orthogonal transformation may be performed by replacing  $x, y$  by  $x, -y$ , i. e. reversing the direction of the axis of  $y$ , and then performing the most

general direct orthogonal transformation. Should, however, a function  $K_1 + K_2$ , which has the above annihilator, become  $K_1 - K_2$  upon making these changes of sign, it is readily seen that  $K_1 - K_2$  also has the annihilator, since this annihilator is only altered in sign by the changes of sign of  $y$  and  $a_1, a_3, a_5, \dots$ . In this case  $K_1$  and  $K_2$  are orthogonal covariants for skew as well as direct substitutions, while  $K_1 + K_2$  and  $K_1 - K_2$  are not.  $K_1$  and  $K_2$  have in fact for their factors powers of the modulus  $\pm 1$  of which one is even and the other odd.

In particular, orthogonal *invariants* are functions of the coefficients only which have the annihilator  $O - \Omega$  and are either unaltered or changed only in sign when the signs of  $a_1, a_3, a_5, \dots$  are altered.

272.] **Annihilator of Boolean covariants and invariants.** We may also find an annihilator of any Boolean covariant or invariant from the consideration that any one is unchanged when the oblique axes are turned through an infinitesimal angle  $\theta$ .

For such a turning the formulae are, by § 266, since  $a = \theta$ ,  
 $\beta = \omega + \theta$ ,  
 $x = X - X \cot \omega \cdot \theta - Y \operatorname{cosec} \omega \cdot \theta$ ,  
 $y = Y + X \operatorname{cosec} \omega \cdot \theta + Y \cot \omega \cdot \theta$ .

Thus  $x$  and  $y$  have the increments

$$\delta x = x \cot \omega \cdot \theta + y \operatorname{cosec} \omega \cdot \theta, \quad \delta y = -x \operatorname{cosec} \omega \cdot \theta - y \cot \omega \cdot \theta.$$

Also it is readily seen that the increments of  $a_0, a_1, a_2, \dots, a_p$  may be exhibited as follows :

$$\begin{aligned} & \delta a_0, \quad \delta a_1, \quad \delta a_2, \quad \dots, \quad \delta a_{p-1}, \quad \delta a_p \\ = & (-pa_0, -(p-1)a_1, -(p-2)a_2, \dots, -a_{p-1}, \quad 0) \cot \omega \cdot \theta \\ + & (0, -a_0, -2a_1, \dots, -(p-1)a_{p-2}, -pa_{p-1}) \operatorname{cosec} \omega \cdot \theta \\ + & (pa_1, (p-1)a_2, (p-2)a_3, \dots, a_p, \quad 0) \operatorname{cosec} \omega \cdot \theta \\ + & (0, a_1, 2a_2, \dots, (p-1)a_{p-1}, pa_p) \cot \omega \cdot \theta. \end{aligned}$$

Hence the expression of the fact that the increment of a Boolean covariant vanishes is that it is annihilated by

$$\begin{aligned} \cot \omega \left\{ x \frac{d}{dx} - y \frac{d}{dy} - pa_0 \frac{d}{da_0} - (p-2)a_1 \frac{d}{da_1} \right. \\ \left. - (p-4)a_2 \frac{d}{da_2} - \dots + (p-2)a_{p-1} \frac{d}{da_{p-1}} + pa_p \frac{d}{da_p} \right\} \\ + \operatorname{cosec} \omega \left\{ y \frac{d}{dx} - x \frac{d}{dy} + O - \Omega \right\}, \end{aligned}$$

which correctly becomes the annihilator of the preceding article when  $\omega = \frac{\pi}{2}$ , i.e. when the covariant is orthogonal.

In particular, Boolean *invariants* have the annihilator

$$O - \Omega - \cos \omega \left\{ pa_0 \frac{d}{da_0} + (p-2)a_1 \frac{d}{da_1} + \dots \right. \\ \left. - (p-2)a_{p-1} \frac{d}{da_{p-1}} - pa_p \frac{d}{da_p} \right\}.$$

The second part of this annihilator has the effect of multiplying every isobaric part of weight  $w$  of a Boolean invariant of degree  $i$  by  $ip - 2w$ . But this multiplier is not constant throughout, as such Boolean invariants as are not full invariants are not isobaric in the coefficients of the quantic to which they belong. They are what isobaric invariants of the quantic and a quadratic become in a case when the idea of weight is banished from the coefficients of the latter.

Boolean covariants and invariants have also to obey a further law, which is best expressed by saying that they must be unaltered or changed at most in sign when  $x$  and  $y$ ,  $a_0$  and  $a_p$ ,  $a_1$  and  $a_{p-1}$ ,  $a_2$  and  $a_{p-2}$ , &c., are interchanged. The supplementary necessity of the last article as to orthogonal covariants and invariants might have been expressed in the same way.

Ex. 4. By turning the axis of  $y$  through an angle  $d\omega$ , keeping that of  $x$  unchanged, prove that, if  $\mu$  be the index of the power to which the modulus  $\frac{\sin \omega'}{\sin \omega}$  enters in the equality expressing that  $F$  is a Boolean covariant,

$$\mu F = \left\{ \left( a_1 \frac{d}{da_1} + 2a_2 \frac{d}{da_2} + \dots + pa_p \frac{d}{da_p} - y \frac{d}{dy} \right) \right. \\ \left. + \sec \omega \left( y \frac{d}{dx} - \Omega \right) + \tan \omega \frac{d}{d\omega} \right\} F.$$

Ans. Express that the increment of  $(\sin \omega)^{-\mu} F$  vanishes.

273.] **Boolean system for linear form.** We proceed by chapter xiv to write down complete irreducible systems of Boolean invariants and covariants for binary quantics of the first few orders, as invariants and covariants of those quantics and the quadratic  $x^2 + 2xy \cos \omega + y^2$ .

Take first the linear form  $ax + by$ .

By § 251 the system consists of five individuals; viz.

(1) the linear form itself  $ax + by$ ,

(2)  $x^2 + 2xy \cos \omega + y^2$ ,

(3) the linear Boolean covariant

$$(a \cos \omega - b)x + (a - b \cos \omega)y,$$

(4) the Boolean invariant

$$a^2 - 2ab \cos \omega + b^2,$$

(5)  $\sin^2 \omega$ .

Any pencil of lines connected with the given line by descriptive or metrical properties has for its equation a rational integral function of these equated to zero. There are none of course whose connexions with the line are purely descriptive.

The Boolean covariant (3), the Jacobian of (1) and (2), is the perpendicular to  $ax + by$ .

The Boolean invariant (4) is the criterion for  $ax + by$  running to one of the circular points at infinity.

A Boolean covariant or invariant is separately homogeneous in  $x, y$ , in  $a, b, c$ , and in  $1, \cos \omega, 1$ , in which last (1), (2), (3), (4), (5) are of degrees 0, 1, 1, 1, 2 respectively.

274.] **Case of the quadratic.** For

$$u \equiv ax^2 + 2bxy + cy^2$$

the Boolean system is that of invariants and covariants of  $u$  and the second quadratic

$$x^2 + 2xy \cos \omega + y^2,$$

and so (§ 257) consists of

(1) the quadratic itself

$$ax^2 + 2bxy + cy^2,$$

(2)  $x^2 + 2xy \cos \omega + y^2$ ,

(3)  $ac - b^2$ , the one full invariant of  $u$ ,

(4)  $\sin^2 \omega$ ,

(5) the Boolean invariant

$$a + c - 2b \cos \omega,$$

(6) the Jacobian

$$(a \cos \omega - b)x^2 + (a - c)xy + (b - c \cos \omega)y^2.$$

Interpretation has already been given (§ 268) to (5) and (6).

Any other Boolean covariant or invariant is a rational integral function of (1) to (6). For a rational integral function of them to be such a covariant or invariant it must be homogeneous in  $x, y$ , in  $a, b, c$ , and in  $1, \cos \omega, 1$ , separately, (1), (2), (3), (4), (5), (6) being looked upon as of degrees 0, 1, 0, 2, 1, 1 respectively in  $1, \cos \omega, 1$ .

Ex. 5. Interpret the Boolean covariant

$$(a+c-2b \cos \omega)(x^2+2xy \cos \omega+y^2) - \sin^2 \omega (ax^2+2bxy+cy^2).$$

*Ans.* The perpendiculars to the lines  $u$ . To see it take  $\omega = \frac{\pi}{2}$ .

Ex. 6. Interpret the Boolean invariant condition

$$(a+c-2b \cos \omega)^2 - 4(ac-b^2) \sin^2 \omega = 0.$$

*Ans.* One of the lines  $u$  runs to a circular point at infinity.

275.] **Linear form and quadratic.** The Boolean system for

$$\xi x + \eta y \text{ and } ax^2 + 2bxy + cy^2$$

is written down from § 259 by taking  $x^2 + 2xy \cos \omega + y^2$  for  $v$ .

We have, besides the system written down in the preceding article, three Boolean invariants, viz.

$$(7) \quad a\eta^2 - 2b\eta\xi + c\xi^2,$$

$$(8) \quad \eta^2 - 2\eta\xi \cos \omega + \xi^2,$$

$$(9) \quad (a \cos \omega - b)\eta^2 - (a-c)\eta\xi + (b-c \cos \omega)\xi^2,$$

of which (7) is a full invariant, and the following Boolean covariants, of which (10) and (11) are full covariants,

$$(10) \quad \xi x + \eta y,$$

$$(11) \quad (a\eta - b\xi)x + (b\eta - c\xi)y,$$

$$(12) \quad (\eta - \xi \cos \omega)x + (\eta \cos \omega - \xi)y,$$

$$(13) \quad \{2(a \cos \omega - b)\eta - (a-c)\xi\}x \\ + \{(a-c)\eta - 2(b-c \cos \omega)\xi\}y.$$

In terms of these thirteen all Boolean invariants and covariants of the linear form and quadratic can be rationally and integrally expressed. They have to be separately homogeneous in  $x$  and  $y$ , in  $\xi$  and  $\eta$ , in  $a, b, c$ , and in  $1, \cos \omega, 1$ .

Besides (1) to (6) which have been interpreted in the last article, and the full invariant and covariants (7) (10) (11) which have been interpreted in chapter xiv, we have (8) and (12) which have been interpreted as the (4) and (3) of § 273.



We have left (9) which is the Boolean invariant criterion for  $\xi x + \eta y$  being one of the bisectors of the angle between the lines  $ax^2 + 2bxy + cy^2$ , and (13) which is the harmonic conjugate of  $\xi x + \eta y$  with regard to these bisectors, as is readily seen by taking the case when  $\xi = 0$ .

276.] **Case of the cubic.** By § 260 the complete Boolean system for the cubic consists of fifteen individuals. Their forms for the cubic

$$ax^3 + 3bx^2y + 3cxy^2 + dy^3$$

are given by the article referred to, upon replacing  $a', b', c', d'$  by  $a, b, c, d$ , and  $a, b, c$  by  $1, \cos \omega, 1$ .

Let us content ourselves with using § 261 to write them down for the cubic

$$ax^3 + dy^3$$

to whose form the given cubic can certainly be reduced by a change of axes, taking the lines represented by the Hessian as new axes of  $x$  and  $y$ , the one case of exception being the special one when these lines coincide, i.e. when the discriminant of the cubic vanishes. We get

- (1)  $x^2 + 2xy \cos \omega + y^2$ ,
- (2)  $\sin^2 \omega$ ,
- (3)  $ax^3 + dy^3$ , the cubic itself,
- (4)  $adx$ ,
- (5)  $ad(ax^3 - dy^3)$ ,
- (6)  $a^2d^2$ ,
- (7)  $(dy^3 - ax^3) \cos \omega + (dy - ax)xy$ ,
- (8)  $ad(x^2 - y^2)$ ,
- (9)  $ax + dy$ ,
- (10)  $(a \cos \omega - d)x + (a - d \cos \omega)y$ ,
- (11)  $ad(ax - dy)$ ,
- (12)  $ad(dx + ay)$ , where  $L'_{23}$  is chosen,
- (13)  $-ad \cos \omega$ ,
- (14)  $a^2 - 2ad \cos \omega + d^2$ , where  $I'_{32}$  is chosen,
- (15)  $ad(a^2 - d^2)$ .

The degrees in  $1, \cos \omega, 1$  are not all clearly indicated by the

powers to which  $\cos \omega$  enters in these canonized forms. A double suffix notation, as in § 261, should be used in compounding them homogeneously.

277.] The geometrical relations of the members of this system may be expressed in various ways.

The full covariants and invariant of the cubic are (3), (4), (5), (6). Of these (5) represents the harmonic conjugates of the lines composing the cubic each with regard to the other two, and (4) represents the double lines of the involution to which corresponding lines of the cubic and (5) belong. The invariant (6) is the discriminant.

Of the other Boolean covariants (7) represents the three lines whose polar lines with regard to the cubic are at right angles to them respectively; (8) represents the bisectors of the angles between the lines represented by the Hessian (4); (9) is the line whose first polar with regard to the cubic (3) is (8); (11) is the harmonic conjugate of this with regard to the Hessian (4); (12) is the polar line of (11) with regard to the cubic; (10) is at right angles to (9).

Of the Boolean invariants (13), best considered in its full form

$$bd - c^2 - (ad - bc) \cos \omega + ac - b^2$$

for the unreduced cubic, is the criterion that the Hessian (4) consist of lines at right angles, i.e. that (3) and (5) are equiangular and oppositely turned pencils; (14) is the criterion that (9) and (12) be at right angles to themselves, i.e. run to the circular points at infinity; and (15) is the criterion that (9) and (12) coincide.

In terms of all these (1) to (15) any geometrical criterion, and any pencil of lines geometrically connected with the pencil forming the cubic, can be rationally and integrally expressed.

278.] By means of § 262 we may in like manner write down the complete Boolean system of a cubic and a linear form.



chosen as in § 27, where we supposed  $\frac{x}{z}, \frac{y}{z}$  to be each of unit weight. We might equally have adopted suffix notations appropriate to cases when  $\frac{y}{x}, \frac{z}{x}$  in the one case, and  $\frac{z}{y}, \frac{x}{y}$  in the other, are of unit weight. Any fact as to weight of concomitants which may be adduced will have two companion facts, arising from it by changes corresponding to cyclical interchange of  $x, y, z$ . In one sense, indeed, companion facts are sixfold, one corresponding to every permutation of  $x, y, z$ .

In much that follows the notation of the cubic will for simplicity of writing be adopted in our work, and the conclusions only indicated in the notation of the general  $p$ -ic. It is important to have before us three ways in which the same cubic may be arranged, namely,

$$\begin{aligned} & a_3 z^3 \\ & + 3(a_2 x + b_2 y) z^2 \\ & + 3(a_1 x^2 + 2b_1 xy + c_1 y^2) z \\ & + a_0 x^3 + 3b_0 x^2 y + 3c_0 xy^2 + d_0 y^3, \end{aligned} \quad \dots \text{(i)}$$

$$\begin{aligned} & a_0 x^3 \\ & + 3(b_0 y + a_1 z) x^2 \\ & + 3(c_0 y^2 + 2b_1 yz + a_2 z^2) x \\ & + d_0 y^3 + 3c_1 y^2 z + 3b_2 yz^2 + a_3 z^3, \end{aligned} \quad \dots \text{(ii)}$$

$$\begin{aligned} & d_0 y^3 \\ & + 3(c_1 z + c_0 x) y^2 \\ & + 3(b_2 z^2 + 2b_1 zx + b_0 x^2) y \\ & + a_3 z^3 + 3a_2 z^2 x + 3a_1 zx^2 + a_0 x^3. \end{aligned} \quad \dots \text{(iii)}$$

The corresponding triple arrangement is general for the ternary  $p$ -ic.

281.] We first notice that, an invariant or covariant being unaltered, except for a power of the modulus as factor, when we substitute for the coefficients and variables  $x, y, z$  the new coefficients and variables given by any linear substitution for  $x, y, z$  whatever, the same is true in particular when the linear substitution is one affecting  $x$  and  $y$  only, leaving  $z$  unaltered.

Consider, to begin with, invariants only. We are thus told,

using the notation of the cubic, that an invariant is a function of

$$a_3 \quad \dots (1)$$

and the coefficients in the quantics

$$a_2x + b_2y, \quad \dots (2)$$

$$a_1x^2 + 2b_1xy + c_1y^2, \quad \dots (3)$$

$$a_0x^3 + 3b_0x^2y + 3c_0xy^2 + d_0y^3, \quad \dots (4)$$

which is unaffected, except by a power of the modulus as factor, when these quantics are simultaneously linearly transformed. From this we gather that it is a rational integral function of  $a_3$  and invariants of the system (2), (3), (4). Or, regarding  $a_3$  as itself a quantic of zero order, which has itself for its one invariant, we may say that an invariant of the ternary cubic is a rational integral function of invariants of the system (1), (2), (3), (4). It is isobaric on the whole (§ 28), and of course homogeneous on the whole (§ 22), but is not to be expected to be homogeneous in  $a_3$  and the coefficients of (2), (3), (4) separately. It is, in fact, a linear function of invariants of (1), (2), (3), (4) of one whole weight and one whole degree, but different partial weights and degrees.

The invariant has then (§ 247) two annihilators which have been hitherto called  $\Sigma\Omega$ ,  $\Sigma O$ , but will for our present purpose be designated differently, viz.

$$\begin{aligned} \Omega_{yx} \equiv & \left( a_0 \frac{d}{db_0} + 2b_0 \frac{d}{dc_0} + 3c_0 \frac{d}{dd_0} \right) \\ & + \left( a_1 \frac{d}{db_1} + 2b_1 \frac{d}{dc_1} \right) + a_2 \frac{d}{db_2}, \end{aligned}$$

$$\begin{aligned} \Omega_{xy} \equiv & \left( 3b_0 \frac{d}{da_0} + 2c_0 \frac{d}{db_0} + d_0 \frac{d}{dc_0} \right) \\ & + \left( 2b_1 \frac{d}{da_1} + c_1 \frac{d}{db_1} \right) + b_2 \frac{d}{da_2}. \end{aligned}$$

In the general notation of the  $p$ -ic we should have

$$\Omega_{yx} \equiv \sum_{r=0}^{r=p-1} \left\{ a_r \frac{d}{db_r} + 2b_r \frac{d}{dc_r} + 3c_r \frac{d}{dd_r} + \dots \right\},$$

$$\begin{aligned} \Omega_{xy} \equiv & \sum_{r=0}^{r=p-1} \\ & \left\{ (p-r)b_r \frac{d}{da_r} + (p-r-1)c_r \frac{d}{db_r} + (p-r-2)d_r \frac{d}{dc_r} + \dots \right\}, \end{aligned}$$

where only coefficients which actually occur in the  $p$ -ic are present.

282.] In like manner a covariant of the ternary cubic is a rational integral function, of constant whole order degree and weight throughout, of covariants and invariants of the system (1), (2), (3), (4) of § 281. And analogously for the ternary  $p$ -ic. A covariant has then (§ 247) the two annihilators

$$\Omega_{yz} - y \frac{d}{dx},$$

$$\Omega_{xy} - x \frac{d}{dy},$$

and these, it is to be noticed, annihilate the ternary  $p$ -ic itself, which is of course to be regarded as one of its own covariants.

This fact has led to the frequent use for the operators  $\Omega_{yz}$ ,  $\Omega_{xy}$  of the symbolical notation  $\left[ y \frac{d}{dx} \right]$ ,  $\left[ x \frac{d}{dy} \right]$ .

283.] Let us now pay attention to the second and third forms (§ 280) in which the cubic or  $p$ -ic may be arranged. Attending to the second form we see, just as in §§ 281, 282, that an invariant, or covariant, is a rational integral function of invariants, or of covariants and invariants, of the system of  $p+1$  binary quantities, which for the case of the cubic are

$$a_0,$$

$$b_0 y + a_1 z,$$

$$c_0 y^2 + 2b_1 yz + a_2 z^2,$$

$$d_0 y^3 + 3c_1 y^2 z + 3b_2 yz^2 + a_3 z^3;$$

that an invariant has two annihilators, which for the case of the cubic are

$$\Omega_{xy} \equiv \left( d_0 \frac{d}{dc_1} + 2c_1 \frac{d}{db_2} + 3b_2 \frac{d}{da_3} \right) + \left( c_0 \frac{d}{db_1} + 2b_1 \frac{d}{da_2} \right) + b_0 \frac{d}{da_1},$$

$$\Omega_{yz} \equiv \left( 3c_1 \frac{d}{dd_0} + 2b_2 \frac{d}{dc_1} + a_3 \frac{d}{db_2} \right) + \left( 2b_1 \frac{d}{dc_0} + a_2 \frac{d}{db_1} \right) + a_1 \frac{d}{db_0};$$

and that a covariant has the annihilators

$$\Omega_{zy} - z \frac{d}{dy},$$

$$\Omega_{yz} - y \frac{d}{dz},$$

which in particular annihilate the cubic itself.

For the  $p$ -ic  $\Omega_{zy}$  and  $\Omega_{yz}$  are

$$\begin{aligned} \Omega_{zy} \equiv & \left\{ \dots + (p-1) c_{p-2} \frac{d}{db_{p-1}} + p b_{p-1} \frac{d}{da_p} \right\} \\ & + \left\{ \dots + (p-2) c_{p-3} \frac{d}{db_{p-2}} + (p-1) b_{p-2} \frac{d}{da_{p-1}} \right\} \\ & + \dots + b_0 \frac{d}{da_1}, \end{aligned}$$

$$\begin{aligned} \Omega_{yz} \equiv & \left\{ \dots + 2 b_{p-1} \frac{d}{dc_{p-2}} + a_p \frac{d}{db_{p-1}} \right\} \\ & + \left\{ \dots + 2 b_{p-2} \frac{d}{dc_{p-3}} + a_{p-1} \frac{d}{db_{p-2}} \right\} + \dots + a_1 \frac{d}{db_0}. \end{aligned}$$

In like manner, regarding the third form in § 280, we see that an invariant has two additional annihilators  $\Omega_{xz}$ ,  $\Omega_{zx}$ , and a covariant the two additional annihilators

$$\Omega_{xz} - x \frac{d}{dz}, \quad \Omega_{zx} - z \frac{d}{dx},$$

where, in the case of the cubic,

$$\Omega_{xz} \equiv \left( a_3 \frac{d}{da_2} + 2 a_2 \frac{d}{da_1} + 3 a_1 \frac{d}{da_0} \right) + \left( b_2 \frac{d}{db_1} + 2 b_1 \frac{d}{db_0} \right) + c_1 \frac{d}{dc_0},$$

$$\Omega_{zx} \equiv \left( 3 a_2 \frac{d}{da_3} + 2 a_1 \frac{d}{da_2} + a_0 \frac{d}{da_1} \right) + \left( 2 b_1 \frac{d}{db_2} + b_0 \frac{d}{db_1} \right) + c_0 \frac{d}{dc_1},$$

and, in the general case of the  $p$ -ic,

$$\begin{aligned} \Omega_{xz} \equiv & \left\{ a_p \frac{d}{da_{p-1}} + \dots + (p-1) a_2 \frac{d}{da_1} + p a_1 \frac{d}{da_0} \right\} \\ & + \left\{ b_{p-1} \frac{d}{db_{p-2}} + \dots + (p-2) b_2 \frac{d}{db_1} + (p-1) b_1 \frac{d}{db_0} \right\} + \dots, \end{aligned}$$

$$\begin{aligned} \Omega_{zx} \equiv & \left\{ p a_{p-1} \frac{d}{da_p} + \dots + 2 a_1 \frac{d}{da_2} + a_0 \frac{d}{da_1} \right\} \\ & + \left\{ (p-1) b_{p-2} \frac{d}{db_{p-1}} + \dots + 2 b_1 \frac{d}{db_2} + b_0 \frac{d}{db_1} \right\} + \dots \end{aligned}$$

284.] The facts as to invariants and covariants expressed by the existence of their six annihilators thus found are not all independent. Information as to the nature of their interdependence can be obtained by forming the fifteen alternants of  $\Omega_{yx}$ ,  $\Omega_{xy}$ ,  $\Omega_{zy}$ ,  $\Omega_{yz}$ ,  $\Omega_{zx}$ ,  $\Omega_{xz}$  in pairs. Taking the forms appropriate to the cubic, we readily obtain first the following triad of alternants ;

$$H_3 \equiv \Omega_{yx}\Omega_{xy} - \Omega_{xy}\Omega_{yz} \equiv \left( 3a_0 \frac{d}{da_0} + b_0 \frac{d}{db_0} - c_0 \frac{d}{dc_0} - 3d_0 \frac{d}{dd_0} \right) + \left( 2a_1 \frac{d}{da_1} - 2c_1 \frac{d}{dc_1} \right) + \left( a_2 \frac{d}{da_2} - b_2 \frac{d}{db_2} \right), \dots (3)$$

$$H_1 \equiv \Omega_{zy}\Omega_{yz} - \Omega_{yz}\Omega_{zy} \equiv \left( 3d_0 \frac{d}{dd_0} + c_1 \frac{d}{dc_1} - b_2 \frac{d}{db_2} - 3a_2 \frac{d}{da_2} \right) + \left( 2c_0 \frac{d}{dc_0} - 2a_2 \frac{d}{da_2} \right) + \left( b_0 \frac{d}{db_0} - a_1 \frac{d}{da_1} \right), \dots (1)$$

$$H_2 \equiv \Omega_{zx}\Omega_{xz} - \Omega_{xz}\Omega_{zx} \equiv \left( 3a_3 \frac{d}{da_3} + a_2 \frac{d}{da_2} - a_1 \frac{d}{da_1} - 3a_0 \frac{d}{da_0} \right) + \left( 2b_2 \frac{d}{db_2} - 2b_0 \frac{d}{db_0} \right) + \left( c_1 \frac{d}{dc_1} - c_0 \frac{d}{dc_0} \right). \dots (2)$$

We here see first that the sum  $H_1 + H_2 + H_3$  vanishes identically, whatever be the function operated upon. Thus any function which is annihilated by two of  $H_1$ ,  $H_2$ ,  $H_3$ , or by two independent sums of multiples of them, is also annihilated by the third, and by any sum of multiples of them.

Before forming the other alternants we proceed to exhibit the information as to invariants of the cubic given by these.

285.] Any invariant of the cubic is annihilated by  $H_1$ ,  $H_2$  and  $H_3$ . For every one of these is a difference of two parts each of which annihilates it, since it has the six annihilators  $\Omega$ . Let us consider the fact that  $H_1 - H_2$ , the difference of the operators written second and third above, annihilates it. We readily see that

$$H_1 - H_2 \equiv 3 \left( a_0 \frac{d}{da_0} + b_0 \frac{d}{db_0} + c_0 \frac{d}{dc_0} + d_0 \frac{d}{dd_0} \right) + a_1 \frac{d}{da_1} + b_1 \frac{d}{db_1} + c_1 \frac{d}{dc_1} + a_2 \frac{d}{da_2} + b_2 \frac{d}{db_2} + a_3 \frac{d}{da_3} - 3 \left( a_1 \frac{d}{da_1} + b_1 \frac{d}{db_1} + c_1 \frac{d}{dc_1} \right) - 6 \left( a_2 \frac{d}{da_2} + b_2 \frac{d}{db_2} \right) - 9 a_3 \frac{d}{da_3}.$$



We notice hence, by Euler's theorem as to homogeneous functions, and the consequent theorem (§ 117) as to isobaric functions, that  $H_1 - H_2$  operating on a function of degree  $i$  and weight (sum of suffixes)  $w$  has the effect of multiplying it by

$$3i - 3w.$$

Thus since the invariant, which is homogeneous (§ 22), is annihilated by  $H_1 - H_2$ , it must have a constant weight  $w$  throughout given by

$$3i - 3w = 0.$$

If we had taken the forms of  $H_1, H_2$  for the  $p$ -ic instead of the cubic, we should have had in like manner

$$pi - 3w = 0.$$

The information given by  $H_1 - H_2$  is then that of § 28.

In like manner we have, for the cubic,

$$\begin{aligned} H_2 - H_3 \equiv & 3 \left( a_0 \frac{d}{da_0} + b_0 \frac{d}{db_0} + c_0 \frac{d}{dc_0} + d_0 \frac{d}{dd_0} \right. \\ & \left. + a_1 \frac{d}{da_1} + b_1 \frac{d}{db_1} + c_1 \frac{d}{dc_1} + a_2 \frac{d}{da_2} + b_2 \frac{d}{db_2} + a_3 \frac{d}{da_3} \right) \\ & - 3 \left( c_0 \frac{d}{dc_0} + b_1 \frac{d}{db_1} + a_2 \frac{d}{da_2} \right) - 6 \left( b_0 \frac{d}{db_0} + a_1 \frac{d}{da_1} \right) - 9a_0 \frac{d}{da_0}, \end{aligned}$$

the annihilation of the invariant by which tells us, upon observation of the form (ii) of the cubic in § 280, that

$$3i - 3w' = 0,$$

where  $w'$  is the weight of the invariant when we consider  $\frac{y}{x}$  and  $\frac{z}{x}$  as of weight unity. Thus

$$w' = w,$$

which would also follow from the fact that the particular substitution which replaces  $x, y, z$  by  $y, z, x$ , whose modulus is unity, does not alter the value of the invariant while it replaces  $c_0, b_1, a_2, b_0, a_1, a_0$  by  $a_1, b_1, c_1, a_2, b_2, a_3$ .

For the general case of the  $p$ -ic we should have had

$$pi - 3w' = 0.$$

In exactly the same way

$$\begin{aligned}
 H_3 - H_1 \equiv & 3 \left( a_0 \frac{d}{da_0} + b_0 \frac{d}{db_0} + c_0 \frac{d}{dc_0} + d_0 \frac{d}{dd_0} \right. \\
 & \left. + a_1 \frac{d}{da_1} + b_1 \frac{d}{db_1} + c_1 \frac{d}{dc_1} + a_2 \frac{d}{da_2} + b_2 \frac{d}{db_2} + a_3 \frac{d}{da_3} \right) \\
 & - 3 \left( b_2 \frac{d}{db_2} + b_1 \frac{d}{db_1} + b_0 \frac{d}{db_0} \right) - 6 \left( c_1 \frac{d}{dc_1} + c_0 \frac{d}{dc_0} \right) - 9 d_0 \frac{d}{dd_0},
 \end{aligned}$$

the annihilation by which requires that an invariant of the cubic have the property

$$3i - 3w'' = 0,$$

where  $w''$  is the weight when each  $a$  is of weight 0, each  $b$  of weight 1, each  $c$  of weight 2, and  $d_0$  of weight 3, i.e. when  $\frac{z}{y}$  and  $\frac{x}{y}$  are regarded as of unit weight. Thus

$$w'' = w' = w.$$

For the  $p$ -ic we should have in like manner

$$pi - 3w'' = 0.$$

Since the sum of  $H_2 - H_3$ ,  $H_3 - H_1$  and  $H_1 - H_2$  vanishes, any function, not necessarily an invariant, which possesses two of these properties must also possess the third.

Ex. 1. From the fact of annihilation by  $H_3$ , which may be written

$$\begin{aligned}
 3 \left( a_0 \frac{d}{da_0} + b_0 \frac{d}{db_0} + c_0 \frac{d}{dc_0} + d_0 \frac{d}{dd_0} \right) &+ 2 \left( a_1 \frac{d}{da_1} + b_1 \frac{d}{db_1} + c_1 \frac{d}{dc_1} \right) \\
 &+ \left( a_2 \frac{d}{da_2} + b_2 \frac{d}{db_2} \right) \\
 - 2 \left\{ \left( b_0 \frac{d}{db_0} + 2c_0 \frac{d}{dc_0} + 3d_0 \frac{d}{dd_0} \right) \right. &+ \left. \left( b_1 \frac{d}{db_1} + 2c_1 \frac{d}{dc_1} \right) + b_2 \frac{d}{db_2} \right\},
 \end{aligned}$$

show that if any invariant of a cubic be written as a sum of parts, each separately homogeneous in the sets

$$a_0, b_0, c_0, d_0,$$

$$a_1, b_1, c_1,$$

$$a_2, b_2,$$

$$a_3,$$

and if  $i_3, i_2, i_1$  be the degrees of any such parts in the first, second, and third of these sets, then throughout the invariant

$$3i_3 + 2i_2 + i_1 = 2w.$$

For an invariant of the  $p$ -ic the corresponding fact is

$$pi_p + (p-1)i_{p-1} + \dots + 2i_2 + i_1 = 2w.$$

Ex. 2. From the fact of annihilation by  $H_1$  show that throughout an invariant of the cubic

$$3i_3' + 2i_2' + i_1' = 2w,$$

where  $i_3', i_2', i_1'$  are the partial degrees of any term in the sets

$$d_0, c_1, b_2, a_3,$$

$$c_0, b_1, a_2,$$

$$b_0, a_1;$$

and state the corresponding fact for the  $p$ -ic.

Ex. 3. From the fact of annihilation by  $H_2$  show that throughout an invariant of the cubic

$$3i_3'' + 2i_2'' + i_1'' = 2w,$$

where  $i_3'', i_2'', i_1''$  are the partial degrees of any term in

$$a_3, a_2, a_1, a_0,$$

$$b_2, b_1, b_0,$$

$$c_1, c_0;$$

and state the corresponding fact for the  $p$ -ic.

286.] We now form the other alternants of the six  $\Omega$ 's. They occur in cyclical sets of three. Taking the case of the cubic, and referring to §§ 281, 283 for the notation, we find

$$\begin{aligned} \Omega_{yx}\Omega_{zx} - \Omega_{zx}\Omega_{yx} &\equiv a_0 \frac{d}{db_1} + 2b_0 \frac{d}{dc_1} + 2a_1 \frac{d}{db_2} \\ &\quad - 2a_1 \frac{d}{db_2} - a_0 \frac{d}{db_1} - 2b_0 \frac{d}{dc_1} \\ &= 0, \end{aligned} \quad \dots (4)$$

$$\Omega_{zy}\Omega_{xy} - \Omega_{xy}\Omega_{zy} = 0, \quad \dots (5)$$

$$\Omega_{xz}\Omega_{yz} - \Omega_{yz}\Omega_{xz} = 0. \quad \dots (6)$$

The same relations hold in the general notation of the  $p$ -ic.

In like manner we have another triad

$$\Omega_{xy}\Omega_{xz} - \Omega_{zx}\Omega_{xy} = 0, \quad \dots (7)$$

$$\Omega_{yz}\Omega_{yx} - \Omega_{yx}\Omega_{yz} = 0, \quad \dots (8)$$

$$\Omega_{zx}\Omega_{zy} - \Omega_{zy}\Omega_{zx} = 0. \quad \dots (9)$$

Next we have, for the cubic,

$$\begin{aligned} \Omega_{xy}\Omega_{zx} - \Omega_{zx}\Omega_{xy} &\equiv 3b_0 \frac{d}{da_1} + 2c_0 \frac{d}{db_1} + d_0 \frac{d}{dc_1} + 4b_1 \frac{d}{da_2} + 2c_1 \frac{d}{db_2} \\ &\quad + 3b_2 \frac{d}{da_3} - 2b_1 \frac{d}{da_2} - 2b_0 \frac{d}{da_1} - c_0 \frac{d}{db_1} \\ &\equiv d_0 \frac{d}{dc_1} + 2c_1 \frac{d}{db_2} + 3b_2 \frac{d}{da_3} \\ &\quad + c_0 \frac{d}{db_1} + 2b_1 \frac{d}{da_2} + b_0 \frac{d}{da_1} \\ &\equiv \Omega_{zy}, \quad \dots (10) \end{aligned}$$

and similarly

$$\Omega_{yz}\Omega_{xy} - \Omega_{xy}\Omega_{yz} \equiv \Omega_{zx}, \quad \dots (11)$$

$$\Omega_{zx}\Omega_{yz} - \Omega_{yz}\Omega_{zx} \equiv \Omega_{yx}. \quad \dots (12)$$

Lastly we have in like manner

$$\Omega_{yx}\Omega_{zx} - \Omega_{zx}\Omega_{yx} \equiv -\Omega_{yz}, \quad \dots (13)$$

$$\Omega_{xy}\Omega_{yx} - \Omega_{yx}\Omega_{xy} \equiv -\Omega_{zx}, \quad \dots (14)$$

$$\Omega_{zx}\Omega_{zy} - \Omega_{zy}\Omega_{zx} \equiv -\Omega_{xy}. \quad \dots (15)$$

All these apply to the general notation of the  $p$ -ic, as well as to that of the cubic.

287.] The fifteen alternants of pairs of  $\Omega$ 's introduce then no new operators except the  $H_1, H_2, H_3$  of § 284. We complete the theory of the annihilators by forming the alternants of these three with one another and the  $\Omega$ 's. It is quite easy to see that

$$H_2H_3 - H_3H_2 = 0,$$

$$H_3H_1 - H_1H_3 = 0,$$

$$H_1H_2 - H_2H_1 = 0;$$

and that

$$\begin{aligned} H_3 \Omega_{yx} - \Omega_{yx} H_3 &\equiv 2 \Omega_{yx}, \\ H_3 \Omega_{xy} - \Omega_{xy} H_3 &\equiv -2 \Omega_{xy}, \\ H_3 \Omega_{zy} - \Omega_{zy} H_3 &\equiv -\Omega_{zy}, \\ H_3 \Omega_{yz} - \Omega_{yz} H_3 &\equiv \Omega_{yz}, \\ H_3 \Omega_{zx} - \Omega_{zx} H_3 &\equiv -\Omega_{zx}, \\ H_3 \Omega_{xz} - \Omega_{xz} H_3 &\equiv \Omega_{xz}, \end{aligned}$$

together with two other sets of six, in the first set of which  $H_1$  occurs and the suffixes  $x, y, z$  are interchanged once cyclically, and in the other set  $H_2$  occurs and a second cyclical interchange is made in the suffixes.

Accordingly the nine operators  $\Omega_{yx}, \Omega_{zy}, \Omega_{zx}, \Omega_{xy}, \Omega_{yz}, \Omega_{xz}, H_1, H_2, H_3$  form a group such that, when we form the alternant of any pair of them, some member of the group with a simple numerical multiplier, or else a zero, is produced.

288.] **Three cyclical annihilators suffice to define invariants.** We can now see that, if a function of the coefficients have a cyclical set of three annihilators, such as  $\Omega_{yx}, \Omega_{zx}, \Omega_{xy}$  or  $\Omega_{zy}, \Omega_{xz}, \Omega_{yx}$ , it has also the other three, and is accordingly, if homogeneous, an invariant.

Suppose, for instance, that  $\Omega_{yx}, \Omega_{zx}, \Omega_{xy}$  annihilate a function. By § 286 (10), since  $\Omega_{zx}$  and  $\Omega_{xy}$  annihilate it, so does  $\Omega_{zy}$ . By (11), since  $\Omega_{xy}$  and  $\Omega_{yz}$  annihilate it, so does  $\Omega_{xz}$ . And by (12), since  $\Omega_{yz}$  and  $\Omega_{zx}$  annihilate it, so does  $\Omega_{yx}$ .

And again, to repeat from § 285, since  $\Omega_{yz}$  and  $\Omega_{zy}$  annihilate it, so does  $H_1$ ; and in like manner so do  $H_2$  and  $H_3$ . Thus the function has necessarily the degree and weight properties expressed in § 285 and the examples which follow that article.

The property of annihilation by  $\Omega_{yz}, \Omega_{zx}$  and  $\Omega_{xy}$  includes then all the facts with regard to invariants of ternary quantities except that of homogeneity, just as that of annihilation by  $\Omega$  and  $O$ , i. e. by  $\Omega_{yx}$  and  $\Omega_{xy}$ , does with regard to invariants of binary quantities.

We shall see later by another method, which might have been here applied, that the property of having  $q$  cyclical annihilators of the  $\Omega$  form includes all the facts but that of homogeneity as to invariants of  $q$ -ary quantities.

289.] The six annihilators, as well as going in two triads, go in three pairs  $\Omega_{yx}, \Omega_{xy}; \Omega_{zy}, \Omega_{yz}; \Omega_{zx}, \Omega_{xz}$ . It is sometimes most convenient to use the fact that if two pairs of these annihilate a function its annihilation by the third pair is necessitated. For instance, if the first four annihilate it, it follows from § 286 (14) that  $\Omega_{zx}$  annihilates it, and from (11) that  $\Omega_{xz}$  does, so that it is an invariant.

The possession of *three* annihilators *not* forming a cyclical set does not suffice.

290.] **Invariant of the ternary quadratic.** Let us exemplify some of the above principles by deciding what invariants the ternary quadratic

$$a_2 z^2 + 2(a_1 x + b_1 y)z + a_0 x^2 + 2b_0 xy + c_0 y^2$$

can possess.

Since  $\Omega_{yx}$  and  $\Omega_{xy}$  annihilate it, an invariant of the quadratic is an invariant of the system

$$a_2, \quad a_1 x + b_1 y, \quad a_0 x^2 + 2b_0 xy + c_0 y^2,$$

and so (§ 251) is a rational integral function of

$$a_2, \quad a_0 c_0 - b_0^2, \quad a_0 b_1^2 - 2b_0 a_1 b_1 + c_0 a_1^2. \quad \dots (1)$$

Again, since  $\Omega_{zx}$  and  $\Omega_{xz}$  annihilate it, it is an invariant of the system

$$c_0, \quad b_1 z + b_0 x, \quad a_2 z^2 + 2a_1 zx + a_0 x^2,$$

and so is a rational integral function of

$$c_0, \quad a_0 a_2 - a_1^2, \quad a_2 b_0^2 - 2a_1 b_1 b_0 + a_0 b_1^2. \quad \dots (2)$$

Consider the first fact, and let

$$a_2^\lambda (a_0 c_0 - b_0^2)^\mu (a_0 b_1^2 - 2b_0 a_1 b_1 + c_0 a_1^2)^\nu$$

be a part of the invariant. By § 285 its weight measured by sum of suffixes must be equal to its weight considering  $a$ 's,  $b$ 's, and  $c_0$  as respectively of weights 0, 1, 2. Thus

$$2\lambda + 2\nu = 2\mu + 2\nu,$$

so that  $\lambda = \mu$ , and the invariant involves only

$$a_2 (a_0 c_0 - b_0^2), \text{ and } a_0 b_1^2 - 2b_0 a_1 b_1 + c_0 a_1^2.$$

Similarly it involves only

$$c_0 (a_0 a_2 - a_1^2), \text{ and } a_2 b_0^2 - 2a_1 b_1 b_0 + a_0 b_1^2.$$

Now any function of the first of these pairs which is also a function of the second pair must in particular be so when  $c_0 = 0$ . Thus a necessity as to such a function is that the said function of

$$-a_2 b_0^2 \text{ and } a_0 b_1^2 - 2b_0 a_1 b_1$$

is a function of

$$0 \text{ and } a_2 b_0^2 - 2a_1 b_1 b_0 + a_0 b_1^2.$$

The difference of the two, and its powers, are obviously the only functions for which this is the case.

Thus

$$a_2(a_0 c_0 - b_0^2) - (a_0 b_1^2 - 2b_0 a_1 b_1 + c_0 a_1^2)$$

and its powers are the only functions which can be invariants, and they can be so only if this is also a function of  $c_0(a_0 a_2 - a_1^2)$  and  $a_2 b_0^2 - 2a_1 b_1 b_0 + a_0 b_1^2$ , as it is, viz. the difference of the two.

The ternary quadratic has then only one irreducible invariant

$$a_0 c_0 a_2 + 2b_1 a_1 b_0 - a_0 b_1^2 - c_0 a_1^2 - a_2 b_0^2,$$

which is, in the more usual notation,

$$abc + 2fgh - af^2 - bg^2 - ch^2,$$

the discriminant.

291.] **The ternary cubic.** The general ternary cubic can (§ 229) be linearly transformed into  $X^3 + Y^3 + Z^3 + 6mXYZ$ .

It cannot then have more than two independent invariants. For if it had three it would have two absolute invariants, i.e. there would be two functions of the coefficients equal to functions of  $m$ ; and by elimination of  $m$  we could find a relation in the coefficients only, which there cannot be as the coefficients are all independent.

Two independent invariants  $S$  and  $T$ , of degrees 4 and 6, will now be found. It will hereafter be seen that not only is there no other independent of these, but that there is no other which cannot be rationally and integrally expressed in terms of them, so that they form the whole system of irreducible invariants.

By § 285, or by § 28, the weight, in either of the three senses, of an invariant of the ternary cubic is equal to its degree. Thus  $S$ , which we seek, is of degree 4 and weight 4.

Now suppose that  $S$  contains a term or terms of degrees

$m$  in  $a_3$ ,  $n$  in  $a_2$  and  $b_2$ ,  $p$  in  $a_1$ ,  $b_1$ ,  $c_1$ , and  $q$  in  $a_0$ ,  $b_0$ ,  $c_0$ ,  $d_0$ . The facts as to degree and weight give us

$$m + n + p + q = 4,$$

$$3m + 2n + p = 4,$$

and the only positive integral, including zero, values of  $m$ ,  $n$ ,  $p$ ,  $q$  which satisfy these equations are given by the scheme

$$\underline{m, n, p, q}.$$

$$1, 0, 1, 2$$

$$0, 2, 0, 2$$

$$0, 1, 2, 1$$

$$0, 0, 4, 0$$

Thus  $S$ , if it exists, must be of the form

$$a_3(1^1 0^2) + (2^2 0^2) + (2^1 1^2 0^1) + (1^4),$$

where, for instance, the notation  $(2^1 1^2 0^1)$  denotes a function of degree 1 in  $a_2$ ,  $b_2$ , degree 2 in  $a_1$ ,  $b_1$ ,  $c_1$ , and degree 1 in  $a_0$ ,  $b_0$ ,  $c_0$ ,  $d_0$ .

Moreover, since  $\Omega_{yx}S = 0$  and  $\Omega_{xy}S = 0$ , the functions  $(1^1 0^2)$ ,  $(2^2 0^2)$ ,  $(2^1 1^2 0^1)$ ,  $(1^4)$  are invariants of the system

$$a_2x + b_2y,$$

$$a_1x^2 + 2b_1xy + c_1y^2,$$

$$a_0x^3 + 3b_0x^2y + 3c_0xy^2 + d_0y^3.$$

Now the invariants of this system are (§ 262) the invariants of the quadratic and cubic given in § 260, and the results of replacing  $x$  and  $y$  by  $b_2$  and  $-a_2$  in the covariants of that article. Those of degree and sum of suffixes not exceeding 4 are given by (2), (4), (6), (9), (10), (13) of the article in question, and are

$$A = a_1c_1 - b_1^2,$$

$$B = (a_0c_0 - b_0^2)b_2^2 - (a_0d_0 - b_0c_0)a_2b_2 + (b_0d_0 - c_0^2)a_2^2,$$

$$C = (a_0d_0 - b_0c_0)^2 - 4(a_0c_0 - b_0^2)(b_0d_0 - c_0^2),$$

$$D = (a_1c_0 - 2b_1b_0 + c_1a_0)b_2 - (a_1d_0 - 2b_1c_0 + c_1b_0)a_2,$$

$$E = \{a_1^2d_0 - 3a_1b_1c_0 + (a_1c_1 + 2b_1^2)b_0 - b_1c_1a_0\}b_2 \\ + \{c_1^2a_0 - 3b_1c_1b_0 + (a_1c_1 + 2b_1^2)c_0 - a_1b_1d_0\}a_2,$$

$$F = a_1(b_0d_0 - c_0^2) - b_1(a_0d_0 - b_0c_0) + c_1(a_0c_0 - b_0^2).$$



We see then that we must have

$$\begin{aligned}(1^1 0^2) &= \lambda F, \\ (2^2 0^2) &= \mu B, \\ (2^1 1^2 0^1) &= \nu E, \\ (1^4) &= \varpi A^2,\end{aligned}$$

where  $\lambda, \mu, \nu, \varpi$  are numerical.

Thus we must have

$$\begin{aligned}S &= \lambda a_3 \{a_1 (b_0 d_0 - c_0^2) - b_1 (a_0 d_0 - b_0 c_0) + c_1 (a_0 c_0 - b_0^2)\} \\ &+ \mu \{(a_0 c_0 - b_0^2) b_2^2 - (a_0 d_0 - b_0 c_0) a_2 b_2 + (b_0 d_0 - c_0^2) a_2^2\} \\ &+ \nu \{a_1^2 d_0 b_2 - a_1 b_1 (3 c_0 b_2 + d_0 a_2) + (a_1 c_1 + 2 b_1^2) (b_0 b_2 + c_0 a_2) \\ &\quad - b_1 c_1 (a_0 b_2 + 3 b_0 a_2) + c_1^2 a_0 a_2\} \\ &+ \varpi (a_1 c_1 - b_1^2)^2.\end{aligned}$$

To determine  $\lambda, \mu, \nu, \varpi$  we may express that  $S$  is annihilated by any of the  $\Omega$ 's, except  $\Omega_{yx}$  and  $\Omega_{xy}$  by which we have already secured its annihilation, whatever  $\lambda, \mu, \nu, \varpi$  be. More easily perhaps we may use the fact that  $S$  must be exactly the same function of

$$\begin{aligned}d_0, \\ c_0, c_1, \\ b_0, b_1, b_2, \\ a_0, a_1, a_2, a_3,\end{aligned}$$

as of

$$\begin{aligned}a_3, \\ a_2, b_2, \\ a_1, b_1, c_1, \\ a_0, b_0, c_0, d_0,\end{aligned}$$

since these sets of coefficients are exactly interchanged by the linear substitution which interchanges  $y$  and  $z$  leaving  $x$  unaltered, whose modulus is  $-1$ , which modulus in the expression of invariancy of an invariant of weight 4 is raised to the fourth power, thus producing  $+1$  for the factor. For  $S$  to be an invariant the above expression must then be the same as

$$\begin{aligned}\lambda d_0 \{b_0 (a_1 a_3 - a_2^2) - b_1 (a_0 a_3 - a_1 a_2) + b_2 (a_0 a_2 - a_1^2)\} \\ + \mu \{(a_0 a_2 - a_1^2) c_1^2 - (a_0 a_3 - a_1 a_2) c_0 c_1 + (a_1 a_3 - a_2^2) c_0^2\} \\ + \nu \{b_0^2 a_3 c_1 - b_0 b_1 (3 a_2 c_1 + a_3 c_0) + (b_0 b_2 + 2 b_1^2) (a_1 c_1 + a_2 c_0) \\ \quad - b_1 b_2 (a_0 c_1 + 3 a_1 c_0) + b_2^2 a_0 c_0\} \\ + \varpi (b_0 b_2 - b_1^2)^2.\end{aligned}$$

We at once see that the two forms are identical if

$$\frac{\lambda}{-1} = \frac{\mu}{1} = \frac{\nu}{1} = \frac{\varpi}{-1}.$$

With these values the expression will have  $\Omega_{xx}$  and  $\Omega_{xz}$  for annihilators, because of its second form, as well as  $\Omega_{yz}$  and  $\Omega_{xy}$  because of its first form. It is then an invariant by § 289. Thus the invariant  $S$  looked for is, adopting the second form of writing it,

$$\begin{aligned} S = & (a_0 a_2 - a_1^2) c_1^2 - (a_0 a_3 - a_1 a_2) c_0 c_1 + (a_1 a_3 - a_2^2) c_0^2 \\ & + b_0^2 a_3 c_1 - b_0 b_1 (3 a_2 c_1 + a_3 c_0) + (b_0 b_2 + 2 b_1^2) (a_1 c_1 + a_2 c_0) \\ & \quad - b_1 b_2 (a_0 c_1 + 3 a_1 c_0) + b_2^2 a_0 c_0 \\ & - d_0 \{ b_0 (a_1 a_3 - a_2^2) - b_1 (a_0 a_3 - a_1 a_2) + b_2 (a_0 a_2 - a_1^2) \} \\ & - (b_0 b_2 - b_1^2)^2. \end{aligned}$$

292.] The cubic can (§ 229) be linearly transformed to the canonical form  $X^3 + Y^3 + Z^3 + 6mXYZ$ .

By a substitution of modulus unity it can consequently be given the form

$$a'(x^3 + y^3 + z^3) + 6m'xyz.$$

Let us consider it in the less particularized form

$$a_0'x^3 + d_0'y^3 + a_3'z^3 + 6b_1'xyz,$$

in which the names of non-vanishing coefficients accord with the notation used in general. The modulus of the transformation which produces this from the general cubic is taken to be unity.

For this form we have, by the above,

$$S = a_0' d_0' a_3' b_1' - b_1'^4 = b_1' (a_0' d_0' a_3' - b_1'^3).$$

For the canonical form itself the value is

$$m(1 - m^3),$$

which is of course equal not to the  $S$  of the untransformed cubic but to that  $S$  multiplied by the fourth power of the modulus, which is no longer unity.

Ex. 4. Show that  $S = 0$  is the condition that the cubic be capable of expression as a sum of three cubes.

293.] The second invariant  $T$  of degree 6, and therefore, since  $3i = 3w$ , also of weight 6, can be found as  $S$  has been.

For our purposes, however, the labour may be avoided by use of a covariant, the Hessian, which it is already known from § 11 that the cubic possesses.

For the semi-canonized form

$$a_0'x^3 + d_0'y^3 + a_3'z^3 + 6b_1'xyz,$$

the Hessian, with the numerical factor  $6^3$  rejected, is

$$\begin{vmatrix} a_0'x & b_1'z & b_1'y \\ b_1'z & d_0'y & b_1'x \\ b_1'y & b_1'x & a_3'z \end{vmatrix},$$

i. e.  $(a_0'd_0'a_3' + 2b_1'^3)xyz - b_1'^2(a_0'x^3 + d_0'y^3 + a_3'z^3).$

The Hessian is then a covariant of the third degree and order.

For the canonical form itself the Hessian is

$$(1 + 2m^3)XYZ - m^2(X^3 + Y^3 + Z^3),$$

i. e. this is equal to the Hessian of the untransformed quantic multiplied by the square of the modulus of the fully canonizing substitution.

Now an invariant of a quantic and a covariant is an invariant of the quantic alone. Also (§ 19) if  $ax^3 + \dots$  and  $Ax^3 + \dots$  are two quantics of the same order, we may derive an invariant of the two from one of the first only by operation

with  $A \frac{d}{da} + \dots$ . Applying this principle to the cubic and its

Hessian, both of order 3, we derive from the invariant  $S$  another invariant of the sixth degree. This is  $T$ , or rather a numerical multiple of  $T$ .

We can at once derive the expression for  $T$  that goes with the semi-canonized form

$$a_0'x^3 + d_0'y^3 + a_3'z^3 + 6b_1'xyz,$$

whose Hessian is as above

$$-b_1'^2a_0'x^3 - b_1'^2d_0'y^3 - b_1'^2a_3'z^3 + (a_0'd_0'a_3' + 2b_1'^3)xyz,$$

by operating with

$$-b_1'^2a_0' \frac{d}{da_0'} - b_1'^2d_0' \frac{d}{dd_0'} - b_1'^2a_3' \frac{d}{da_3'} + \frac{1}{6}(a_0'd_0'a_3' + 2b_1'^3) \frac{d}{db_1'}$$

on  $S$ , which is

$$b_1'(a_0'd_0'a_3' - b_1'^3).$$

The reason for this lawfulness of working with the reduced number of coefficients is that the full expression for  $S$  contains, besides the terms  $b_1(a_0d_0a_3 - b_1^3)$ , only terms which involve coefficients which vanish for the semi-canonized form—indeed only powers and products of such coefficients, a fact which will be useful later—and that the terms in the generating operator other than those in  $\frac{d}{da_0}$ ,  $\frac{d}{dd_0}$ ,  $\frac{d}{da_3}$ ,  $\frac{d}{db_1}$  have for coefficients coefficients in the Hessian which vanish for the semi-canonized form.

The result of performing the operation above, and multiplying by 6, is to obtain

$$T = (a_0'd_0'a_3')^2 - 20b_1'^3(a_0'd_0'a_3') - 8b_1'^6.$$

For the canonical form itself this becomes

$$1 - 20m^3 - 8m^6;$$

but this is equal, not to the  $T$  of the untransformed cubic, but to the  $T$  multiplied by the sixth power of the canonizing modulus.

There is no difficulty in obtaining the lengthy expression for  $T$  in the notation of the general cubic, but only tediousness. It will not be here written down. Reference may be made for it to Salmon's Higher Plane Curves, §§ 221, 223.

Ex. 5. Prove that  $b_1'$  is given by the quartic in  $b_1'^2 = \beta$

$$27\beta^4 + 18S\beta^2 + T\beta - S^2 = 0,$$

and that when  $b_1'$  is found the corresponding product  $a_0'd_0'a_3'$  is uniquely determined.

Ex. 6. The discriminant of this quartic is a perfect square, namely, a numerical multiple of the square of  $64S^3 + T^2$ , which is a numerical multiple of its catalecticant.

Ex. 7. The discriminant of the ternary cubic is

$$a_0'd_0'a_3'(a_0'd_0'a_3' + 8b_1'^3)^3, \text{ i.e. } 64S^3 + T^2.$$

Ex. 8. The  $S$  of the Hessian of a ternary cubic is a numerical multiple of  $48S^3 + T^2$ .

294.] The result of Ex. 5 above leads us to expect that there are eight distinct ways of reducing the cubic to the

form  $ax^3 + by^3 + cz^3 + 6mxyz$  by substitutions of such modulus that *S* and *T* are absolutely unaltered. (Note that we do not reckon as distinct different ways in which the product  $xyz$  is the same.) If *M* be the modulus of such a substitution, the facts with regard to *S* and *T* give us respectively  $M^4 = 1$  and  $M^6 = 1$ . These lead to  $M^2 = 1$ , i.e.  $M = \pm 1$ .

It is easy to see that there really are eight ways, and to exhibit their connexion. Take, for instance, the cubic

$$x^3 + y^3 + z^3 + 6mxyz.$$

The substitutions of modulus 1

$$x\sqrt{-3} = X + Y + Z, \quad y\sqrt{-3} = X + \omega Y + \omega^2 Z, \\ z\sqrt{-3} = X + \omega^2 Y + \omega Z,$$

$$x\sqrt{-3} = X + \omega Y + \omega Z, \quad y\sqrt{-3} = X + \omega^2 Y + Z, \\ z\sqrt{-3} = X + Y + \omega^2 Z,$$

$$x\sqrt{-3} = X + \omega^2 Y + \omega^2 Z, \quad y\sqrt{-3} = X + Y + \omega Z, \\ z\sqrt{-3} = X + \omega Y + Z,$$

produce three forms whose *m*'s are respectively

$$m_2 = -\frac{1-m}{\sqrt{-3}}, \quad m_3 = -\frac{\omega^2-m}{\sqrt{-3}}, \quad m_4 = -\frac{\omega-m}{\sqrt{-3}},$$

so that

$$mm_2m_3m_4 = \frac{m(1-m^3)}{3\sqrt{-3}} = \frac{S}{3\sqrt{-3}}.$$

Also  $x = x', y = y', z = -z'$ , whose modulus is  $-1$ , gives a form with  $-m$  for  $m$ , those with  $-m_2, -m_3, -m_4$  for  $m$  being obtained in like manner. The product of the eight *m*'s is then

$$(mm_2m_3m_4)^2 = -\frac{1}{27}S^2,$$

which accords with § 293, Ex. 5.

295.] *S* and *T* the only irreducible invariants. We may prove as follows that any other invariant of the ternary cubic is a rational integral function of *S* and *T*.

Write the semi-canonical form of the cubic with the notation of coefficients

$$ax^3 + by^3 + cz^3 + 6mxyz, \quad \dots (1)$$

so that

$$S = m(abc - m^3),$$

$$T = (abc)^2 - 20m^3abc - 8m^6,$$

and (§ 293, Ex. 5)

$$27m^8 + 18Sm^4 + Tm^2 - S^2 = 0. \quad \dots (2)$$

We notice here, and from the last article, that the product of the eight values of  $m$  for substitutions which leave  $S$  and  $T$  unaltered, modulus  $+1$  or  $-1$ , is a numerical multiple of  $S^2$ . The product of the four for modulus  $+1$  is a numerical multiple of  $S$ .

Now if an invariant vanish when an  $m$  vanishes it must when any of the  $m$ 's vanishes. For its form for (1) has  $m$  for a factor, and it is the same thing,  $a, b, c$  being really independent, that its form for

$$a_2x^3 + b_2y^3 + c_2z^3 + 6m_2xyz$$

have  $m_2$  for a factor. An invariant divisible by  $m$  is then divisible by  $mm_2m_3m_4$ , i.e. by  $S$ ; and the quotient as well as itself must be an invariant. Equally if divisible by  $abc - m^3$ , a product of three  $m$ 's, an invariant is divisible by the fourth, and so by  $S$ .

We have then only to consider the reducibility of invariants which are not divisible by  $m$  or by  $abc - m^3$ , which latter call  $k$ . In this notation we have

$$S = mk, \quad \dots (3)$$

$$\begin{aligned} T &= k^2 - 18m^3k - 27m^6 \\ &= k^2 - 18Sm^2 - 27m^6. \end{aligned} \quad \dots (4)$$

Consider then an invariant  $I$  which is not divisible by  $m$  or by  $k$ . Its degree must be a multiple of 3. For  $k$  is of degree 3, so that, if its term free from  $m$  is  $k^n$ , this is of degree  $3n$ , and this degree must be preserved throughout. We may suppose then that the invariant is

$$I = k^n + pk^{n-1}m^3 + qk^{n-2}m^6 + \dots + tm^{3n},$$

for it is a function of  $S$  and  $T$  (§ 291), and consequently of the independent  $k$  and  $m$ , which are all that  $S$  and  $T$  involve. It would not be integral were  $k$  or  $m$  involved fractionally.

Now first use (4) to depress  $I$  to the first degree in  $k$ , or to the degree zero in  $k$  if no odd powers of  $k$  occur in  $I$  as already

exhibited, by substitution for  $k^2$  of  $T + 18Sm^2 + 27m^6$ . We thus get

$$I = kf(S, T, m) + \phi(S, T, m),$$

where the functions  $f$  and  $\phi$  are rational and integral, and where the former may or may not actually occur.

Firstly, if  $f(S, T, m)$  do not occur, we have

$$I = \phi(S, T, m).$$

By aid of (2) we may depress this equation below the eighth degree in  $m$  by successive substitutions such as that of  $\frac{1}{27}(S^2m^r - Tm^{r+2} - 18Sm^{r+4})$  for  $m^{8+r}$ , where  $r$  is zero or a positive integer. We thus get eventually

$$I = Am^7 + Bm^6 + \dots + K,$$

where  $A, B, \dots K$ , if they do not vanish, are rational integral functions of  $S$  and  $T$ .

Now there are (§ 294) eight values of  $m$  which must satisfy this equation of degree 7 at most. It must then be an identity. Hence, taking the terms free from  $m$ ,

$$I = K,$$

i.e.  $I$  is a rational integral function of  $S$  and  $T$ .

Secondly, if  $f(S, T, m)$  do occur, let  $\frac{1}{m}S$  be put for  $k$ , by (3), in  $kf(S, T, m)$ . We have

$$I = \frac{S}{m}F(S, T) + \psi(S, T, m),$$

where  $F$  and  $\psi$  are rational and integral; and this again may be reduced by (2) to

$$Im = A'm^7 + B'm^6 + \dots + H'm + K',$$

where  $A', B', \dots H', K'$ , if non-vanishing, are rational integral functions of  $S$  and  $T$ . This equation of degree 7 must be an identity by reasoning as before, and therefore, taking coefficients of  $m$ ,

$$I = H',$$

so that the conclusion as before is that  $I$  is a rational integral function of  $S$  and  $T$ .





covariant is annihilated by  $\Omega_{yx}$  and  $\Omega_{xy}$ . By the same reasoning the coefficient of  $x^w$  is annihilated by  $\Omega_{zy}$  and  $\Omega_{yz}$ .

Thus the coefficient of  $x^w$  in the covariant has the four annihilators

$$\Omega_{yx}, \Omega_{zx}, \Omega_{zy}, \Omega_{yz}.$$

In like manner the coefficient of  $y^w$  has the four annihilators

$$\Omega_{zy}, \Omega_{xy}, \Omega_{zx}, \Omega_{xz};$$

and the coefficient of  $z^w$  has the four annihilators

$$\Omega_{xz}, \Omega_{yz}, \Omega_{yx}, \Omega_{xy}.$$

§97.] A covariant is given by an end coefficient. When one of these three coefficients is known the whole covariant is known.

Consider the arrangement, (1) of the preceding article, by powers of  $z$ . Expressing the fact of annihilation by  $\Omega_{xz} - x \frac{d}{dz}$ , we have, by taking the coefficients of the successive powers of  $z$ ,

$$\Omega_{xz}P_w - wxP_{w-1} = 0,$$

$$\Omega_{xz}P_{w-1} - (w-1)xP_{w-2} = 0,$$

. . . . .

$$\Omega_{xz}P_1 - xP_0 = 0,$$

$$\Omega_{xz}P_0 = 0,$$

which tell us that (1) may be written

$$\left\{ 1 + \frac{z}{x} \Omega_{xz} + \frac{1}{1 \cdot 2} \frac{z^2}{x^2} \Omega_{xz}^2 + \dots + \frac{1}{w!} \frac{z^w}{x^w} \Omega_{xz}^w + \dots \right\} P_w,$$

the terms beyond the last written down vanishing because  $\Omega_{xz}P_0 = 0$ , i.e.  $\Omega_{xz}^{w+1}P_w = 0$ , and consequently  $\Omega_{xz}^{w+r}P_w = 0$ , for any positive value of the integer  $r$ .

Now this expression for the covariant may be written

$$e^{\frac{z}{x} \Omega_{xz}} P_w.$$

Again consider  $P_w$ . It is by the last article annihilated by  $\Omega_{yx} - y \frac{d}{dx}$  and  $\Omega_{xy} - x \frac{d}{dy}$ . Hence as above, or as in § 110, if  $S$  be the coefficient of  $x^w$  in  $P_w$ , i.e. in the covariant,

$$P_w = x^w e^{\frac{y}{x} \Omega_{xy}} S.$$

Consequently, upon insertion of this value for  $P_{\omega}$ , the covariant is

$$x^{\omega} e^{\frac{z}{x}\Omega_{xz}} e^{\frac{y}{x}\Omega_{xy}} S,$$

which, since (§ 286)  $\Omega_{xz}\Omega_{xy} = \Omega_{xy}\Omega_{xz}$ , so that  $\Omega_{xy}$  and  $\Omega_{xz}$  are commutative, and since they do not operate on  $x, y, z$ , may without ambiguity be written

$$x^{\omega} e^{\frac{z}{x}\Omega_{xz} + \frac{y}{x}\Omega_{xy}} S.$$

In like manner, if  $S'$  be the coefficient of  $y^{\omega}$  in the covariant, and  $S''$  (the  $P_0$  above) be the coefficient of  $z^{\omega}$ , the covariant may also be written in either of the forms

$$y^{\omega} e^{\frac{x}{y}\Omega_{yx} + \frac{z}{y}\Omega_{yz}} S',$$

$$z^{\omega} e^{\frac{y}{z}\Omega_{zy} + \frac{x}{z}\Omega_{zx}} S''.$$

Ex. 9. Prove that the covariant may also be written

$$z^{\omega} e^{\frac{x}{y}\Omega_{yx}} e^{\frac{y}{z}\Omega_{zy}} S'',$$

and in two similar forms derived from  $S$  and  $S'$ , but that since  $\Omega_{yx}$  and  $\Omega_{zy}$  are not commutative this must *not* be written

$$z^{\omega} e^{\frac{x}{y}\Omega_{yx} + \frac{y}{z}\Omega_{zy}} S''.$$

298.] Another method of obtaining the full expression for a covariant from the coefficient of the highest power of  $z$  in it is analogous to that of § 160. Substitute, in the final coefficient  $S''$  or  $P_0$  of a covariant of the ternary  $p$ -ic  $u$ ,

$u$  for  $a_p$ ,

$$\frac{1}{p} \frac{du}{dx}, \frac{1}{p} \frac{du}{dy} \text{ for } a_{p-1}, b_{p-1},$$

$$\frac{1}{p(p-1)} \frac{d^2u}{dx^2}, \frac{1}{p(p-1)} \frac{d^2u}{dxdy}, \frac{1}{p(p-1)} \frac{d^2u}{dy^2}$$

for  $a_{p-2}, b_{p-2}, c_{p-2},$

$$\frac{1}{p!} \frac{d^p u}{dx^p}, \frac{1}{p!} \frac{d^p u}{dx^{p-1} dy}, \dots \text{ for } a_0, b_0, \dots,$$

and divide through by the power of  $z$  which occurs as a factor in the result.

The proof is easy. We at once see that

$$\begin{aligned}
 u &= z^p e^{\frac{x}{z}\Omega_{zx} + \frac{y}{z}\Omega_{zy}} a_p, \\
 \frac{1}{p} \frac{du}{dx} &= \frac{1}{p} z^{p-1} e^{\frac{x}{z}\Omega_{zx} + \frac{y}{z}\Omega_{zy}} \Omega_{zx} a_p \\
 &= z^{p-1} e^{\frac{x}{z}\Omega_{zx} + \frac{y}{z}\Omega_{zy}} a_{p-1}, \\
 \frac{1}{p} \frac{du}{dy} &= z^{p-1} e^{\frac{x}{z}\Omega_{zx} + \frac{y}{z}\Omega_{zy}} b_{p-1}, \\
 \frac{1}{p(p-1)} \frac{d^2u}{dx^2} &= z^{p-2} e^{\frac{x}{z}\Omega_{zx} + \frac{y}{z}\Omega_{zy}} a_{p-2}, \\
 &\text{\&c., \&c.}
 \end{aligned}$$

Also, if  $P$  and  $Q$  be two functions of the coefficients,

$$e^{\frac{x}{z}\Omega_{zx} + \frac{y}{z}\Omega_{zy}} PQ = e^{\frac{x}{z}\Omega_{zx} + \frac{y}{z}\Omega_{zy}} P \cdot e^{\frac{x}{z}\Omega_{zx} + \frac{y}{z}\Omega_{zy}} Q;$$

for

$$\begin{aligned}
 \left(\frac{x}{z}\Omega_{zx} + \frac{y}{z}\Omega_{zy}\right)^m PQ &= P \cdot \left(\frac{x}{z}\Omega_{zx} + \frac{y}{z}\Omega_{zy}\right)^m Q \\
 + m \left(\frac{x}{z}\Omega_{zx} + \frac{y}{z}\Omega_{zy}\right) P \cdot \left(\frac{x}{z}\Omega_{zx} + \frac{y}{z}\Omega_{zy}\right)^{m-1} Q &+ \dots \\
 + \left(\frac{x}{z}\Omega_{zx} + \frac{y}{z}\Omega_{zy}\right)^m P \cdot Q, &
 \end{aligned}$$

by the method of Leibnitz's theorem. Hence the conclusion is immediate.

Since  $z^w = z^\varpi z^{w-\varpi}$ , the power of  $z$  which divides through is  $z^{w-\varpi}$ , where  $w$  is the weight of the final coefficient on the supposition that  $z$  has zero weight, i.e. of the covariant, and  $\varpi$  is the order of the latter. The difference  $w - \varpi$  is the weight, on the same supposition, of the coefficient of  $x^\varpi$ .

299.] The search for covariants of ternary quantics is then, as in the case of binary quantics, coextensive with the search for their leading or end coefficients, it being equally reasonable to consider the coefficient of  $x^\varpi$  or  $y^\varpi$  or  $z^\varpi$  a leader.

Take  $S''$ , the coefficient of  $z^\varpi$ . It has, as has been seen, the four annihilators  $\Omega_{zx}, \Omega_{yz}, \Omega_{yx}, \Omega_{xy}$ .

By § 286 (13) if  $\Omega_{yx}$  and  $\Omega_{zx}$  annihilate a function, then  $\Omega_{yz}$

must. Thus we may say that  $S''$  has the three annihilators

$$\Omega_{xz}, \Omega_{yx}, \Omega_{xy},$$

and, as a consequence, also the fourth  $\Omega_{yz}$ .

We proceed to see that any rational integral function  $S''$  with these properties is the last coefficient in a covariant, or a sum of last coefficients in more covariants than one. (This latter will be the state of things when  $S''$  is a sum of parts for which  $\varpi$  in (5) below has different values.)

Let us adopt the notation of the cubic for simplicity. That  $\Omega_{yx}$  and  $\Omega_{xy}$  are annihilators tells us that  $S''$  is a full *invariant* of the system

$$\left. \begin{aligned} a_3, \\ a_2x + b_2y, \\ a_1x^2 + 2b_1xy + c_1y^2, \\ a_0x^3 + 3b_0x^2y + 3c_0xy^2 + d_0y^3; \end{aligned} \right\} \dots (1)$$

and that  $\Omega_{xz}$  is an annihilator tells us that it is a seminvariant of the system

$$\begin{aligned} d_0, \\ c_1z + c_0x, \\ b_2z^2 + 2b_1zx + b_0x^2, \\ a_3z^3 + 3a_2z^2x + 3a_1zx^2 + a_0x^3, \end{aligned}$$

or, let us say, that it is an anti-seminvariant of the system

$$\left. \begin{aligned} d_0, \\ c_0x + c_1z, \\ b_0x^2 + 2b_1xz + b_2z^2, \\ a_0x^3 + 3a_1x^2z + 3a_2xz^2 + a_3z^3. \end{aligned} \right\} \dots (2)$$

The consequence that  $\Omega_{yz}$  is an annihilator tells us that in virtue of having these properties  $S''$  must be also an anti-seminvariant of the system

$$\left. \begin{aligned} a_0, \\ b_0y + a_1z, \\ c_0y^2 + 2b_1yz + a_2z^2, \\ d_0y^3 + 3c_1y^2z + 3b_2yz^2 + a_3z^3. \end{aligned} \right\} \dots (3)$$

Taking for  $S''$  any solution of  $\Omega_{xz}S'' = 0, \Omega_{yz}S'' = 0, \Omega_{xy}S'' = 0,$

let us form from it the function

$$e^{\frac{y}{z}\Omega_{xy} + \frac{x}{z}\Omega_{xz}} S'',$$

remembering that  $\Omega_{xy}$  and  $\Omega_{xz}$  are commutative. We proceed to see that this, made integral by the lowest necessary power  $z^\omega$  of  $z$ , is a covariant.

Because  $\Omega_{xz}S'' = 0$  we can form from  $S''$  a covariant

$$z^\omega e^{\frac{x}{z}\Omega_{xz}} S''$$

of the system (2). Call this

$$x^\omega S + \omega x^{\omega-1} z S_1 + \frac{\omega(\omega-1)}{1 \cdot 2} x^{\omega-2} z^2 S_2 + \dots \\ + \omega x z^{\omega-1} S_{\omega-1} + z^\omega S''. \quad \dots (4)$$

The order  $\omega$  here is given by

$$\omega = 3i_3 + 2i_2 + i_1 - 2w, \quad \dots (5)$$

where  $w$  is the sum of the suffixes in  $S$ , or, to express by what is known,  $w + \omega$  is the sum of the suffixes in  $S''$ , and  $i_3, i_2, i_1$  are the degrees of  $S''$  in  $a$ 's, in  $b$ 's, and in  $c$ 's respectively.  $\omega$  is non-negative by the known theory of binary quantics. If for different parts of  $S''$  this expression for  $\omega$  has different values, the present and following reasoning applies to those parts separately. By  $S''$  we now mean such a part.

Now  $S, S_1, S_2, \dots, S_{\omega-1}$  are all seminvariants of the system (1), of which  $S''$  is an invariant. For, § 286 (4), whatever be the function operated on

$$\Omega_{yx}\Omega_{zx} = \Omega_{zx}\Omega_{yx};$$

whence

$$\Omega_{yx}S_{\omega-1} = \Omega_{yx}\Omega_{zx}S'' = \Omega_{zx}\Omega_{yx}S'' = 0,$$

because  $\Omega_{yx}$  annihilates  $S''$ ; and

$$\Omega_{yx}S_{\omega-2} = \Omega_{yx} \cdot \frac{1}{2} \Omega_{zx} S_{\omega-1} = \frac{1}{2} \Omega_{zx} \Omega_{yx} S_{\omega-1} = 0,$$

&c., &c.

Again  $S, S_1, S_2, \dots, S_{\omega-1}, S''$  are all annihilated by  $\Omega_{yz}$ . That  $S''$  is so has been seen above. Also it has been seen, § 286 (12), that

$$\Omega_{yz}\Omega_{zx} - \Omega_{zx}\Omega_{yz} = -\Omega_{yx}.$$

Thus

$$\Omega_{yz}S_{\omega-1} = \Omega_{yz}\Omega_{zx}S'' = (\Omega_{zx}\Omega_{yz} - \Omega_{yx})S'' = 0,$$

because  $\Omega_{yz}$  and  $\Omega_{yx}$  annihilate  $S''$ ;

$$\Omega_{yz}S_{\varpi-2} = \Omega_{yz} \cdot \frac{1}{2} \Omega_{zx}S_{\varpi-1} = \frac{1}{2}(\Omega_{zx}\Omega_{yz} - \Omega_{yx})S_{\varpi-1} = 0,$$

because  $\Omega_{yz}$  and  $\Omega_{yx}$  annihilate  $S_{\varpi-1}$ ; and so on.

Thus  $S, S_1, S_2, \dots, S_{\varpi-1}, S''$  are all anti-seminvariants of the system (3). The first, as we shall presently see, is an invariant of the system.

All of these when operated on by  $e^{\frac{y}{z}\Omega_{zy}}$  and made integral by multiplication by just adequate powers of  $z$  will then produce covariants (invariants a particular case) of the system (3).

Now these covariants are of orders 0, 1, 2, ...  $\varpi-1$ ,  $\varpi$  respectively. For in the first place  $S''$ , an invariant of the system (1), is unaltered, except at most in sign, when interchanges are made in it equivalent to the interchange of  $x$  and  $y$  in the system (1). Thus  $S''$  is the same function of the coefficients in the system (3), but for sign at most, as of those of the system (2). It is, as we have seen, an anti-seminvariant of both systems. The covariant of the system (3) of which it is the last coefficient is then of the same order  $\varpi$  as that of the system (2) of which it is also the last coefficient, and which has been written above (4). This proves what is wanted as to  $S''$ . Now  $S_{\varpi-1}$ , which is obtained from  $S''$  by operation with  $\Omega_{zx}$ , i. e. with

$$a_0 \frac{d}{da_1} + 2a_1 \frac{d}{da_2} + 3a_2 \frac{d}{da_3} + b_0 \frac{d}{db_1} + 2b_1 \frac{d}{db_2} + c_0 \frac{d}{dc_1},$$

is of weight (sum of suffixes) one less than  $w'$  the weight of  $S''$ . Also if  $i_3', i_2', i_1'$  be the degrees of  $S''$  in the coefficients of the cubic the quadratic and the linear form of the system (3), so that

$$\begin{aligned} \varpi &= 3i_3' + 2i_2' + i_1' - 2(w' - \varpi) \\ &= 2w' - 3i_3' - 2i_2' - i_1', \end{aligned}$$

the sum  $3i_3' + 2i_2' + i_1'$  for  $S_{\varpi-1}$  or  $\Omega_{zx}S''$  is  $3i_3' + 2i_2' + i_1' - 1$ , for the operation replaces in each term one of the coefficients of the cubic by one in the quadratic, or one in the quadratic by one in the linear, or, &c. Thus the order of the covariant of (3), which  $S_{\varpi-1}$  ends, is

$$\begin{aligned} \varpi' &= 2(w' - 1) - 3i_3' - 2i_2' - i_1' + 1 \\ &= \varpi - 1. \end{aligned}$$

In like manner  $S_{\varpi-2}, S_{\varpi-3}, \dots, S_2, S_1, S$  produce covariants of (3) of orders  $\varpi-2, \varpi-3, \dots, 2, 1, 0$  respectively, the last being therefore an invariant of the set (3).

It hence follows that the covariants of the set (3) in which  $S, S_1, \dots, S_{\varpi-1}, S''$  are the last coefficients are respectively

$$S \text{ or } e^{\frac{y}{z}\Omega_{zy}} S, z e^{\frac{y}{z}\Omega_{zy}} S_1, \dots, z^{\varpi-1} e^{\frac{y}{z}\Omega_{zy}} S_{\varpi-1}, z^{\varpi} e^{\frac{y}{z}\Omega_{zy}} S''.$$

Thus the expression (4),

$$x^{\varpi} S + \varpi x^{\varpi-1} z S_1 + \frac{\varpi(\varpi-1)}{1 \cdot 2} x^{\varpi-2} z^2 S_2 + \dots + \varpi x z^{\varpi-1} S_{\varpi-1} + z^{\varpi} S'',$$

i.e.  $z^{\varpi} e^{\frac{x}{z}\Omega_{zx}} S''$ , is the part free from  $y$ , and consequently  $z^{\varpi} S''$  the part free from  $x$  and  $y$ , in an integral expression

$$z^{\varpi} e^{\frac{y}{z}\Omega_{zy}} e^{\frac{x}{z}\Omega_{zx}} S'',$$

or

$$z^{\varpi} e^{z\Omega_{zy} + \frac{x}{z}\Omega_{zx}} S'',$$

which is of the form (§ 297) of a covariant of the ternary quantic.

The notation of the cubic has been used, but the argument is general.

The expression found from  $S''$  is easily seen to obey all the conditions for a covariant. It has been constructed so as to be a function of  $x$  and covariants of the system (3), so that it is annihilated by  $\Omega_{yz} - y \frac{d}{dz}$  and  $\Omega_{zy} - z \frac{d}{dy}$ . The symmetry of its form in  $x$  and  $y$  tells us that it is also annihilated by  $\Omega_{xz} - x \frac{d}{dz}$  and  $\Omega_{zx} - z \frac{d}{dx}$ . Now annihilation by these pairs necessitates annihilation by the third pair  $\Omega_{xy} - x \frac{d}{dy}$ ,  $\Omega_{yx} - y \frac{d}{dx}$ , just as in the case (§ 289) when  $x, y, z$  did not occur. This can be seen by the properties of alternants contained in the first example following.

Ex. 10. Prove from §§ 284, 286 the five triads of facts as to alternants of which the types are

$$\begin{aligned} (\Omega_{zy} - z \frac{d}{dy}) (\Omega_{yz} - y \frac{d}{dz}) - (\Omega_{yz} - y \frac{d}{dz}) (\Omega_{zy} - z \frac{d}{dy}) \\ = H_1 - y \frac{d}{dy} + z \frac{d}{dz}, \end{aligned}$$

$$\begin{aligned}
& (\Omega_{yx} - y \frac{d}{dx}) (\Omega_{zx} - z \frac{d}{dx}) - (\Omega_{zx} - z \frac{d}{dx}) (\Omega_{yx} - y \frac{d}{dx}) = 0, \\
& (\Omega_{xy} - x \frac{d}{dy}) (\Omega_{xz} - x \frac{d}{dz}) - (\Omega_{xz} - x \frac{d}{dz}) (\Omega_{xy} - x \frac{d}{dy}) = 0, \\
& (\Omega_{xy} - x \frac{d}{dy}) (\Omega_{zx} - z \frac{d}{dx}) - (\Omega_{zx} - z \frac{d}{dx}) (\Omega_{xy} - x \frac{d}{dy}) = \Omega_{zy} - z \frac{d}{dy}, \\
& (\Omega_{yx} - y \frac{d}{dx}) (\Omega_{xz} - x \frac{d}{dz}) - (\Omega_{xz} - x \frac{d}{dz}) (\Omega_{yx} - y \frac{d}{dx}) \\
& \hspace{20em} = -\Omega_{yz} + y \frac{d}{dz}.
\end{aligned}$$

Ex. 11. Vary the argument in the preceding article so as to find the covariant whose last coefficient is  $S''$  in the form

$$z^\varpi e^{\frac{x}{y} \Omega_{yx}} e^{\frac{y}{z} \Omega_{zy}} S''.$$

Ex. 12. As in § 285 prove the fact already known from chapter iii. that throughout a covariant of degree  $i$ , order  $\varpi$ , and weight  $w$

$$ip - 3w + 2\varpi = 0,$$

weight being estimated in either of the three ways.

Ex. 13. The order of any covariant of a ternary cubic is a multiple of 3.

Ex. 14. The excess of weight over degree in the coefficient of  $z^\varpi$  in a covariant of a ternary cubic,  $z$  being of weight zero, is non-negative and even.

300.] Has a ternary quadratic any covariants? Let us examine for covariants the ternary quadratic

$$u \equiv a_0 x^2 + 2b_0 xy + c_0 y^2 + 2(a_1 x + b_1 y)z + a_2 z^2.$$

The coefficient  $S''$  of the highest power of  $z$  in any covariant is an invariant of the system

$$a_0 x^2 + 2b_0 xy + c_0 y^2,$$

$$a_1 x + b_1 y,$$

$$a_2.$$

It is consequently a rational integral function of

$$a_0 c_0 - b_0^2 \equiv \alpha, \text{ say,}$$

$$a_0 b_1^2 - 2b_0 a_1 b_1 + c_0 a_1^2 \equiv \beta, \text{ say,}$$

and

$$a_2.$$



It is also annihilated by

$$\Omega_{xz} \equiv a_2 \frac{d}{da_1} + 2a_1 \frac{d}{da_0} + b_1 \frac{d}{db_0}.$$

Now

$$\Omega_{xz} a = 2a_1 c_0 - 2b_1 b_0,$$

$$\Omega_{xz} \beta = 2a_2 (a_1 c_0 - b_1 b_0),$$

and

$$\Omega_{xz} a_2 = 0.$$

If then it be

$$f(a_2, a, \beta)$$

$$\begin{aligned} \Omega_{xz} f(a_2, a, \beta) &= \frac{df}{da_2} \Omega_{xz} a_2 + \frac{df}{da} \Omega_{xz} a + \frac{df}{d\beta} \Omega_{xz} \beta \\ &= 2(a_1 c_0 - b_1 b_0) \left( \frac{df}{da} + a_2 \frac{df}{d\beta} \right). \end{aligned}$$

Thus

$$\frac{df}{da} + a_2 \frac{df}{d\beta} = 0,$$

so that  $a$  and  $\beta$  only occur in  $f$  in the connexion  $a_2 a - \beta$ . Consequently

$$S'' = F(a_2, a_2 a_0 c_0 - a_2 b_0^2 - a_0 b_1^2 + 2b_0 a_1 b_1 - c_0 a_1^2),$$

where the second argument is the one invariant (§ 290), the discriminant  $D$ , of  $u$ .

$S''$  is then such an expression as  $a_2^m D^n$ , or a sum of such terms. It can, in fact, be only one such term. For it has to be homogeneous and isobaric, the  $z^\omega$  which it multiplies being taken as of weight zero, and these two facts give  $m + 3n = \text{constant}$  and  $2m + 2n = \text{constant}$ , so that  $m$  and  $n$  are constant.

Now the covariant

$$z^\omega e^{\frac{y}{z} \Omega_{xy} + \frac{x}{z} \Omega_{xz}} S''$$

determined from a final coefficient  $S''$  is unique. Also  $u^m D^n$  is a covariant of  $u$  with  $a_2^m D^n$  for final coefficient. There is then no covariant which is not of the form  $u^m D^n$ .

In other words, a ternary quadratic has no covariant which is not a mere product of powers of its discriminant and itself.

301.] **Covariants of the ternary cubic.** The cubic is taken as before to be

$$\begin{aligned} a_0 x^3 + 3b_0 x^2 y + 3c_0 x y^2 + d_0 y^3 + 3(a_1 x^2 + 2b_1 x y + c_1 y^2) z \\ + 3(a_2 x + b_2 y) z^2 + a_3 z^3. \end{aligned}$$

It contains ten coefficients and three variables, together thirteen. The general scheme of linear substitution contains nine constants. These, eliminated between thirteen equations expressive of the identity of old and new forms, leave four equations connecting old and new coefficients and variables.

We must be prepared then to meet with four absolute covariants and invariants of the cubic, i.e. to meet with five quite independent covariants and invariants, including the cubic itself. We have already met with four, the cubic itself, the invariants  $S$  and  $T$ , and one covariant the Hessian. Another quite independent one must be expected.

Before seeking it let us illustrate the methods of §§ 296–299 by finding the one covariant which we already know, i.e. the Hessian, of degree 3 and order 3, and consequently (§ 299, Ex. 12) of whole weight  $\frac{1}{3}(9+6) = 5$ . This must also be the weight of the coefficient of  $z^3$  in it.

We seek this coefficient of  $z^3$ , i.e. an invariant of the system

$$\left. \begin{aligned} a_0 x^3 + 3b_0 x^2 y + 3c_0 xy^2 + d_0 y^3, \\ a_1 x^2 + 2b_1 xy + c_1 y^2, \\ a_2 x + b_2 y, \\ a_3, \end{aligned} \right\} \dots (1)$$

which is annihilated by

$$\begin{aligned} \Omega_{xz} &\equiv a_3 \frac{d}{da_2} + 2a_2 \frac{d}{da_1} + 3a_1 \frac{d}{da_0} + b_2 \frac{d}{db_1} + 2b_1 \frac{d}{db_0} + c_1 \frac{d}{dc_0} \\ &\equiv a_3 \frac{d}{da_2} + \mathcal{S}, \text{ say.} \end{aligned}$$

It must involve  $a_3$ . Moreover it cannot involve  $a_3^2$ , for the weight of this exceeds 5. Let it be

$$a_3 P + Q.$$

Here  $P$  and  $Q$  must be separately invariants of the binary system above.  $P$  is an invariant of the system whose degree and weight are both 2. It must then be  $a_1 c_1 - b_1^2$ . Now, expressing the annihilation of

$$a_3(a_1 c_1 - b_1^2) + Q$$

by  $a_3 \frac{d}{da_2} + \mathfrak{S}$ , we have, by taking the terms in  $a_3^2$ ,  $a_3$ , 1 separately,

$$\frac{d}{da_2} (a_1 c_1 - b_1^2) = 0, \text{ which is obvious,}$$

$$\frac{d}{da_2} Q + \mathfrak{S} (a_1 c_1 - b_1^2) = 0,$$

$$\mathfrak{S} Q = 0;$$

of which the second gives

$$\frac{d}{da_2} Q = -2a_2 c_1 + 2b_2 a_1,$$

i. e.  $Q = -a_2^2 c_1 + 2a_2 b_2 b_1 + R,$

where  $R$  is free from  $a_2$ , and has so to be chosen that  $Q$  is an invariant of the system (1). It is made one by taking  $R = -b_2^2 a_1$ ; for which  $\mathfrak{S} Q$  is seen, as it should, to vanish.

Thus  $\{a_3(a_1 c_1 - b_1^2) - a_2^2 c_1 + 2a_2 b_2 b_1 - b_2^2 a_1\} z^3$

is the last term in a covariant, the whole expression of which is obtained by operating on the term with

$$e^{\frac{y}{z} \Omega_{zy} + \frac{x}{z} \Omega_{zx}}.$$

The coefficient of  $z^3$  is correctly

$$\begin{vmatrix} a_1, & b_1, & a_2 \\ b_1, & c_1, & b_2 \\ a_2, & b_2, & a_3 \end{vmatrix},$$

and the whole covariant is, as it should be,

$$\frac{1}{216} \begin{vmatrix} \frac{d^2 u}{dx^2}, & \frac{d^2 u}{dxdy}, & \frac{d^2 u}{dxdz} \\ \frac{d^2 u}{dxdy}, & \frac{d^2 u}{dy^2}, & \frac{d^2 u}{dydz} \\ \frac{d^2 u}{dxdz}, & \frac{d^2 u}{dydz}, & \frac{d^2 u}{dz^2} \end{vmatrix} \equiv H.$$

Ex. 15. Prove that for the canonical form  $x^3 + y^3 + z^3 + 6mxyz$  the covariants  $-Tu + 24SH$  and  $8S^2u + 3TH$  are

$$(1 + 8m^3) \{(4m^3 - 1)(x^3 + y^3 + z^3) + 18mxyz\}$$

and

$$(1 + 8m^3) \{m^2(5 + 4m^3)(x^3 + y^3 + z^3) + 3(1 - 10m^3)xyz\}. \text{ (Cayley.)}$$

302.] We have to seek another covariant of the ternary cubic, independent of  $u, S, T$  and the Hessian  $H$ .

We have seen that there is no other invariant independent of  $S$  and  $T$ . There can also be no further covariant of order 3. For, by consideration of the canonical form

$$X^3 + Y^3 + Z^3 + 6mXYZ,$$

it may be proved that any cubic covariant must be a linear function of  $X^3 + Y^3 + Z^3$  and  $XYZ$ , and consequently of  $u$  and  $H$ .

Now (§ 299, Ex. 13) the order of any covariant of the cubic is a multiple of 3. The next possible order is then 6. We proceed to see that there is a covariant of order 6 and degree in the coefficients 8, which is independent of  $u, H, S$  and  $T$ .

We have already two covariants of this order and degree; viz.  $uHS$  and  $u^2T$ . We seek a third, by looking for the coefficient in it of  $z^6$ .

By § 299, Ex. 12 the weight of the covariant is 12, which is the weight of  $uHS$  and  $u^2T$ . The weights of the coefficients of  $z^6$  are equally 12.

The highest power of  $a_3$  which can occur in the coefficient sought is then  $a_3^4$ , whose weight is 12. The coefficient of  $a_3^4$  in it must be of weight zero, so that that coefficient is a function of  $a_0, b_0, c_0, d_0$  only, and, being an invariant of

$$\left. \begin{aligned} a_0x^3 + 3b_0x^2y + 3c_0xy^2 + d_0y^3, \\ a_1x^2 + 2b_1xy + c_1y^2, \\ a_2x + b_2y, \end{aligned} \right\} \dots (1)$$

must be an invariant of the first only, and so, being of degree 4, must be a numerical multiple of

$$(\alpha_0d_0 - b_0c_0)^2 - 4(\alpha_0c_0 - b_0^2)(b_0d_0 - c_0^2).$$

Now  $a_3^4$  times this is the corresponding coefficient in  $u^2T$ . Thus, after subtracting a numerical multiple of  $u^2T$ , we have left in the coefficient of  $z^6$  no term involving  $a_3^4$ . It suffices then to look for a covariant in which the coefficient of  $z^6$  is of the form

$$a_3^3P_1 + a_3^2Q_1 + a_3R_1 + S_1.$$

We have to determine  $P_1, Q_1, R_1, S_1$ , as invariants of the system (1), in such a way that this may be annihilated by

$$\Omega_{xz}, \text{ i.e. } a_3 \frac{d}{du_2} + \mathfrak{F}.$$

As in § 301 we must have

$$\frac{d}{da_2} P_1 = 0, \quad \dots (2)$$

$$\frac{d}{da_2} Q_1 + \mathfrak{S} P_1 = 0, \quad \dots (3)$$

$$\frac{d}{da_2} R_1 + \mathfrak{S} Q_1 = 0, \quad \dots (4)$$

$$\frac{d}{da_2} S_1 + \mathfrak{S} R_1 = 0, \quad \dots (5)$$

$$\mathfrak{S} S_1 = 0. \quad \dots (6)$$

The first of these tells us that  $P_1$  does not involve  $a_2$ , and consequently is an invariant of the cubic and quadratic in the set (1) only. Its degree is 5 and its weight 3. Now (§ 260) the only invariants of the quadratic and cubic of this degree and sum of suffixes are

$$AF \equiv (a_1 c_1 - b_1^2) \{ a_1 (b_0 d_0 - c_0^2) - b_1 (a_0 d_0 - b_0 c_0) + c_1 (a_0 c_0 - b_0^2) \}$$

and (§ 261, end)

$$G \equiv a_1^3 d_0^2 + c_1^3 a_0^2 - 6 a_1^2 b_1 c_0 d_0 - 6 b_1 c_1^2 a_0 b_0 + 2(a_1 c_1 + 2b_1^2)(a_1 b_0 d_0 + c_1 a_0 c_0) + (a_1 c_1 + 8b_1^2)(a_1 c_0^2 + c_1 b_0^2) - 2a_1 b_1 c_1 a_0 d_0 - 2b_1(5a_1 c_1 + 4b_1^2) b_0 c_0;$$

and of these the first is the coefficient of  $a_3^3$ , the highest power of  $a_3$  which occurs, in the coefficient of  $z^6$  in  $uHS$ . We may subtract this covariant, and look for a covariant in which the coefficient of  $z^6$  has the form

$$a_3^3 G + a_3^2 Q + a_3 R + S.$$

With some labour, by successive use of (3), (4), (5), (6), we can determine  $Q$ ,  $R$ ,  $S$  as invariants of the system (1). For their expression we need, besides  $G$  above and  $A$  to  $F$  of § 291, the following other invariants of the system, taken from § 260 by aid of § 262,

$$K = c_1 a_2^2 - 2b_1 a_2 b_2 + a_1 b_2^2, \quad [\S 260 (1)],$$

$$L = a_0 b_2^3 - 3b_0 a_2 b_2^2 + 3c_0 a_2^2 b_2 - d_0 a_2^3, \quad [\S 260 (3)],$$

$$M = (a_1 b_0 - a_0 b_1) b_2^3 - (2a_1 c_0 - b_1 b_0 - c_1 a_0) a_2 b_2^2 + (a_1 d_0 + b_1 c_0 - 2c_1 b_0) a_2^2 b_2 - (b_1 d_0 - c_1 c_0) a_2^3, \quad [\S 260 (7)].$$

We find that

$$Q = 4AB - 3D^2 - 2FK - 8AE + 9A^3,$$

$$R = 3DL + 8AM - BK + 6EK - 11A^2K,$$

$$S = 3AK^2 - L^2 - 6KM.$$

The only point of difficulty which presents itself in proceeding by means of (3), (4), (5), (6) is the determination of the coefficient of  $A^3$  in  $Q$ . This has to be chosen so that the eventual value of  $S$  shall be annihilated by  $\mathfrak{S}$ .

The above found are not of course the only, or probably the simplest, expressions for  $Q, R, S$  in terms of invariants of the binary quadratic and cubic, of which there are five besides the ten  $A, B, \dots, L, M$ , as these are connected by many syzygies.

303.] From this coefficient of  $z^6$  in the new covariant the full expansion of the covariant, which call  $\Phi$ , may be obtained by either of the methods already detailed, i.e. by operating on it either with  $z^6 e^{\frac{y}{z}\Omega_{zy} + \frac{x}{z}\Omega_{zx}}$  or with  $z^6 e^{\frac{x}{y}\Omega_{yx}} e^{\frac{y}{z}\Omega_{zy}}$ , or by substituting in it for  $a_1, b_1, c_1, a_2, b_2, a_3$  the expressions  $\frac{1}{6} \frac{d^2u}{dx^2}$ ,  $\frac{1}{6} \frac{d^2u}{dx dy}$ ,  $\frac{1}{6} \frac{d^2u}{dy^2}$ ,  $\frac{1}{3} \frac{du}{dx}$ ,  $\frac{1}{3} \frac{du}{dy}$ ,  $u$ , and dividing through by the power of  $z$  which occurs as a factor in the result, i.e.  $z^6$ .

For the canonical form  $x^3 + y^3 + z^3 + 6mxyz$  we at once see that  $A = -m^2$ ,  $C = 1$ ,  $F = -m$ , while  $B, D, E, G, K, L, M$  all vanish, and  $a_3 = 1$ . Thus the coefficient of  $z^6$ , and therefore of  $x^6 + y^6 + z^6$ , in the covariant  $\Phi$  of the canonical form is  $-9m^6$ .

For the semi-canonized form  $ax^3 + by^3 + cz^3 + 6mxyz$  the coefficient of  $z^6$  in  $\Phi$  is in like manner  $-9c^2m^6$ , so that  $\Phi$  has the three terms  $-9m^6(a^2x^6 + b^2y^6 + c^2z^6)$ .

For the canonical form the coefficients of  $z^6$  in  $uHS$  and  $u^2T$  are  $-m^3 + m^6$  and  $1 - 20m^3 - 8m^6$  respectively. The covariant  $\Phi$  is the  $\Theta u$  of Cayley's third memoir. The  $\Theta$  of Salmon's Higher Plane Curves, § 231, has for its  $z^6$  coefficient  $3m^3 + 6m^6$ , and is  $-(\Phi + 3uSH)$ .  $\Theta$  itself might with equal reason be taken as fundamental.

To find the full expression of  $\Phi$  for the canonical form  $x^3 + y^3 + z^3 + 6mxyz$  we have to put, in the general expression

for the final coefficient found in the last article,

$$x^3 + y^3 + z^3 + 6mxyz; \quad x^2 + 2myz, \quad y^2 + 2mzx, \quad z^2 + 2mxy; \\ x, \quad mz, \quad y; \quad 1, \quad 0, \quad 0, \quad 1,$$

for  $a_3; a_2, b_2; a_1, b_1, c_1; a_0, b_0, c_0, d_0,$

respectively, and divide by  $z^6$ . The result is that for the canonical form

$$\Phi = -9m^6(x^3 + y^3 + z^3)^2 \\ - (2m + 5m^4 + 20m^7)(x^3 + y^3 + z^3)xyz \\ - (15m^2 + 78m^5 - 12m^8)x^2y^2z^2 \\ + (1 + 8m^3)^2(y^3z^3 + z^3x^3 + x^3y^3).$$

304.] The system  $u, H, S, T, \Phi$  is an algebraically complete one of invariants and covariants of the cubic. Any other covariant is a function of them. But there is another covariant which is irreducible. It was obtained by Brioschi, and is, for the semi-canonized form  $ax^3 + by^3 + cz^3 + 6mxyz,$

$$(abc + 8m^3)^3(by^3 - cz^3)(cz^3 - ax^3)(ax^3 - by^3).$$

305.] **Contravariants.** A contravariant of a ternary quantic  $u$  is (§ 66) an invariant of the system consisting of  $u$  and the linear form

$$\xi x + \eta y + \zeta z$$

in which the coefficients of the latter are present.

Now the annihilators of invariants of two ternary quantics  $u, v$  are

$$\Omega_{yx} + \Omega'_{yx}, \quad \Omega_{xy} + \Omega'_{xy}, \quad \Omega_{zy} + \Omega'_{zy}, \\ \Omega_{yz} + \Omega'_{yz}, \quad \Omega_{xz} + \Omega'_{xz}, \quad \Omega_{zx} + \Omega'_{zx},$$

where unaccented  $\Omega$ 's are the annihilators of invariants of  $u$ , and accented  $\Omega$ 's are the corresponding annihilators of invariants of  $v$ . This is proved exactly as in §§ 281, &c.

A contravariant of  $u$  has then the six annihilators

$$\Omega_{yx} + \xi \frac{d}{d\eta}, \quad \Omega_{xy} + \eta \frac{d}{d\xi}, \quad \Omega_{zy} + \eta \frac{d}{d\xi}, \\ \Omega_{yz} + \zeta \frac{d}{d\eta}, \quad \Omega_{xz} + \zeta \frac{d}{d\xi}, \quad \Omega_{zx} + \xi \frac{d}{d\zeta},$$

and all properties of contravariants, except the one fact  $ip + \omega' = \text{constant}$ , where  $\omega'$  is the order in  $\xi, \eta, \zeta$ , are consequences of these six facts of annihilation.

Notice the distinction between corresponding facts of annihilation as to covariants and contravariants. In corresponding annihilators

$$\Omega_{yx} - y \frac{d}{dx}, \quad \Omega_{yx} + \xi \frac{d}{d\eta},$$

$x$  and  $y$  correspond to  $\eta$  and  $-\xi$ , and not to  $\xi$  and  $\eta$  or  $\xi$  and  $-\eta$ .

It is not hard to see, by proceeding as in § 296, that the coefficient of  $\zeta^{\omega'}$ , the highest power of  $\zeta$  which occurs in any *contravariant*, is to be determined so as to have the four annihilators

$$\Omega_{yx}, \quad \Omega_{xy}, \quad \Omega_{zx}, \quad \Omega_{zy},$$

whereas the four annihilators of the last coefficient in a *covariant* are

$$\Omega_{yx}, \quad \Omega_{xy}, \quad \Omega_{zx}, \quad \Omega_{yz}.$$

A function of the coefficients which is annihilated by

$$\Omega_{yx}, \quad \Omega_{xy}, \quad \Omega_{zx}$$

is necessarily also annihilated by  $\Omega_{zy}$  by § 286 (10). Thus three facts of annihilation suffice for the coefficient of  $\zeta^{\omega'}$ .

It can also be seen, as in the case of covariants, that, when the final coefficient in a contravariant is known, found as any homogeneous function annihilated by  $\Omega_{yx}$ ,  $\Omega_{xy}$  and  $\Omega_{zx}$ , the whole contravariant is determined in the form

$$\zeta^{\omega'} e^{-\frac{\xi}{\zeta} \Omega_{zx} - \frac{\eta}{\zeta} \Omega_{yz}} \Sigma,$$

where  $\Sigma$  is the coefficient in question.

In a covariant the last coefficient is the one of greatest weight (sum of suffixes). In a contravariant, on the other hand, it is the one of least weight. This is reasonable, for, to make

$$\xi x + \eta y + \zeta z$$

isobaric when we take  $x, y, z$  of weights 1, 1, 0, we naturally take  $\xi, \eta, \zeta$  of weights 0, 0, 1.

Ex. 16. A ternary  $p$ -ic ( $p > 2$ ) cannot have more than, and is to be expected to have exactly,  $\frac{1}{2}(p+1)(p+2) - 5$  algebraically independent contravariants and invariants together, i.e. the same number as of algebraically independent covariants and invariants together.



306.] **Contravariant of ternary quadratic.** The method of evectants (§ 67) is a fruitful one for the discovery of contravariants.

The ternary quadratic

$$a_0x^2 + 2b_0xy + c_0y^2 + 2(a_1x + b_1y)z + a_2z^2$$

has only one contravariant. It is the evectant of the discriminant

$$a_0c_0a_2 + 2b_0a_1b_1 - a_0b_1^2 - c_0a_1^2 - b_0^2a_2,$$

i. e. is formed by operation on this with

$$\xi^2 \frac{d}{da_0} + \xi\eta \frac{d}{db_0} + \eta^2 \frac{d}{dc_0} + \xi\zeta \frac{d}{da_1} + \eta\zeta \frac{d}{db_1} + \zeta^2 \frac{d}{da_2},$$

and is

$$(c_0a_2 - b_1^2)\xi^2 + 2(a_1b_1 - b_0a_2)\xi\eta + (a_0a_2 - a_1^2)\eta^2 \\ + 2(b_0b_1 - c_0a_1)\xi\zeta + 2(b_0a_1 - a_0b_1)\eta\zeta + (a_0c_0 - b_0^2)\zeta^2.$$

Geometrically its vanishing expresses the tangential equation of the conic denoted by the quadratic, or the point-coordinate equation of a reciprocal conic. Such a contravariant has been called the *reciprocant* of a ternary quadratic. This word has been lately also used in a totally different sense of wide application.

Ex. 17. The result of substituting  $\frac{du}{dx}$ ,  $\frac{du}{dy}$ ,  $\frac{du}{dz}$  for  $\xi$ ,  $\eta$ ,  $\zeta$  in the reciprocant of a ternary quadratic  $u$  is four times the product of  $u$  and its discriminant. (*Cayley.*)

307.] **Contravariants of the ternary cubic.** The method of evectants also gives three contravariants of the cubic, in terms of which and the invariants  $S$  and  $T$  all other contravariants can be expressed, not however rationally and integrally.

We cannot expect more than three contravariants absolutely independent of one another and the invariants. For the cubic

$$a_0x^3 + 3b_0x^2y + 3c_0xy^2 + d_0y^3 + 3(a_1x^2 + 2b_1xy + c_1y^2)z \\ + 3(a_2x + b_2y)z^2 + a_3z^3$$

and

$$\xi x + \eta y + \zeta z$$

contain together thirteen coefficients, and the scheme of linear substitution contains nine. Now elimination of nine quantities

from thirteen equations leaves four only; and if there were more than five independent invariants and contravariants there would be more than four independent absolute invariants and contravariants, i. e. more than four independent results of elimination of the nine constants of substitution from the thirteen equations.

Now the first evectant of  $S$ , and the first and second evectants of  $T$ , are three independent contravariants.

We may readily form two of these three contravariants for the canonical form of the cubic.

For the semi-canonized form

$$ax^3 + by^3 + cz^3 + 6mxyz$$

the invariants are (§§ 292, 293)

$$S = m(abc - m^3),$$

$$T = (abc)^2 - 20m^3abc - 8m^6.$$

Now we know that it is not safe in general to assume that we can correctly obtain, by use of canonical or particularized forms, concomitants from other concomitants by processes which use differentiation with regard to coefficients. For, though a part of a concomitant may vanish when coefficients which vanish in the case of a particularized form are made zero, it is not as a rule the case that the derivatives of that function with regard to those coefficients vanish.

If we regard, however, the expression for  $S$  in § 291, we notice that it consists of the part  $b_0(a_0d_0a_3 - b_1^3)$ , which does not involve coefficients which vanish for the semi-canonized form, and other terms all of which are of the second or higher degrees in these coefficients. The derivatives with regard to all these coefficients will then vanish when they vanish. Moreover, if we regard the process of formation of  $T$  from  $S$  by means of the Hessian, we see that the full expression for  $T$ , too, involves, besides the terms which do not vanish for the semi-canonized form, only terms of the second and higher degrees in the coefficients which vanish for that form.

For the semi-canonized form

$$ax^3 + by^3 + cz^3 + 6mxyz$$

we consequently correctly form the first evectants of  $S$  and  $T$

by operation on the expressions above for those invariants with

$$\xi^3 \frac{d}{da} + \eta^3 \frac{d}{db} + \zeta^3 \frac{d}{dc} + \xi\eta\zeta \frac{d}{dm}.$$

Thus the first evectant of  $S$  is

$$P = m(bc\xi^3 + ca\eta^3 + ab\zeta^3) + (abc - 4m^3)\xi\eta\zeta,$$

and the first evectant of  $T$ , divided by 2, is

$$Q = (abc - 10m^3)(bc\xi^3 + ca\eta^3 + ab\zeta^3) - m^2(30abc + 24m^3)\xi\eta\zeta.$$

To obtain correctly the second evectant of  $T$  it would be necessary to retain in the full expression for  $T$ , not only the terms in  $a, b, c, m$ , but those which involve to the second degree coefficients which vanish for the semi-canonized form.

The coefficient of  $\zeta^6$  in the full expression for this second evectant of  $T$  is

$$(a_0d_0 - b_0c_0)^2 - 4(a_0c_0 - b_0^2)(b_0d_0 - c_0^2),$$

for this is the coefficient of  $a_3^2$  in  $T$ . The contravariant is then

$$\zeta^6 e^{-\frac{\xi}{\zeta}\Omega_{xz} - \frac{\eta}{\zeta}\Omega_{yz}} \{(a_0d_0 - b_0c_0)^2 - 4(a_0c_0 - b_0^2)(b_0d_0 - c_0^2)\}.$$

Another way of finding a contravariant which proves to be the same is suggested by geometry. Its vanishing is the condition that the line

$$\xi x + \eta y + \zeta z = 0$$

should touch the cubic. Thus, to find it for the semi-canonized form

$$ax^3 + by^3 + cz^3 + 6mxyz,$$

we may express that

$$\zeta^3(ax^3 + by^3) - c(\xi x + \eta y)^3 - 6m(\xi x + \eta y)\zeta^2xy,$$

considered as a binary quantic in  $x, y$ , may have a square factor, i.e. take the discriminant of this cubic in  $x, y$ . This discriminant, divided by  $\zeta^6$ , is

$$F = b^2c^2\xi^6 + c^2a^2\eta^6 + a^2b^2\zeta^6 \\ - (2abc + 32m^3)(a\eta^3\xi^3 + b\xi^3\zeta^3 + c\xi^3\eta^3) \\ - 24m^2\xi\eta\zeta(bc\xi^3 + ca\eta^3 + ab\zeta^3) - 24m(abc + 2m^3)\xi^2\eta^2\zeta^2.$$

For the fully canonized form

$$x^3 + y^3 + z^3 + 6mxyz$$

the three contravariants are

$$P = m(\xi^3 + \eta^3 + \zeta^3) + (1 - 4m^3)\xi\eta\zeta,$$

$$Q = (1 - 10m^3)(\xi^3 + \eta^3 + \zeta^3) - m^2(30 + 24m^3)\xi\eta\zeta,$$

$$F = \xi^6 + \eta^6 + \zeta^6 - (2 + 32m^3)(\eta^3\zeta^3 + \zeta^3\xi^3 + \xi^3\eta^3) \\ - 24m^2\xi\eta\zeta(\xi^3 + \eta^3 + \zeta^3) - (24m + 48m^4)\xi^2\eta^2\zeta^2.$$

$P$  is called by Cayley (who takes  $-P$ ) the Pippian, and by other writers the Cayleyan.  $Q$  is called the Quippian.  $F$  is the reciprocant.

308.] In terms of  $P$ ,  $Q$ ,  $F$  and the invariants  $S$  and  $T$  all contravariants can be expressed. There is, however, one more irreducible contravariant which is not a rational integral function of them, obtained by Hermite. For the semi-canonized form  $ax^3 + by^3 + cz^3 + 6mxyz$  its expression is

$$(abc + 8m^3)^3 (c\eta^3 - b\zeta^3)(a\zeta^3 - c\xi^3)(b\xi^3 - a\eta^3).$$

Ex. 18. Prove that  $4SQ - 3TP$  and  $TQ + 48S^2P$  are cubic contravariants whose canonical forms are

$$(1 + 8m^3)^2 \{m(\xi^3 + \eta^3 + \zeta^3) - 3\xi\eta\zeta\}, (1 + 8m^3)^2 \{(1 + 2m^3)(\xi^3 + \eta^3 + \zeta^3) \\ + 18m^2\xi\eta\zeta\}. \quad (\text{Aronhold.})$$

Ex. 19. The result of putting  $\frac{du}{dx}$ ,  $\frac{du}{dy}$ ,  $\frac{du}{dz}$  for  $\xi$ ,  $\eta$ ,  $\zeta$  in  $F$  the reciprocant of a ternary cubic  $u$  is the product of  $u$  and a covariant; and the same is true as to the reciprocant of any ternary quantic.

(Cayley.)

309.] **Mixed concomitants.** A mixed concomitant of a ternary quantic  $u$  may be regarded as a covariant of the system consisting of  $u$  and the linear form

$$\xi x + \eta y + \zeta z.$$

It has then the six annihilators

$$\Omega_{yx} + \xi \frac{d}{d\eta} - y \frac{d}{dx}, \quad \Omega_{zy} + \eta \frac{d}{d\zeta} - z \frac{d}{dy}, \quad \Omega_{zx} + \zeta \frac{d}{d\xi} - x \frac{d}{dz}, \\ \Omega_{xy} + \eta \frac{d}{d\xi} - x \frac{d}{dy}, \quad \Omega_{yz} + \zeta \frac{d}{d\eta} - y \frac{d}{dz}, \quad \Omega_{zx} + \xi \frac{d}{d\zeta} - z \frac{d}{dx}.$$

If a mixed concomitant of a ternary  $p$ -ic be of orders  $\varpi$ ,  $\varpi'$  in  $x$ ,  $y$ ,  $z$  and  $\xi$ ,  $\eta$ ,  $\zeta$  respectively, it readily follows that the terms of it in  $\zeta^{\varpi'}$  have the annihilators

$$\Omega_{yx} - y \frac{d}{dx}, \quad \Omega_{xy} - x \frac{d}{dy}, \quad \Omega_{zx} - z \frac{d}{dx}, \quad \Omega_{zy} - z \frac{d}{dy},$$

of which the first two and one of the others necessitate the fourth.

If  $P\zeta^{\varpi'}$  denote the aggregate of these terms the whole concomitant is

$$\zeta^{\varpi'} e^{-\frac{\xi}{\zeta}(\Omega_{zx} - x \frac{d}{dz}) - \frac{\eta}{\zeta}(\Omega_{yz} - y \frac{d}{dz})} P.$$

If  $Sz^{\varpi}$  be the highest term in  $z$  which occurs in  $P$ , then  $S$  has the annihilators  $\Omega_{yx}$ ,  $\Omega_{xy}$ , and is consequently an invariant of the system

$$(a_0, b_0, c_0, d_0, \dots) (x, y)^p,$$

$$(a_1, b_1, c_1, \dots) (x, y)^{p-1},$$

$$(a_2, b_2, \dots) (x, y)^{p-2},$$

$$\dots \dots \dots$$

$$a_{p-1}x + b_{p-1}y,$$

$$a_p.$$

The whole expression for  $P$  is  $z^{\varpi} e^{\frac{x}{z}\Omega_{zx} + \frac{y}{z}\Omega_{zy}} S$ .

Consequently if  $Sz^{\varpi}\zeta^{\varpi'}$  be the last term in any concomitant the whole can be derived from it, and is

$$\zeta^{\varpi'} e^{-\frac{\xi}{\zeta}(\Omega_{zx} - x \frac{d}{dz}) - \frac{\eta}{\zeta}(\Omega_{yz} - y \frac{d}{dz})} \cdot z^{\varpi} e^{\frac{x}{z}\Omega_{zx} + \frac{y}{z}\Omega_{zy}} S.$$

As to  $\varpi$  and  $\varpi'$  the former may be taken arbitrarily not below a certain limit; viz. not below  $m$  where  $m$  is the first integer for which  $(x\Omega_{xx} + y\Omega_{yy})^{m+1}S = 0$ .  $\varpi'$  is then determinate and has a constant difference from  $\varpi$ . If  $K$  is the concomitant for the lowest value  $m$  of  $\varpi$ , the concomitant for any higher value of  $\varpi$  is merely  $(\xi x + \eta y + \zeta z)^{\varpi-m} K$ . In fact the whole concomitant, which may be written

$$\zeta^{\varpi'} e^{\frac{1}{\zeta}(\xi x + \eta y) \frac{d}{dz}} \cdot z^{\varpi} e^{-\frac{\xi}{\zeta}\Omega_{zx} - \frac{\eta}{\zeta}\Omega_{yz}} e^{\frac{x}{z}\Omega_{zx} + \frac{y}{z}\Omega_{zy}} S,$$

is by Taylor's theorem

$$\zeta^{\omega'-\omega} (\xi x + \eta y + \zeta z)^{\omega} e^{-\frac{\xi}{\zeta} \Omega_{xz} - \frac{\eta}{\zeta} \Omega_{yz}} \left[ e^{\frac{x}{z} \Omega_{zx} + \frac{y}{z} \Omega_{zy}} S \right],$$

where in the square bracket  $\frac{1}{\zeta} (\xi x + \eta y + \zeta z)$  is put for  $z$ .

Any invariant of the system of binary quantics written above, i.e. any gradient annihilated by  $\Omega_{yx}$  and  $\Omega_{xy}$ , is the final coefficient in a concomitant of some kind, i.e. an invariant, covariant, contravariant, or mixed concomitant.

A valuable authority on this subject is a paper by Forsyth entitled 'Systems of Ternariants that are Algebraically Complete' (*American Journal*, Vol. XII).

Ex. 20. The number of algebraically independent concomitants, including the  $p$ -ic itself and  $\xi x + \eta y + \zeta z$ , of a ternary  $p$ -ic is

$$\frac{1}{2} (p+1) (p+2) - 2. \quad (\text{Forsyth.})$$

Ex. 21. The ternary quadratic has a mixed concomitant whose last term is  $(a_0 b_1^2 - 2 b_0 a_1 b_1 + c_0 a_1^2) z^2 \zeta^2$ ; and in terms of this, the quadratic itself, the discriminant, and  $\xi x + \eta y + \zeta z$  all concomitants whatever of the quadratic can be expressed.

Ex. 22. Any concomitant of the ternary cubic can be algebraically expressed in terms of  $\xi x + \eta y + \zeta z$  and seven concomitants whose last coefficients are functions of the results of replacing  $x, y$  by  $b_2, -a_2$  in

$$(a_0, b_0, c_0, d_0) (x, y)^3, (a_1, b_1, c_1) (x, y)^2, (a_2, b_2) (x, y), a_3$$

and their successive derivatives with regard to  $x$ . (Cf. § 263.)

(Forsyth.)

310.] The whole system of *irreducible*, pure and mixed, concomitants of the ternary cubic has been found to consist of thirty-four forms. This was established by Gordan (*Math. Ann.* I). The system was systematically exhibited by Gundelfinger (*Math. Ann.* VI); and calculated for the form  $ax^3 + by^3 + cz^3 + 6mxyz$  by Cayley (*Am. J.* IV).

311.] **Quantics in more than three variables.** With regard to  $q$ -ary quantics in general we confine ourselves to one proposition due to Sylvester.

*For a homogeneous function of the coefficients in a  $q$ -ary*

*p*-ic to be an invariant it is necessary and sufficient that it have a cyclical set of *q* annihilators of the  $\Omega$  type.

Let  $x_1, x_2, x_3, \dots, x_q$  be the variables, and denote a cyclical set of  $\Omega$ 's, whose symbolical forms as in § 282 are

$$\left[ x_2 \frac{d}{dx_1} \right], \left[ x_3 \frac{d}{dx_2} \right], \dots \left[ x_q \frac{d}{dx_{q-1}} \right], \left[ x_1 \frac{d}{dx_q} \right],$$

by  $\Omega_{2,1}, \Omega_{3,2}, \dots, \Omega_{q,q-1}, \Omega_{1,q}$ .

It can be readily proved by the theory of multiplication of determinants that the modulus of the resultant substitution which is the equivalent of a succession of substitutions is the product of their moduli. But this is not essential to the argument, in virtue of the theorem of § 23.

By § 22 a homogeneous function of the coefficients has only to be multiplied by a power of *l*, to become the same function of the coefficients in the quantic which is obtained by substituting  $lx_1, lx_2, \dots, lx_q$  for  $x_1, x_2, \dots, x_q$  in the given quantic.

By chapter vi,  $\Omega_{2,1}I = 0$  is the necessary and sufficient condition that *I* persist in form after the substitution of

$$x_1 + mx_2, x_2, x_3, \dots, x_q$$

for

$$x_1, x_2, x_3, \dots, x_q.$$

Thus  $\Omega_{2,1}I = 0$  is the necessary and sufficient condition for persistence of the homogeneous *I*, but for a power of *l*, after the substitution of

$$lx_1 + lmx_2, lx_2, lx_3, \dots, lx_q.$$

$\Omega_{3,2}I = 0$  is in like manner the necessary and sufficient condition for persistence after the further substitution of  $l'x_1, l'x_2 + m'x_3, l'x_3, \dots, l'x_q$ , but for a power of *l'*, i.e. for persistence, but for a function of the constants as factor, after the resultant substitution of

$$l'l'x_1 + lml'x_2 + lmm'x_3, l'l'x_2 + lm'x_3, l'l'x_3, \dots, l'l'x_q.$$

Repeat in like manner for  $\Omega_{4,3}I, \Omega_{5,4}I, \dots, \Omega_{q,q-1}I$ . We get eventually that there is persistence, but for a function of the constants as factor, if and only if

$$\Omega_{2,1}I = 0, \Omega_{3,2}I = 0, \dots, \Omega_{q,q-1}I = 0,$$

after substitutions of which that for  $x_1$  is

$$\lambda_1 x_1 + \lambda_2 x_2 + \dots + \lambda_q x_q,$$

where  $\lambda_1, \lambda_2, \dots, \lambda_q$  are arbitrary, each involving an  $l$  or an  $m$  not involved in any previous one of them, while the substitutions for  $x_2, x_3, \dots, x_q$  though restricted are consistent.

In like manner, if and only if

$$\Omega_{3,2} I = 0, \Omega_{4,3} I = 0, \dots, \Omega_{q,q-1} I = 0, \Omega_{1,q} I = 0,$$

there is like persistence when for  $x_2$  a general substitution

$$\mu_1 x_1 + \mu_2 x_2 + \dots + \mu_q x_q$$

is made, and for  $x_1, x_3, \dots, x_q$  restricted but consistent substitutions.

Again, similarly,

$$\Omega_{4,3} I = 0, \Omega_{5,4} I = 0, \dots, \Omega_{2,1} I = 0$$

express that there is like persistence when  $x_3$  is generally substituted for, and  $x_1, x_2, x_4, \dots, x_q$  consistently. And so on. Repeat this process  $q$  times.

Now the result of this succession of substitutions is the general substitution of

$$\lambda'_1 x_1 + \lambda'_2 x_2 + \dots + \lambda'_q x_q,$$

$$\mu'_1 x_1 + \mu'_2 x_2 + \dots + \mu'_q x_q,$$

$$\dots \dots \dots$$

$$\omega'_1 x_1 + \omega'_2 x_2 + \dots + \omega'_q x_q,$$

for  $x_1, x_2, \dots, x_q$ .

The possession of the  $q$  annihilators

$$\Omega_{2,1}, \Omega_{3,2}, \dots, \Omega_{q,q-1}, \Omega_{1,q}$$

is then necessary and sufficient for a homogeneous function  $I$  to persist in form, but for a function of the constants of substitution as factor, after the general linear substitution, i.e. for it to be an invariant of the  $q$ -ary  $p$ -ic.

In like manner for  $C$ , for which  $ip - \pi$  is constant, to be a covariant it is necessary and sufficient that  $C$  have  $q$  cyclical annihilators

$$\Omega_{2,1} - x_2 \frac{d}{dx_1}, \text{ \&c.}$$




Also for  $\Gamma$ , for which  $ip + \varpi'$  is constant, to be a contra-variant it is necessary and sufficient that it have the  $q$  cyclical annihilators

$$\Omega_{2,1} + \xi_1 \frac{d}{d\xi_2}, \text{ \&c.}$$

And for  $K$ , for which  $ip + \varpi' - \varpi$  is constant, to be a mixed concomitant it is necessary and sufficient that it have the  $q$  cyclical annihilators

$$\Omega_{2,1} + \xi_1 \frac{d}{d\xi_2} - x_2 \frac{d}{dx_1}, \text{ \&c.}$$


---



# INDEX

(The references are to pages.)

- Absolute covariants, 52.  
 A rational integral a. c. is a power of a quantic, 52.  
 A fractional one given by two covariants, 53.  
 Limit to number of independent, 55.
- Absolute invariants, 40.  
 One given by any two invariants, 40.  
 Limit to number of independent, 41.
- Algebraically complete systems,  
 for  $n$  linear forms, 339.  
 for quantics with orders in A. P., 349.
- Alternants, defined, 152.  
 of  $\Omega$  and  $O$ , 151.  
 of  $\Omega$  and  $O^r$ , 153.  
 of powers of  $\Omega$  and  $O$ , 154.  
 of  $\Sigma\Omega$  and  $(\Sigma O)^r$ , 163.  
 of ternary annihilators, 374, 377, 397.
- American Journal, 185.
- Anharmonic ratios,  
 are irrational invariants, 94, 103.  
 of roots of quartic, 292.
- Annihilation, by  $\Omega$  interpreted, 112.  
 by  $O$  interpreted, 116.
- Annihilators, of invariants, 112, 116,  
 121, 126.  
 expressed by means of roots, 237, 240.  
 by means of sums of powers of roots,  
 238, 240.  
 derive these sums in succession, 240.  
 of covariants, 128, 140.  
 of non-unitary parts of invariants,  
 224.  
 of orthogonal concomitants, 361.  
 of Boolean concomitants, 363.
- Annihilators, of all gradients, 251.  
 of ternary concomitants, 371, 372,  
 373, 405, 410.  
 of end coefficients in the same, 391,  
 406, 412.  
 of  $q$ -ary concomitants, 412.
- Anti-seminvariants, 133.
- Aronhold, 79, 112, 410.
- Aszygetic, or linearly independent,  
 invariants, 124.  
 seminvariants, 137.
- Boole, 28, 64, 353, 355, 358, 360.
- Boolean invariants, &c., 355.  
 Annihilator of, 363.
- Boolean system for linear form, 364.  
 for quadratic, 365.  
 for linear form and quadratic, 366.  
 for cubic, 367.
- Booth, 275.
- Brill, 297.
- Brioschi, 165, 405.
- Burnside and Panton, 33, 102, 195, 238.
- Canonical forms, definition of, 261.  
 of binary cubic, 263, &c.  
 of binary  $(2n-1)$ -ic, 268.  
 of quintic and septic, 271, 274.  
 of binary  $2n$ -ics, 275, 294, 299.  
 of binary quartic, 277, &c.  
 of binary sextic, 295, 297.  
 of binary octavic, 297.  
 Hammond's of quintic, 305.  
 of two quadratics, 343.  
 of ternary cubic, 300, 384, 387.

- Canonizants, 25, 27, 272.  
     with repeated factors, 270, 274.
- Canonizing equations of  $2n$ -ics, 289,  
 296, 299, 300.
- Cartesian geometry, Substitutions of,  
 354
- Catalecticants, 24, 25, 275, 276.  
     interpreted, 275.  
     of ternary &c. quantics, 301, 303.
- Cayley, 16, 38, 50, 63, 78, 99, 103, 111,  
 112, 142, 156, 162, 165, 178,  
 186, 214, 216, 218, 224, 231,  
 251, 256, 268, 285, 290, 304,  
 305, 306, 320, 322, 325, 326,  
 353, 401, 404, 407, 410, 412.
- Cayleyan, of ternary cubic, 410.
- Clebsch, 79, 80, 217, 304, 322.
- Coefficients in quantic freed from second  
 term are seminvariants, 209.
- Cogredieny, defined, 64.  
     of  $x, y$  with  $\partial_y, -\partial_x$ , 59, 88.  
     of roots with variables, 65.  
     Orthogonal the same as contragredieny,  
     358, 360.
- Combinants, 351.
- Concomitants, defined, 85.
- Contragredieny, defined, 82.  
     Geometrical, 83.  
     of  $x, y, z, \dots$  and  $\partial_x, \partial_y, \partial_z, \dots$ , 84.  
     of  $x, y$  and  $-y, x$ , 88.  
     Orthogonal the same as cogredieny,  
     358, 360.
- Contravariants, defined, 85.  
     of binary quantics not distinct from  
     covariants, 87.  
     of binary quantics found as in-  
     variants of linear form and the  
     quantics, 329, 332, 333.  
     of ternary quantics, 405.
- Convention as to numerical multiples of  
 concomitants, 90.
- Correspondence, of seminvariants and  
 non-unitaries, 245.  
     of seminvariants and power enders,  
     253.
- Covariants, defined, 4.  
     to be expected to exist, 7.  
     of several quantics from those of one,  
     28, &c.  
     of one quantic from those of several,  
     30, 75
- Covariants, of cubic quartic &c., see  
     Cubic, &c.
- Absolute, see Absolute.
- of two or more quantics, 56.
- of covariants, 57.
- derived from emanants, 71.
- of second degree in coefficients, 76,  
 79.
- as invariants, 89.
- are given by one coefficient, 131-134,  
 391.
- as functions of differences, 105.
- Sum of numerical coefficients in, 107.
- are derivable from sources by substi-  
 tutions, 207, 392.
- Covariancy of factors of a binary  
 quantic, 91.
- Cubic, Binary,  
     Canonical form of, 16, 263.  
     Cubicovariant of, 58, 63.  
     has only one invariant, 98, 173, 198,  
     228.  
     has two covariants besides itself,  
     109.  
     has no other irreducible covariant,  
     139, 175, 218.  
     has a syzygy among concomitants,  
     175, 219, 267.  
     Boolean system for, 367.
- Cubic, Ternary, Canonical form of, 300,  
 384, 387.  
     The invariant  $S$  of, 381.  
     The Hessian of, 385, 400.  
     The invariant  $T$  of, 385.  
     The irreducible system of invariants  
     of, 387.  
     The sextic covariant of, 402.  
     The contravariants of, 407.
- Cubic equation, solution of, 16, 265.
- Cubic protomorphs, 215.
- Cubicovariant of cubic, 58, 63.
- Degree, of an invariant, 32.  
     of a covariant, 47.  
     of an invariant of a binary  $(2n+1)$ -ic  
     is even, 36.  
     of a covariant of odd order of a binary  
     quantic is odd, 50.  
     of a covariant of even order of a  
     binary  $(2n+1)$ -ic is even, 50.
- Constancy of, see Homogeneity.

- Differentiation of seminvariants to produce others, 231, 332, 350.  
 Use of to obtain irreducible systems, 333.
- Diophantine Equations, 188, &c.  
 Simple sets of solutions of, 189.
- Discriminants, are invariants, 20, 17, 97.  
 freed from numerical factors, 99.  
 Weight and degree of, 99.  
 found in succession, 231.  
 of quadratics, 17.  
 of binary cubic, 58, 63.  
 of binary quartic, 102.  
 of binary quintic, 311.  
 of ternary cubic, 386.
- D'Ocagne, 256.
- Dual Substitutions, 83.
- Duality in Geometry, 83.
- Durfee, 247.
- Elemental products of differences, 194.
- Eliminants, are invariants, 17.  
 are combinants, 351.  
 of linear forms, 11.
- Emanants, 65.  
 are absolute covariants, 66.  
 Geometry of, 68.  
 Invariants of, 71.  
 Boolean invariants of, 357.
- End coefficients in ternary concomitants, 391, 393.
- Euler, 165.
- Euler's Theorem, 84, 146, 375.
- Evectants, 30, 86.
- Excess, of a gradient, 148.  
 of an invariant vanishes, 122, 152.  
 of a seminvariant is not negative, 136, 154, 163.
- Extent of a gradient, 145.
- Faà de Bruno, 21, 144, 165, 207, 247, 304, 308.
- Factor in expression of covariancy or invariancy a power of the modulus, 5, 33, 43, 47, 56, 122.  
 an integral power for rational integral concomitants, 36, 43, 49, 56.
- Factors of binary quantic, covariancy of, 91.  
 of a seminvariant are seminvariants, 144.
- Ferrers, 160.  
 Ferrers' diagrams, 160, 164, 251.
- Final coefficients in covariants have annihilator  $O$ , 133.
- Finiteness of systems of concomitants, 80, 196, 200, 201.  
 of numbers of solutions of Diophantine equations, 189, &c.
- Formes-types, 313, 316.
- Forms, defined and classified, 1.
- Forsyth, 349, 350, 412.
- Functional determinants. See Jacobians.
- Franklin, 164, 165, 181, 185.
- Generating functions, defined, 166.  
 for  $(w; i, p)$ , 166.  
 for numbers of seminvariants, 168, 170, 248, 250.  
 for concomitants of given degree and order, 178.  
 Reduced, 179, 181.  
 Representative, 181.  
 Real, 186.  
 in case of two quantics, 185.  
 for perpetuants, 249, 259.
- Generators of all seminvariants, 232, 253.
- Geometry of binary systems, 7, 68.  
 of emanants, 68.  
 of Hessians, 73.  
 of concomitants of cubic, 267.  
 of concomitants of quartic, 291.  
 of a quintic for which  $I_{18}=0$ , 318.  
 of a sextic for which  $I_{15}=0$ , 327.  
 of two quadratics, 344.  
 of Boolean concomitants, 356, 357, 365, 366, 368.  
 Metrical, chap. xv.
- Gordan, 43, 79, 80, 178, 188, &c., 304, 322.
- Gordan's Theorem, 188, &c.  
 Proof of for invariants of one binary quantic, 196.  
 for invariants and covariants, 200.  
 for more quantics than one, 201.
- Gradients, defined, 145.  
 annihilated by  $\Omega$  and  $O$  are invariants, 118.  
 of positive excess are of form  $\Omega G$ , 123, 157.  
 always of form  $\Omega G$  when extent of  $\Omega$  is infinite, 253.

- Ground forms, see Irreducible systems.  
 Gundelfinger, 412.
- Hammond, 165, 181, 184, 305, 309,  
 316, 347, 348.
- Hermite, 74, 87, 117, 160, 162, 163,  
 304, 321, 410.
- Hesse, 300.
- Hessians, are covariants, 14.  
 are discriminants of second emanants,  
 72.  
 of quadratics are discriminants, 17.  
 of cubic, Roots of, 111.  
 of quartic, Concomitants of, 288.  
 of ternary cubic, 385, 400.  
 Geometry of, 73.
- Hilbert, 43, 154, 188, &c., 232, 236.
- Hirsch, Meyer, 247.
- Homogeneity, of an invariant of one  
 quantic, 32.  
 of a complete system of invariants of  
 several quantics, 44.  
 in variables of complete systems of  
 covariants, 46, 56.  
 in coefficients of the same, 47, 56.
- Hyperdeterminants, 77, 161.
- Independent Covariants, &c., Limit to  
 number of, 53.  
 Exact number of, 210.
- Independent invariants, Limit to num-  
 ber of, 42.
- Infinite order, Binary quantic of, 246, &c.
- Intermediate invariants and covariants,  
 27, &c., 144.  
 are not combinants, 352.
- Invariants, defined, 4.  
 to be expected to exist, 6.  
 of several quantics from those of one,  
 27, &c.  
 of one quantic from those of several,  
 30, 75.  
 Homogeneity of, see Homogeneity.  
 Isobarism of, see Isobarism.  
 Absolute, see Absolute.  
 Irreducible, see Irreducible.  
 of quadratic, cubic, &c., see Quad-  
 ratic, &c.  
 Limit to number of independent, 42.  
 of two or more quantics, 43.  
 of covariants, 57.
- Invariants, of second degree, 61, 63.  
 Lineo-linear, 62.  
 as functions of differences, 93, 94, 111.  
 Sum of numerical coefficients in, 94.  
 Symmetry of, 117.  
 of odd weight are skew, 117.  
 Formation of by aid of  $\Omega$ , 124, 127.  
 Number of Aszygetic of degree  $i$ ,  
 125.  
 involve all coefficients, 149.  
 of third degree, 162.  
 of fourth degree, 162.  
 of invariants of  $\lambda u + \mu v$  are combi-  
 nants, 352.  
 of ternary quantics are defined by  
 three cyclical or two pairs of  
 annihilators, 379, 380.
- Involution, Criterion of, 25, 353.
- Irreducible concomitants are finite in  
 number, 196, 200, 201.
- Irreducible invariants, 42.  
 covariants, 55.
- Irreducible systems,  
 for linear form, 171.  
 for quadratic, 108, 171.  
 for cubic, 172, 173, 218.  
 for quartic, 176, 180, 219.  
 for quintic, 178, 184, 306.  
 for sextic, 178, 322.  
 for two linear forms, 331.  
 for linear form and quadratic, 332.  
 for linear form and cubic, 333.  
 for linear form and quartic, 338.  
 for  $n$  linear forms, 339.  
 for two quadratics, 340.  
 for linear form and two quadratics,  
 345.  
 for quadratic and cubic, 346.
- Isobarism, of invariants, 37, 39, 43,  
 375.  
 of covariants, 49, 51, 56.  
 Triple of invariants of ternary quan-  
 tics, 375.
- Jacobians, are covariants, 12.
- Joubert, 327.
- Kempe, 197, 349.
- Leading coefficients of covariants are  
 seminvariants, 132.

- Linear covariants,  
 of binary  $(2n + 1)$ -ic, 74.  
 of binary quintic, 74, 313.  
 of quadratic and cubic, 141.
- Linear form, has no invariant, 171.  
 Boolean system for, 364.
- Linear substitution, defined, 3.  
 Modulus of, 3.  
 Reversed, 33.  
 Cartesian, 354.  
 Dual, 83.
- Linear transformation, defined, 3.
- Linearly independent, see Aszygetic.
- Lineo-linear invariants, covariants and seminvariants, 62, 75, 79, 127, 142.  
 of  $u, v$  are transvectants of  $u, v$ , 142.
- MacMahon, 165, 246, 247, 248, 249, 251, 255, 256.
- Mixed concomitants, defined, 85.  
 of binary quantics found as covariants of quantics and linear form, 329, 332, 334.  
 of ternary quantics, 410.
- Modulus of linear substitution, 3.  
 must not vanish, 3.  
 irresoluble into factors, 19.  
 Powers of, 21.  
 Cartesian, 555.  
 Orthogonal, 358.
- Non-unitary symmetric functions, 245.  
 Correspondence of with seminvariants, 245.
- Non-unitary terms, determine a seminvariant, 208, 221.  
 in an invariant are given by an annihilator, 225, 227, 228, 243.
- Number of aszygetic invariants of given degree, 125.  
 of aszygetic seminvariants of given type, 137, 156, 163.  
 of seminvariants and invariants of given degree, 169.  
 of independent covariants, &c., 53, 211.  
 of independent invariants, 41.  
 of independent concomitants of ternary  $p$ -ic, 406, 412.
- Operation with one quantic on another, 60, 307.  
 with contravariants on covariants, 85.  
 with covariants on contravariants, 85.
- Operators, which effect linear transformation, 115, 121, 130.  
 which derive covariants from end coefficients, 133, 141, 392.  
 which derive ternary concomitants from end coefficients, 392, 406, 411.  
 which annihilate gradients of positive excess, 123, 158, 232.  
 which annihilate all gradients, 251.  
 which derive cubic protomorphs from quadratic, 215.  
 which derive seminvariants from non-unitary terms, 223.  
 which derive seminvariants from seminvariants, 226, 231, 232.  
 which generate seminvariants, 232, 253.  
 which derive sums of powers in succession, 240, 241.
- Order, of a quantic, 1.  
 of a covariant, 47.  
 of a covariant of a binary  $2n$ -ic is even, 50.  
 of a covariant of even degree is even, 50.  
 of a covariant of a binary  $(2n + 1)$ -ic is even or odd together with its degree, 50.
- Orthogonal, invariants, &c., 356.  
 cogredency and contragredency identical, 358, 360.  
 substitutions direct and skew, 358.
- Partitions, 125.  
 Conjugate, 159.  
 Reciprocal, 160, 164, 169.  
 Perpetuants, 249, 258, 259.  
 Pippian. See Cayleyan.
- Polar curves, 68.
- Power ending products, 250.  
 One to one correspondence of with seminvariants, 253.
- Protomorphs, defined, 212.  
 Systems of, 210, 214, 217.  
 of lowest degrees, 214.  
 for systems of quantics, 217, 332.

- Quadratic, Binary,  
   The one invariant of, 98, 172.  
   has no covariant but itself, 108, 172.  
   Boolean system for, 365.
- Quadratic, Ternary,  
   The one invariant of, 380.  
   Question of covariants of, 398.  
   Contravariant of, 407.
- Quadratic protomorphs, 214.  
   are irreducible, 221.
- Quantics, defined and classified, 1.
- Quartic, Binary,  
   Catalecticant of, 24, 63, 276.  
   Quadratic invariant of, 62.  
   The two invariants of, 101, 125.  
   Discriminant of, 102.  
   The equi-anharmonic for which  $I = 0$ ,  
     104, 282, 292.  
   The harmonic for which  $J = 0$ , 104,  
     283, 292.  
   transformed into itself, 142.  
   has only two irreducible invariants,  
     177.  
   Irreducible system and syzygy for,  
     180, 219, 287.  
   Canonical form of, 277, 285.  
   with square factor, 284.  
   Linear factor of, 290.  
   Geometry of, 291.
- Quartic, Ternary,  
   Catalecticant of, 302.  
   not in general a sum of 5 fourth  
     powers, 301.
- Quartic equation, Solution of, 289.
- Quartics, Quaternary and quinary, not  
   in general sums of 9, 14 fourth  
     powers, 303.
- Quintic, Binary,  
   Canonizant of, 25, 274.  
   Invariants of, 58, 82, 140, 307, 308.  
   Linear covariants of, 74, 82, 307, 308.  
   Generating function for invariants of,  
     177, 184.  
   Representative G. F. for, 183.  
   Canonical form of, 271, 274, 305.  
   freed from two middle coefficients,  
     305.  
   can only be freed from these and  
     another adjacent when  $I_{12} = 0$ ,  
     314.  
   List of concomitants of, 306.
- Quintic, Binary,  
   Forms of concomitants of when  $c = 0$ ,  
      $d = 0$ , 309.  
   Discriminant of, 311.  
   Syzygy among invariants of, 311.  
   expressed with invariant coefficients,  
     313.  
   for which  $I_{12} = 0$ , 314, 316, 321.  
   for which  $I_4 = 0$ , 317.  
   for which  $I_8 = 0$ , 317.  
   for which  $I_{18} = 0$ , 318, 321.  
   Skew invariant of in terms of roots,  
     320.
- Quippan, of ternary cubic, 410.
- Rational integral invariants, &c.,  
   form a complete system, 6.
- Reciprocant, of ternary quadratic, 407.  
   of ternary cubic, 409.
- Reciprocity, Hermite's law of, 160, 169,  
   249.
- Resultants. See Eliminants.
- Roberts, M., 131, 140, 239.
- Roberts, S., 256.
- Roots of binary quantic, 91.  
   Anharmonic ratios of, 94.
- Salmon, 28, 64, 78, 80, 304, 305, 308,  
   347, 404.
- Semi-canonical form of quintic, 305,  
   309.  
   of ternary cubic, 384, &c.
- Seminvariants, 132, 134.  
   lead covariants, 134, 156.  
   Formation of by aid of  $\Omega$ , 136, 141.  
   Number of aszygetic of given type,  
     137.  
   of second degree, 137.  
   of several quantics, 140.  
   as particular gradients, 146.  
   have no letters absent, 148.  
   as eliminants between  $u$  and deriva-  
     tives, 205.  
   are given by non-unitary terms, 208,  
     222, 223.  
   as integrals of  $\Omega S = 0$ , 216.  
   as quantics in  $a_p : 1$ , 229.  
   of seminvariants so regarded are  
     seminvariants, 230.  
   all generated by certain operators,  
     233, 253.



- Seminvariants, One to one correspondence of with power enders, 253.  
 derived from others by differentiation, 231, 332.  
 as invariants of a quantic and its derivatives, 349.
- Seminvariant operators, 226, 242, 257.
- Sextic, Binary,  
 Invariants of, 25, 62, 324.  
 Generating function for invariants of, 178.  
 The 26 irreducible concomitants of, 322.  
 The skew invariant of interpreted, 327.
- Skew invariant, 117.  
 of a quintic unique, 319.
- Skew orthogonal substitutions, 358.
- Solubility of quintic for which  $I_{18}=0$ , 318.
- Source of a covariant, 133.
- Stephanos, 297.
- Stroh, 249, 258, 259.
- Substitution of derivatives for variables, 64.  
 of derivatives for coefficients, 205, 207, 330, 392.
- Sufficiency of conditions of annihilation, 118, 130.
- Sum of numerical coefficients in invariants, 94.  
 in covariants and seminvariants, 107.
- Sum of powers, Condition that a  $2n$ -ic be a, 275, 301, 303.
- Sums of powers of roots, 238, &c.  
 Differentiation with regard to, 243.
- Sylvester, 64, 86, 87, 112, 144, 148, 154, 156, 161, 165, 181, 185, 231, 257, 268, 301, 304, 322, 326, 351, 353, 360, 361, 412.
- Symbolical representation, 77, 78, 79.
- Symmetry of in- and co-variants, 117, 127, 130, 140.
- Szyzgy, for binary cubic, 110, 175, 267.  
 for quartic, 181, 221.  
 among invariants of quintic, 311.
- Tamisage, 183.
- Ternary quantics, Triple arrangement of, 370.  
 Annihilators of invariants &c. of, 370, 372.  
 Covariants of, 389.
- Transvectants, 78.  
 are lineo-linear invariants of emanants, 79.  
 give all covariants and invariants, 80.  
 of a  $p$ -ic and itself, 139.  
 of two binary quantics, 142.
- Ueberschiebung, 79.
- Uniqueness of covariant led by a seminvariant, 135.
- Universal Concomitant, The, 85.
- Von Gall, 325.
- Weight, 36, 38, 48, 51, 370.  
 Constancy of, see Isobarism.

Oxford

PRINTED AT THE CLARENDON PRESS

BY HORACE HART, PRINTER TO THE UNIVERSITY

34  
0









PLEASE DO NOT REMOVE  
CARDS OR SLIPS FROM THIS POCKET

---

UNIVERSITY OF TORONTO LIBRARY

---

QA  
201  
E44

Elliott, Edwin Bailey  
An introduction to the  
algebra of quantics

Physical &  
Applied Sci.

